

Research Article

Cite this article: Houston MM, Norsworthy JK, Barber T, Brabham C (2019) Field evaluation of preemergence and postemergence herbicides for control of protoporphyrinogen oxidase-resistant Palmer amaranth (*Amaranthus palmeri* S. Watson). *Weed Technol* **33**: 610–615. doi: [10.1017/wet.2019.37](https://doi.org/10.1017/wet.2019.37)

Received: 26 February 2019

Revised: 18 April 2019

Accepted: 19 April 2019

Associate Editor:

Lawrence E. Steckel, University of Tennessee

Nomenclature:

2,4-D; atrazine; dicamba; flumioxazin; glufosinate; glyphosate; paraquat; pyriproxyfen sodium; pyroxasulfone; Palmer amaranth, *Amaranthus palmeri* S. Watson; corn, *Zea mays* L.; cotton, *Gossypium hirsutum* L.; soybean, *Glycine max* (L.) Merr

Keywords:

Multiresistance; selection pressure

Author for correspondence:

Michael M. Houston, Altheimer Laboratory, 1366 West Altheimer Dr., Fayetteville, AR 72704. Email: mmh027@uark.edu

© Weed Science Society of America, 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Field evaluation of preemergence and postemergence herbicides for control of protoporphyrinogen oxidase-resistant Palmer amaranth (*Amaranthus palmeri* S. Watson)

Michael M Houston¹, Jason K Norsworthy², Tom Barber³ and Chad Brabham⁴

¹Graduate Student, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; ²Professor, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; ³Professor, University of Arkansas, Department of Crop, Soil, and Environmental Sciences, University of Arkansas Lonoke Agricultural Center, Lonoke, AR, USA and ⁴Postdoctoral Research Associate, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA

Abstract

Palmer amaranth accessions resistant to protoporphyrinogen oxidase (PPO), 5-enolpyruvyl-shikimate-3-phosphate synthase, and acetolactate synthase (ALS)-inhibitor herbicides are widespread in the Midsouth, making control difficult. Field experiments were conducted in Marion and Crawfordsville, AR, in 2016 and 2017 to assess PRE and POST herbicides labeled for use in corn, cotton, or soybean for control of multiresistant Palmer amaranth. Accessions at both locations were resistant to glyphosate and ALS inhibitors and segregating for both the R128 and ΔG210 PPO resistance mechanisms. Of the 15 herbicide treatments tested, only atrazine (1,120 g ai ha⁻¹), pyroxasulfone (149 g ha⁻¹), and flumioxazin (144 g ha⁻¹) provided 85% or greater Palmer amaranth control 14 days after treatment (DAT). Visible control ratings at 35 DAT declined sharply, with no treatment providing more than 84% control, suggesting POST applications should be made no later than 28 DAT. Glufosinate (594 and 818 g ha⁻¹), dicamba (560 g ae ha⁻¹), 2,4-D plus glyphosate (784 g ae ha⁻¹ plus 834 g ae ha⁻¹), and paraquat (700 g ha⁻¹) applied POST to 7- to 10-cm plants reduced Palmer amaranth density 83% or more 14 DAT. Both glyphosate (1,266 g ha⁻¹) and pyriproxyfen sodium (73 g ha⁻¹) provided less than 7% Palmer amaranth control. Although flumioxazin alone at a labeled rate controlled Palmer amaranth 82% in the PRE experiment, PPO inhibitors by themselves applied POST provided no more than 37% control at 14 DAT. Effective foliar herbicides applied POST, including residual herbicides, should be made when Palmer amaranth are less than 10-cm tall for optimal control of these multiresistant Palmer amaranth accessions.

Introduction

Multiple herbicide-resistant Palmer amaranth in corn, cotton, and soybean fields is difficult to control and can drastically reduce yields if not controlled (Culpepper et al. 2010; Fast et al. 2009; Forseth et al. 1984; Klingaman and Oliver 1994; Massinga et al. 2001; Ward et al. 2013). In the United States alone, Palmer amaranth has evolved resistance to herbicides from eight site-of-action (SOA) groups: inhibitors of 5-enolpyruvyl-shikimate-3-phosphate synthase (EPSPS), acetolactate synthase (ALS), 4-hydroxyphenylpyruvate dioxygenase, photosystem II, protoporphyrinogen oxidase (PPO), synthetic auxins, very-long chain fatty acid, and microtubule assembly, and many of these accessions exhibit multiple resistance mechanisms (Heap 2019). In Arkansas and surrounding states, Palmer amaranth accessions resistant to both EPSPS- and ALS-inhibiting herbicides are common (Burgos et al. 2001; Norsworthy et al. 2008), and this widespread infestation has caused a dynamic shift in weed control programs (Hoffner et al. 2012; Neve et al. 2011). For example, after the spread of multiresistant Palmer amaranth in Georgia cotton fields, Sosnoskie and Culpepper (2014) reported a 10-fold increase in use of PPO-inhibiting herbicides, such as flumioxazin and fomesafen.

Consequently, the increased reliance on PPO inhibitors for Palmer amaranth control selected for PPO resistance. Fomesafen resistance in Palmer amaranth was initially identified in an accession collected in 2011 (Salas et al. 2016). Since then PPO-resistant Palmer amaranth accessions have been positively identified in 18 of the 29 agricultural counties in Arkansas and in three states (Arkansas, Tennessee, and Illinois) (Heap 2019; Varanasi et al. 2018). The major resistant mechanisms are the ΔG210 (210) glycine amino acid deletion, and the Arg-128-Gly/Met (R128) amino acid substitution, which are both target-site mechanisms and confer broad-spectrum resistance to PPO-inhibiting herbicides (Salas et al. 2016; Salas et al. 2017;

Varanasi et al. 2018). These mechanisms are well known, with the 210 deletion also identified in waterhemp [*Amaranthus tuberculatus* (Moq.) J.D. Sauer] and the R128/R98 mutation present in common ragweed (*Ambrosia artemisiifolia* L.). Both mechanisms are widespread in Palmer amaranth in Arkansas and have even been found in the same locations together (Varanasi et al. 2018). In fact, Varanasi et al. (2018) reported that of the 167 accessions screened in Arkansas in 2017, 28% harbored the R128 mutation and 49% harbored the 210 deletion.

PRE control of PPO-resistant Palmer amaranth can still be achieved in part with specific PPO-inhibiting herbicides. Umphres et al. (2017) concluded that flumioxazin and sulfentrazone would still provide some residual control of PPO-resistant Palmer amaranth as seen previously for waterhemp (Wuerffel et al. 2015). Even so, reliance on these two specific herbicides could lead to control failures and difficulty in achieving zero-tolerance weed control programs (Norsworthy et al. 2012; Norsworthy et al. 2014). Unlike residual PRE activity, effective control of PPO-resistant Palmer amaranth with PPO inhibitors POST is only achievable at much higher-than-labeled rates (Schwartz-Lazaro et al. 2017). Fomesafen, a commonly used POST herbicide in soybean, did not control Palmer amaranth progeny originating from Crittenden County, AR, when applied at a higher-than-labeled rate (420 g ha⁻¹) (Schwartz-Lazaro et al. 2017).

The widespread distribution of EPSPS-, ALS-, and PPO-resistant Palmer amaranth has severely limited PRE and POST options for Palmer amaranth control in soybean, corn, and cotton. Therefore, PRE and POST fallow experiments were conducted to determine how to control multiresistant Palmer amaranth

accessions harboring both the 210 and R128 PPO resistance mechanisms.

Materials and Methods

Field experiments were conducted in 2016 and 2017 near Crawfordsville (35.227029°N, 90.344904°W) and Marion, AR (35.209203°N, 90.187199°W) (Crittenden County), at on-farm sites on a Dundee silt loam soil (fine-silty, mixed, active, thermic Typic Endoaqualfs) and a Dubbs silt loam soil (fine-silty, mixed, active, thermic Typic Hapludalfs), respectively. The soil near Marion had a pH of 5.8 with 1.6% organic matter, and the soil near Crawfordsville had a pH of 5.3 with an organic matter content of 1.8%. Each site-year contained a PRE-only and a POST-only experiment to determine herbicide efficacy on multiresistant Palmer amaranth. All experiments were established into a crop-free environment. Herbicides labeled for use in soybean, cotton, and corn production were applied at the typical field-use rates listed in the herbicide index (Table 1). Visible control ratings and density reduction data were gathered in both experiments, each on a 0% to 100% scale relative to the nontreated control, with 0% being no control and 100% being complete weed mortality. Nontreated controls were included in each replication to assess relative control.

PRE Trials

PRE trials were conducted in a randomized complete block design with four replications. Plots were 3.9-m wide by 7.6-m long on 97-cm-wide beds. Herbicide treatments were applied to weed-free

Table 1. Herbicides, rates, timing, and manufacturer details for PRE and POST experiments.

Herbicide	Trade name	Rate	Manufacturer	Location	Application timing
		g ai or ae ha ⁻¹			
Imazaquin	Scepter® 70 DG	138	BASF Corporation	Research Triangle Park, NC	PRE
Pendimethalin	Prowl® H20	1,603	BASF Corporation	Research Triangle Park, NC	PRE
Dicamba	Xtedimax®	560	Monsanto Company, Inc.	St. Louis, MO	PRE
Atrazine	Aatrex® 4L	1,121	Syngenta Crop Protection, LLC	Greensboro, NC	PRE
Metribuzin	Tricore® DF	420	UPL NA Inc.	King of Prussia, PA	PRE
Diuron	Diuron 4L	841	Loveland Products, Inc.	Loveland, CO	PRE
Flumioxazin	Valor® SX	72 and 144	Valent U.S.A. Corporation	Walnut Creek, CA	PRE
Fomesafen	Reflex®	280 and 560	Syngenta Crop Protection, LLC	Greensboro, NC	PRE
Saflufenacil	Sharpen®	50 and 100	BASF Corporation	Research Triangle Park, NC	PRE
Sulfentrazone	Spartan® 4F	280 and 560	FMC Corporation	Philadelphia, PA	PRE
Acetochlor	Warrant®	1,261	Monsanto Company, Inc.	St. Louis, MO	PRE
S-metolachlor	Dual Magnum®	1,389	Syngenta Crop Protection, LLC	Greensboro, NC	PRE
Pyroxasulfone	Zidua®	149	BASF Corporation	Research Triangle Park, NC	PRE
Isoxaflutole	Balance Flexx®	88	Bayer CropScience	Research Triangle Park, NC	PRE
Mesotrione	Callisto® 480 SC	211	Syngenta Crop Protection, LLC	Greensboro, NC	PRE
Pyrithiobac sodium	Staple® LX	73	E.I. du Pont de Nemours and Co.	Wilmington, DE	POST
Dicamba	Engenia®	280 and 560	BASF Corporation	Research Triangle Park, NC	POST
Atrazine	Aatrex® 4L	1,121	Syngenta Crop Protection, LLC	Greensboro, NC	POST
Diuron	Diuron 4L	841	Loveland Products, Inc.	Loveland, CO	POST
Glyphosate	Roundup PowerMAX®	1,266	Monsanto Company, Inc.	St. Louis, MO	POST
Glufosinate	Liberty® 280 SL	594 and 818	Bayer CropScience	Research Triangle Park, NC	POST
Carfentrazone	Aim® EC	22	FMC Corporation	Philadelphia, PA	POST
Flumioxazin	Valor® SX	72	Valent U.S.A. Corporation	Walnut Creek, CA	POST
Fomesafen	Flexstar®	263 and 396	Syngenta Crop Protection, LLC	Greensboro, NC	POST
Saflufenacil	Sharpen®	25	BASF Corporation	Research Triangle Park, NC	POST
Paraquat	Gramoxone® SL 2.0	700	Syngenta Crop Protection, LLC	Greensboro, NC	POST
Mesotrione	Callisto® 480 SC	105	Syngenta Crop Protection, LLC	Greensboro, NC	POST
Tembo	Laudis®	92	Bayer CropScience	Research Triangle Park, NC	POST
TCM + tembo	Capreno®	15 + 76	Bayer CropScience	Research Triangle Park, NC	POST
2,4-D + glyphosate	Enlist Duo®	784 + 834	BASF Corporation	Research Triangle Park, NC	POST

^a Abbreviations: TCM, thienencarbazone-methyl; tembo, tembotrione.

Table 2. PRE Palmer amaranth control and density reduction averaged over experiments in Crawfordsville and Marion, AR, in 2016 and 2017.

Herbicide	Rate g ai or ae ha ⁻¹	WSSA group ^a	Density reduction ^{b,c}		Control ^b			
			28 DAT		28 DAT	35 DAT		
			%					
Imazaquin	138	2	51	c	53	d	38	ef
Pendimethalin	1,603	3	57	bc	61	cd	43	d-f
Dicamba	560	4	67	a-c	71	a-d	50	c-f
Atrazine	1,121	5	96	a	91	a	84	a
Metribuzin	420	5	88	ab	78	a-c	60	a-f
Diuron	841	7	83	a-c	76	a-c	59	a-f
Flumioxazin	72	14	89	ab	82	a-c	61	a-f
Flumioxazin	144	14	91	ab	85	ab	70	a-c
Fomesafen	280	14	68	a-c	65	b-d	43	c-f
Fomesafen	560	14	81	a-c	76	a-c	59	a-f
Saflufenacil	50	14	65	a-c	54	d	34	f
Saflufenacil	100	14	83	a-c	73	a-d	57	a-f
Sulfentrazone	280	14	78	a-c	64	b-d	50	c-f
Sulfentrazone	560	14	84	a-c	77	a-c	63	a-e
Acetochlor	1,261	15	77	a-c	73	a-d	56	b-f
S-metolachlor	1,389	15	80	a-c	73	a-d	65	a-e
Pyroxasulfone	149	15	92	ab	88	a	79	ab
Isoxaflutole	88	27	83	a-c	82	a-c	70	a-d
Mesotrione	211	27	80	a-c	71	a-d	62	a-e
P value				<0.0001		<0.0001		<0.0001

^a Abbreviations: DAT, days after treatment; WSSA, Weed Science Society of America.

^b Means within a column followed by the same lowercase letter are not different based on Tukey honestly significant difference test ($P = 0.05$).

^c Density ratings were not available for Crawfordsville, AR, in 2016.

ground that had been sprayed for winter annual weed control and tilled using standard production practices. Fifteen different herbicides were applied at standard in-crop labeled use rates (Tables 2 and 3). In addition, the four PPO-inhibiting herbicides were applied at two times the recommended rate (2X) to evaluate the effect of increased dose. A nontreated control was included for comparison. At Marion, herbicides were applied in spray solution at 112 L ha⁻¹ with a Bowman Mudmaster™ (Bowman Manufacturing, 2450 Jackson 36, Newport, AR, 72112) sprayer. At Crawfordsville, herbicides were applied in spray solution at 140 L ha⁻¹ using a CO₂-pressurized backpack sprayer. All

treatments were applied with a four-nozzle boom equipped with 110015 TeeJet® air induction extended-range nozzles at an application speed of 4.8 km h⁻¹. Density data were not available for Crawfordsville in 2016.

Because irrigation was not available at either of these sites, rainfall was the only source of water to incorporate herbicides into soil solution. Rainfall data for all experiments were recorded from in-field and local weather stations and are shown in Figures 1 and 2. At each location, a 2-cm or more rainfall event occurred within 1 to 2 weeks after herbicide treatments were applied in both years.


Table 3. Comparing sites-of-action, referenced by WSSA group number, for control of Palmer amaranth PRE in Crawfordsville and Marion, AR in 2016 and 2017.

Herbicide group contrasts ^a	Density reduction ^b		Control ^b	
	28 DAT		28 DAT	35 DAT
%				
2 vs. 14 (Low)	0.0029 (51 vs. 75)	0.0102 (53 vs. 66)	0.1430 (38 vs. 47)	
3 vs. 14 (Low)	0.0255 (57 vs. 75)	0.3014 (61 vs. 66)	0.4712 (43 vs. 47)	
4 vs. 14 (Low)	0.347 (67 vs. 75)	0.2600 (71 vs. 66)	0.6470 (50 vs. 47)	
5 & 7 vs. 14 (Low)	0.0075 (89 vs. 75)	<0.0001 (82 vs. 66)	<0.0001 (68 vs. 47)	
15 vs. 14 (Low)	0.1759 (83 vs. 75)	0.0055 (78 vs. 66)	<0.0001 (67 vs. 47)	
27 vs. 14 (Low)	0.2798 (82 vs. 75)	0.0054 (77 vs. 66)	<0.0001 (66 vs. 47)	
14 (Low vs. high) ^c	0.0422 (75 vs. 85)	<0.0001 (66 vs. 78)	<0.0001 (47 vs. 62)	

^a For contrasts, $P < 0.05$ was considered significant.

^b Means for contrasts are shown in parentheses.

^c Low is equivalent to the labeled rate of the respective Group 14 herbicides and the high is twice the labeled rate.

**Figure 1.** Rainfall data for Marion and Crawfordsville, AR, in 2016 each week after PRE treatment.




Figure 2. Rainfall data for Marion and Crawfordsville, AR, in 2017 each week after PRE treatment.

POST Trials

POST trials were conducted in a randomized complete block design with plots 2-m wide by 7.6-m long on flat, bed-absent ground. Trials at Marion contained three replications, whereas trials in Crawfordsville contained four replications each year. POST herbicide treatments were applied when Palmer amaranth reached 7 to 10 cm in height. S-metolachlor was applied at 1,064 g ha⁻¹

across each trial at the time of POST application to prevent emerging Palmer amaranth from becoming a factor in density reduction and visible control ratings. All POST herbicides were applied at labeled crop use rates and included crop oil concentrate at 1% vol/vol, except for treatments containing glufosinate and glyphosate (Table 4). POST applications were made at 140 L ha⁻¹ spray volume with a four-nozzle boom at 4.8 km h⁻¹ attached to a pressurized backpack. Treatments containing dicamba or 2,4-D were made with 110015 turbo Teejet® induction nozzles; all other treatments were applied with 110015 TeeJet® air induction extended-range nozzles.

Data from both experiments were separately subjected to an ANOVA using JMP Genomics, version 8 (SAS Institute Inc., SAS Campus Drive, Cary, North Carolina 27513). Data from both experiments were analyzed with a randomized site-year effect and randomized replications. Densities measured from each experiment were converted to a percentage of the nontreated control for the corresponding replication. Density data were not available in Marion in 2016 and Crawfordsville in 2017. Data from the nontreated control for each experiment were not included in analyses.

Before experimentation, there was particular interest in comparing the effects of application of the labeled rate (1X) and the 2X rate of PPO inhibitors and in determining the effectiveness of labeled PPO-inhibiting herbicides versus other SOAs. In addition, there was interest in comparing glufosinate, 2,4-D, and dicamba POST for potential efficacy differences when used for control of multiresistant Palmer amaranth. Thus, contrasts were used to test these specific interactions in both experiments. Data were separated using the Tukey honestly significant difference test at an α level value of 0.05.

Table 4. POST Palmer amaranth control and density reduction averaged over experiments in Crawfordsville and Marion, AR, in 2016 and 2017.

Herbicide ^{a,b,c}	Rate g ai or ae ha ⁻¹	WSSA group	Control ^d		Density reduction ^{d,e}
			7 DAT ^f	14 DAT	14 DAT
				%	
Pyriithiobac sodium	73	2	18 i	6 gh	50 b-d
Dicamba	280	4	56 d-g	62 a-c	83 ab
Dicamba	560	4	56 c-g	66 ab	86 ab
Atrazine	1,121	5	66 b-e	46 b-e	76 a-c
Diuron	841	7	80 a-c	63 a-c	92 a
Glufosinate	1,266	9	24 hi	1 h	33 cd
Glufosinate	594	10	70 a-d	48 b-e	89 ab
Glufosinate	818	10	82 ab	57 a-d	91 a
Carfentrazone	22	14	32 e-i	13 f-h	28 d
Flumioxazin	72	14	62 b-f	37 c-f	54 a-d
Fomesafen	263	14	42 f-i	19 e-h	36 cd
Fomesafen	396	14	37 e-i	15 f-h	36 cd
Saflufenacil	25	14	42 e-i	12 f-h	40 cd
Paraquat	700	22	91 a	75 a	85 a
Mesotrione	105	27	39 f-i	30 d-g	46 b-d
Tembotrione	92	27	33 e-i	17 f-h	34 d
Thiencarbazone-methyl + tembotrione	15 76	2 27	49 d-h	42 b-f	78 a-c
2,4-D + glyphosate	784 834	4 9	59 b-f	58 a-c	83 ab
P value			<0.0001	<0.0001	<0.0001

^a All treatments, excluding those containing glyphosate or glufosinate, were applied with a crop oil concentrate at 1% vol/vol.

^b Herbicide Group 4 vs. Group 10 contrast ($P < 0.05$ was considered significant; means for contrasts are shown in parentheses): Control: 7 DAT, $P < 0.0001$ (58 vs. 76) 14 DAT $P = 0.1195$ (62 vs 53). Density reduction 14 DAT, $P = 0.7795$ (85 vs 90).

^c All treatments, excluding those containing glyphosate or glufosinate, were applied with a crop oil concentrate at 1% vol/vol.

^d Means within a column followed by the same lowercase letter are not different based on Tukey honestly significant difference test ($P = 0.05$).

^e Crawfordsville, AR in 2017 and Marion, AR in 2016 density reduction data not available.

^f Abbreviations: DAT, days after treatment; WSSA, Weed Science Society of America.

Results and Discussion

PRE Trial

Timely and adequate rainfall occurred for incorporation of PRE herbicides into soil solution in both years at both locations. Precipitation was comparable across both years and locations in the first 2 weeks, averaging between 2.5 and 3 cm. Both locations also shared similar soil textures, field history, and management practices. The Marion and Crawfordsville locations were silt loam soil textures, previously in continuous soybean production, and were prepared by conventional tillage each year. The most notable difference in soil characteristics between the two locations was soil pH, but that only differed by 0.5 units. At both locations, PPO-resistant Palmer amaranth with target-site resistance mechanisms was confirmed previously (Varanasi et al. 2018).

Herbicide activity was evaluated at 28 and 35 DAT because POST herbicides typically need to be applied by 4 weeks after emergence of corn, cotton, or soybean to protect crop yields (Halford et al. 2001; Mulugeta and Boerboom 2000). Control of Palmer amaranth with the labeled rates of imazaquin, pendimethalin, fomesafen, saflufenacil, and sulfentrazone was less than 66% at 28 DAT and no more than 50% at 35 DAT (Table 2). Palmer amaranth has confirmed resistance to the SOAs of these herbicides at these locations (Heap 2019; Varanasi et al. 2018). Interestingly, flumioxazin, another PPO-inhibiting herbicide, controlled Palmer amaranth 82% at 72 g ha⁻¹ and 85% at 144 g ha⁻¹. Of the 19 herbicide treatments evaluated, only atrazine at 1,120 g ha⁻¹ (91%) and pyroxasulfone at 149 g ha⁻¹ (88%) provided greater than 85% control at 28 DAT (Table 2). In crop systems without the presence of an effective POST option for multiresistant Palmer amaranth, such as conventional soybean, these PRE herbicide treatments would not be enough to ensure extended control alone. In this experiment, emphasis was placed on comparing the efficacy of multiple PRE PPO-inhibiting herbicides at a labeled rate versus a 2X rate and comparing the results with activity at other SOAs (Table 2).

Regardless of the variation in response between individual herbicides within a group, contrasts for control at 28 and 35 DAT identified separation in only three group pairings (Table 2). Contrasts indicated that the 2X rate of Group 14 herbicides was more effective than the 1X rate, based on density reduction and visible weed control ratings at 28 DAT and control at 35 DAT (Table 2). When comparing Palmer amaranth densities and control at 28 DAT and control at 35 DAT, Group 2, 3, and 4 herbicides were never more effective than Group 14 herbicides at a labeled rate, based on contrasts. Conversely, Group 5 and 7 herbicides, collectively along with Group 15 and Group 27 herbicides, were more effective in controlling multiresistant Palmer amaranth at 28 DAT than the labeled rate of Group 14 herbicides. The ineffectiveness of the Groups 2 and 3 SOAs is attributed to herbicide resistance within these populations, whereas the lack of effectiveness of dicamba (Group 4) is its low adsorption coefficient and the occurrence of rainfall soon after application in both years, likely moving the herbicide well below the depth that Palmer amaranth germination and emergence occurs (Figures 1 and 2).

POST Trial


Visible control ratings for Palmer amaranth 7- to 10-cm tall at application ranged from 18% to 91% at 7 DAT (Table 3). Control with paraquat (91%) was equal to control with diuron

at 841 g ha⁻¹ (80%) and both rates of glufosinate (70% and 82%, respectively). Conversely, control with glyphosate, pyriithiobac sodium, tembotrione, or carfentrazone did not exceed 32%. As expected, contrasts between auxin herbicides and glufosinate at 7 DAT indicated glufosinate provided greater initial control. More rapid control of Palmer amaranth with glufosinate than with dicamba and 2,4-D at 7 DAT was not surprising, because glufosinate is a contact herbicide and the other two are systemic (Coetzer and Al-Khatib 2001; Grossman 2010).

At the 14 DAT timing, no treatment provided greater than 75% control (Table 3). Glyphosate and pyriithiobac sodium at 14 DAT provided less than 7% Palmer amaranth control. The lack of control from treatments evaluated infer that reliance on POST herbicides alone for control of these resistant Palmer amaranth accessions is not an acceptable weed control program. No single application of an herbicide, including dicamba, glufosinate, and 2,4-D, provided 85% control, demonstrating their lack of viability as stand-alone options. Density reduction data at 14 DAT followed trends similar to those of visible control data (Table 3). Contrast at 14 DAT between auxin herbicides and glufosinate revealed the options were comparable for controlling these Palmer amaranth accessions ($P = 0.1195$). Although efficacy of dicamba and 2,4-D at 14 DAT was evident, Palmer amaranth control with these herbicides would likely have increased with continued ratings at 21 and 28 DAT.

Practical Implications

Suppression of multiresistant Palmer amaranth with Group 5, 7, 15, and 27 herbicides is possible at 28 DAT, but not as stand-alone options. Although flumioxazin still provided good control of PPO-resistant Palmer amaranth accessions, applying flumioxazin alone could result in escapes and high selection pressure for additional PPO-inhibitor resistance. Because of lack of control of multiresistant Palmer amaranth with ALS- and PPO-inhibiting herbicides, earlier POST applications may be necessary to ensure that herbicides are applied in a timely manner to small weeds. Even when using the most effective PRE herbicide tested, it was not possible to maintain a high level of Palmer amaranth control through 35 DAT. For this reason, POST herbicides should be applied no later than 28 days after PRE application and residuals should be overlapped to provide season-long control (Aulakh and Jhala 2015; Halford et al. 2001). Overlapping effective residuals is key for Palmer amaranth control, demonstrated by the lack of control options in the POST experiment. ALS-, EPSPS-, and PPO-inhibiting herbicides should not be relied on POST in areas with confirmed or suspected Palmer amaranth resistance to these SOAs, because poor weed control will result. Even though S-metolachlor was applied PRE to all plots in the POST experiment, one application of any POST herbicide was not found to be efficacious in controlling these Palmer amaranth accessions. Based on these data, an optimum herbicide program will likely require multiple effective residuals at planting followed by two applications of an effective herbicide POST. In addition to using multiple SOA for weed control, it will also delay the onset of herbicide resistance. Control of multiresistant Palmer amaranth is not possible with ALS-, EPSPS-, or PPO-inhibiting herbicides alone.

Author ORCIDs. Michael M Houstonl,  <https://orcid.org/0000-0001-8968-9892>

Acknowledgements. Funding for this research was provided by the Arkansas Soybean Promotion Board. No conflicts of interest have been declared.

References

- Aulakh JS, Jhala AJ (2015) Comparison of glufosinate-based herbicide programs for broad-spectrum weed control in glufosinate-resistant soybean. *Weed Technol* 29:419–430
- Burgos NR, Kuk YI, Talbert RE (2001) *Amaranthus palmeri* resistance and differential tolerance to *Amaranthus palmeri* and *Amaranthus hybridus* to ALS-inhibitor herbicides. *Pest Manag Sci* 57:449–457
- Coetzer E, Al-Khatib K (2001) Photosynthetic inhibition and ammonium accumulation in Palmer amaranth after glufosinate application. *Weed Sci* 49:454–459
- Culpepper AS, Webster TM, Sosnoskie LM, York AC (2010) Glyphosate-resistant Palmer amaranth in the US. Pages 195–212 in Nandula VK, ed. *Glyphosate Resistance: Evolution, Mechanisms, and Management*. Hoboken, NJ: J. Wiley
- Fast BJ, Murdock SW, Farris RL, Willis JB, Murray DS (2009) Critical timing of Palmer amaranth (*Amaranthus palmeri*) removal in second-generation glyphosate-resistant cotton. *J Cotton Sci* 13:32–36
- Forseth IN, Ehleringer JR, Werk KS, Cook CS (1984) Field water relations of Sonoran Desert annuals. *Ecology* 65:1436–1444
- Grossman K (2010) Auxin herbicides: current status of mechanism and mode of action. *Pest Manag Sci* 66:113–120
- Halford C, Hamill AS, Zhang J, Doucet C (2001) Critical period of weed control in no-till soybean (*Glycine max*) and corn (*Zea mays*). *Weed Technol* 15:737–744
- Heap IM (2019) International survey of herbicide resistant weeds. <http://www.weedscience.org/in.asp> Accessed: April 9, 2019
- Hoffner AE, Jordan DL, Chandi A, York AC, Dunphy EJ, Everman WJ (2012) Management of Palmer amaranth (*Amaranthus palmeri*) in glufosinate-resistant soybean (*Glycine max*) with sequential applications of herbicides. *ISRN Agron* 2012:131650, [10.5402/2012/131650](https://doi.org/10.5402/2012/131650)
- Klingaman TE, Oliver LR (1994) Palmer amaranth (*Amaranthus palmeri*) interference in soybean (*Glycine max*). *Weed Sci* 42:523–527
- Massinga RA, Currie RS, Horak MJ, Boyer J (2001) Interference of Palmer amaranth in corn. *Weed Sci* 49:202–208
- Mulugeta D, Boerboom CM (2000) Critical time of weed removal in glyphosate-resistant *Glycine max*. *Weed Sci* 48:35–42
- Neve P, Norsworthy JK, Smith KL, Zelaya IA (2011) Modeling glyphosate resistance management strategies for Palmer amaranth (*Amaranthus palmeri*) in cotton. *Weed Technol* 25:335–343
- Norsworthy JK, Griffith GM, Scott RC, Smith KL, Oliver LR (2008) Confirmation and control of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) in Arkansas. *Weed Technol* 22:108–113
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60(SP 1):31–62
- Norsworthy JK, Griffith GM, Griffin T, Bagavathiannan M, Gbur EE (2014) In-field movement of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) and its impact on cotton lint yield: evidence supporting a zero-threshold strategy. *Weed Sci* 62:237–249
- Salas RA, Burgos NR, Tranel PJ, Singh S, Glasgow L, Scott RC, Nichols RL (2016) Resistance to PPO-inhibiting herbicide in Palmer amaranth from Arkansas. *Pest Manag Sci* 72:864–869
- Salas RA, Burgos NR, Rangani G, Singh S, Refatti JP, Piveta L, Tranel PJ, Mauroumoustakos A, Scott R (2017) Frequency of gly-210 deletion mutation among protoporphyrinogen oxidase inhibitor-resistant Palmer amaranth (*Amaranthus palmeri*) populations. *Weed Sci* 65:718–731
- Schwartz-Lazaro LM, Norsworthy JK, Scott RC, Barber LT (2017) Resistance of two Arkansas Palmer amaranth populations to multiple herbicide sites of action. *Crop Prot* 96:158–163
- Sosnoskie LM, Culpepper AS (2014) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) increases herbicide use, tillage, and hand-weeding in Georgia cotton. *Weed Sci* 62:393–402
- Umphres AM, Steckel LE, Mueller TC (2017) Control of protoporphyrinogen oxidase inhibiting herbicide resistant and susceptible Palmer amaranth (*Amaranthus palmeri*) with soil-applied protoporphyrinogen oxidase-inhibiting herbicides. *Weed Technol* 32:95–100
- Varanasi VK, Brabham C, Norsworthy JK, Nie H, Young BG, Houston MM, Barber LT, Scott RC (2018) A statewide survey of PPO-inhibitor resistance and the prevalent target-site mechanisms in Palmer amaranth (*Amaranthus palmeri*) accessions from Arkansas. *Weed Sci* 66:149–158
- Ward SM, Webster TM, Steckel LE (2013) Palmer amaranth (*Amaranthus palmeri*): a review. *Weed Technol* 27:12–27
- Wuerffel RJ, Young JM, Tranel PJ, Young BG (2015) Soil-residual protoporphyrinogen oxidase-inhibiting herbicides influence the frequency of associated resistance in waterhemp (*Amaranthus tuberculatus*). *Weed Sci* 63:529–538