

Psychosis/Antipsychotics. Antipsychotic intrauterine exposure is not significantly associated with increased risk of major congenital malformations. Minimum effective doses are recommended.

Case reports. (1)Pre-pregnancy counselling. Schizoaffective disorder receiving perphenazine, quetiapine and lithium. (2)First-trimester pregnancy. Discontinuation of treatment in major depressive disorder. (3-4)Second/third trimester. Occurrence of anxiety symptoms in posttraumatic stress disorder. (5)Postpartum/Breastfeeding. Discontinuation of antidepressants.

Conclusions: Shared decision-making models for antidepressants and antipsychotics prescription represent patient-centered approaches to be recommended in perinatal period.

Disclosure of Interest: None Declared

EPV1104

Postmenopausal women's perceptions regarding menopause within the context of cognitive behavioral model: A qualitative evaluation

F. Atkan* and F. Oflaz

Psychiatric Nursing, Koç University, Istanbul, Türkiye

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2394

Introduction: Women experience some physiological, cognitive and emotional symptoms due to changes during menopause, which is a critical period in their lives. Every woman experience differences in symptoms and severity of them. Women's individual perceptions and attitudes towards menopause affect their quality of life.

Objectives: The aim of this study is to describe women's perceptions and feelings about menopausal process and symptoms experienced and their behaviours to cope with these symptoms within framework of Cognitive Behavioral Model.

Methods: Phenomenology design was used in this study. Three focus group interviews were held June-September 2022, using Zoom platform. Purposive sampling was used as the sampling method of study. A total of 13 women in a menopausal period of 1-15 years, 5-6 people in each group, were included in the study. Focus group interviews were held for 1-1.5 hours and once with each group. In the focus group interviews, a 'Semi-Structured Interview Form' consisting of 4 questions was used to evaluate perceptions, changes and behaviors of postmenopausal women regarding menopause process. Zoom recordings were taken during the interviews and data were written down. The data were evaluated by thematic analysis method within framework of Cognitive Behavioral Model.

Results: In this study, 11 themes were defined as women's perception of menopause at individual and social level. These themes are negative automatic beliefs about female identity such as menopause reduces woman and loses her femininity, woman is not understood and struggles alone, woman has been stigmatized, women need support, hidden and spoken in a low voice, long and difficult process, also beliefs that physiological changes occur in the body associated with the fact that it's a natural process and that it's necessary to use new coping strategies to effectively manage these changes, onset of diseases, changing relationships, and high self-observation process. Emotions such as irritability, tension, sadness, and hypersensitivity are negative automatic beliefs that decrease

woman and loss of femininity, accompanied by emotions such as irritability, sadness, and behaviors such as crying crises, social withdrawal, that woman is not understood and is a process that she struggles alone, avoidance and social withdrawal behaviors such as reading books, listening music, walking alone.

Conclusions: In this study, it was observed that women had difficulties with gender identity along with physiological and psychological changes during menopause, and there was an increase in self-observation. It was determined that they used some new behavioral and psychological strategies to cope with this new situation. It was evaluated that these changes in emotions, thoughts and behaviors could be well formulated within the cognitive behavioral model and this model would be useful in supporting women.

Disclosure of Interest: None Declared

EPV1105

Mental Health problems and psychotherapy in female victims of human trafficking

F. S. Ali

Psychiatry, BorgElArab Central Hospital, Alexandria, Egypt

doi: 10.1192/j.eurpsy.2023.2395

Introduction: Human trafficking is the trade of humans for the purpose of forced labor, sexual slavery, or commercial sexual exploitation for the trafficker or others. This may encompass providing a spouse in the context of forced marriage or the extraction of organs.

Research into the mental health impact of trafficking has consistently found high rates of mental health problems, most commonly depression, anxiety, and posttraumatic stress disorder (PTSD).

While there is some evidence that these factors can contribute the diagnosis of PTSD, exposure to trauma is the most important feature in the development of PTSD.

Other Mental Health Problems in victims of Trafficking:

1. Dissociative disorders
2. Substance-related disorders
3. Complex trauma

Objectives: Although mental health problems among victims of trafficking have been shown to be high, recovery without treatment is rare, particularly in those who have developed PTSD.

Where there is comorbidity, recovery often does not occur even when rehabilitation has been attempted. This is unsurprising given the multiplicity of trauma that victims of trafficking have experienced, which often includes trauma prior to the trafficking situation.

Methods: Evidence-Based Therapeutic Treatment Options for PTSD:

Some evidence suggests that selective serotonin reuptake inhibitors can effectively complement the psychotherapeutic treatment of PTSD as well as other anxiety and mood disorders.

Cognitive-behavioral therapy

Exposure therapy

Eye movement desensitization and reprocessing

Narrative Exposure Therapy (NET)

Results: In the absence of research pertaining to the mental health treatment of victims of human trafficking, mental health