



Asymptotic estimate of solutions in a 4th-order parabolic equation with the Frobenius norm of a Hessian matrix

Ke Li, Bingchen Liu , and Jiaxin Dou

Abstract. This paper deals with a 4th-order parabolic equation involving the Frobenius norm of a Hessian matrix, subject to the Neumann boundary conditions. Some threshold results for blow-up or global or extinction solutions are obtained through classifying the initial energy and the Nehari energy. The bounds of blow-up time, decay estimates, and extinction rates are studied, respectively.

1 Introduction

In this paper, we study the following 4th-order parabolic problem involving the Frobenius norm of a Hessian matrix:

$$(1.1) \quad \begin{cases} u_t - \Delta u + \Delta^2 u - 2|\Delta u|^2 + 2|D^2 u|^2 = |u|^{p-1}u, & (x, t) \in \Omega \times (0, T), \\ u = 0, \quad \frac{\partial u}{\partial \eta} = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = u_0(x), & x \in \Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ ($1 \leq N \leq 3$) is a general bounded domain with smooth boundary, η is the unit outward normal vector on $\partial\Omega$, the initial datum $u_0 \in H_0^2(\Omega)$, the exponent p is a positive constant, T is the maximal existence time of (1.1), and the Frobenius norm of the Hessian matrix is defined as

$$|D^2 u| := \left[\sum_{i,j=1}^3 \left(\frac{\partial^2 u}{\partial x_i \partial x_j} \right)^2 \right]^{\frac{1}{2}}.$$

By direct computation, problem (1.1) can be rewritten as

$$(1.2) \quad \begin{cases} u_t - \Delta u + \Delta^2 u - \operatorname{div}(2\Delta u \nabla u) + \Delta|\nabla u|^2 = |u|^{p-1}u, & (x, t) \in \Omega \times (0, T), \\ u = 0, \quad \frac{\partial u}{\partial \eta} = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = u_0(x), & x \in \Omega. \end{cases}$$

Problem (1.1) or (1.2) could be used to describe the growth of thin surfaces when exposed to molecular beam epitaxy (see [13, 17, 9, 5]). In particular, u can either

Received by the editors February 5, 2024; revised August 9, 2024; accepted August 13, 2024.

AMS subject classification: 35K25, 35A01, 35B40, 35B44.

Keywords: 4th-order parabolic equation, Hessian matrix, global existence, blow-up time, extinction rate.



represent the absolute thickness of the film or rather the relative surface height – that is, the deviation of the film height at the point x from the mean film thickness at time t (see [14]); $-\Delta u$ indicates the diffusion due to evaporation-condensation (see [12]); $\Delta^2 u$ indicates capillarity-driven surface diffusion (see [12]); $\Delta|\nabla u|^2 - \operatorname{div}(2\Delta u \nabla u)$ is related to the equilibration of the inhomogeneous concentration of the diffusing particles on the surface (see [1]); the source term u^p denotes the mean deposition flux of the superlinear growth conditions with respect to u at ∞ , which could lead to the singularity of solutions or their derivatives to (1.1) or (1.2).

The parabolic equation in (1.1) or (1.2) is a typical equation of the continuum model of motion for the evolution of the film surface height $u(x, t)$:

$$u_t + A_1 \Delta u + A_2 \Delta^2 u + A_3 \operatorname{div}(|\nabla u|^2 \nabla u) + A_4 \Delta|\nabla u|^2 = f + \eta, \quad x \in \Omega, \quad t > 0,$$

where f is the deposition flux and η is the Gaussian random variable which describes the fluctuations in the average deposition flux.

In the case $A_1, A_2 > 0, A_3 < 0$ and $A_4 = 0$, Kohn and Yan in [10] considered

$$u_t + \Delta^2 u + \operatorname{div}(2(1 - |\nabla u|^2) \nabla u) = 0, \quad x \in \Omega, \quad t > 0,$$

where $\Omega \subset \mathbb{R}^2$ is a square domain. They obtained the decay of energy in time.

In the case $A_1, A_3 < 0, A_2 > 0$, and $A_4 = 0$, Liu and Li in [11] studied

$$u_t - \Delta u + \Delta^2 u - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = f(u), \quad x \in \Omega, \quad t > 0,$$

where $\Omega \subset \mathbb{R}^N$ ($N \geq 1$) is a square domain. They obtain the sufficient conditions on the global existence, asymptotic behavior, and finite time blow-up of weak solutions, but also show exact descriptions of smallness conditions on the initial data.

In the case $A_1, A_2, A_4 > 0$, and $A_3 = 0$, Winkler in [14] investigated the following equation by using the computational methods

$$(1.3) \quad u_t + \mu \Delta u + \Delta^2 u + \lambda \Delta|\nabla u|^2 = f(x), \quad x \in \Omega, \quad t > 0,$$

where $\Omega \subset \mathbb{R}^N$ ($1 \leq N \leq 3$) is a bounded convex domain with smooth boundary, $\mu \geq 0$ and $\lambda > 0$. Under appropriate assumptions on f , the global existence of weak solutions was obtained. Under an additional smallness condition on μ and the size of f , it was shown that there exists a bounded set which is absorbing for (1.3) in some sense for any solution. Blomker and Gugg in [3] also studied the related problem

$$u_t + A_1 \Delta u + \Delta^2 u + \Delta|\nabla u|^2 = \eta, \quad x \in \Omega, \quad t > 0,$$

where $\Omega \subset \mathbb{R}$ is a bounded interval. The global existence of weak solutions was proved. This result was extended by Blomker et al. in [4] to the parabolic equation

$$u_t + A_1 \Delta u + A_2 \Delta^2 u + A_4 \Delta|\nabla u|^2 = \nu|\nabla u|^2 + \eta, \quad x \in \Omega, \quad t > 0,$$

where $\Omega \subset \mathbb{R}$ is a bounded interval and $\nu > 0$.

In the case $A_1, A_2, A_4 > 0$ and $A_3 < 0$, Agélas in [1] dealt with

$$u_t + \nu \Delta u + \nu_2 \Delta^2 u - \nu_3 \operatorname{div}(|\nabla u|^2 \nabla u) + \nu_4 \Delta|\nabla u|^2 = \nu_5 |\nabla u|^2, \quad x \in \Omega, \quad t > 0,$$

where $\Omega = \mathbb{R}^N$ ($N = 1, 2$). He proved the existence, uniqueness, and regularity of global weak solutions. Moreover, under the condition $v_2 v_3 > v_4^2$, the author proved the existence and uniqueness of global strong solutions for sufficiently smooth initial data.

In [6], Escudero dealt with both the initial and initial-boundary value problems for the partial differential equation $u_t + \Delta^2 u = \det(D^2 u)$ posed either on \mathbb{R}^2 or on a bounded subset of the plane, where $\det(D^2 u)$ is the determinant of the Hessian matrix $D^2 u$. The author studied the blow-up behavior including the complete blow-up in either finite or infinite time. Moreover, he refined a blow-up criterium that was proved for this evolution equation. The interested authors could find other results in [16, 18] and the papers cited therein.

To our knowledge, the 4th-order parabolic problem (1.1) involving a Frobenius type nonlinearity has been rarely considered before. Moreover, the mean deposition flux of the superlinear growth conditions would play an important role in the property of the solutions, including the existence of blow-up, extinction solutions. Inspired by the works [14, 3], we want to study the threshold results on the initial data with respect to the existence of blow-up, global, and extinction solutions of (1.1) or (1.2). Throughout this paper, we denote by $\|\cdot\|_p$ the $L^p(\Omega)$ norm and by (\cdot, \cdot) the inner product in $L^2(\Omega)$, respectively. For $u \in H_0^2(\Omega)$ with norm $\|u\|_{H_0^2(\Omega)} = \|\Delta u\|_2$, we define the energy functional and the Nehari functional, respectively,

$$(1.4) \quad \begin{aligned} J(u) &:= \frac{1}{2} \|\nabla u\|_2^2 + \frac{1}{2} \|\Delta u\|_2^2 + \int_{\Omega} |\nabla u|^2 \Delta u \, dx - \frac{1}{p+1} \|u\|_{p+1}^{p+1}, \\ I(u) &:= \|\nabla u\|_2^2 + \|\Delta u\|_2^2 + 3 \int_{\Omega} |\nabla u|^2 \Delta u \, dx - \|u\|_{p+1}^{p+1}, \end{aligned}$$

which satisfy

$$J(u) := \frac{1}{3} I(u) + \frac{1}{6} \|\nabla u\|_2^2 + \frac{1}{6} \|\Delta u\|_2^2 + \frac{p-2}{3(p+1)} \|u\|_{p+1}^{p+1}.$$

We give the weak solutions of problem (1.1) as follows.

Definition 1.1 Let $T > 0$. A function $u(x, t) \in L^\infty(0, T; H_0^2(\Omega))$ with $u_t \in L^2(0, T; L^2(\Omega))$ is the so-called weak solution to (1.1) or (1.2) in $\Omega \times [0, T)$, if $u(x, 0) = u_0(x) \in H_0^2(\Omega)$, for any $\varphi(x) \in H_0^2(\Omega)$,

$$(1.5) \quad \begin{aligned} (u_t, \varphi) + (\nabla u, \nabla \varphi) + (\Delta u, \Delta \varphi) - (2|\Delta u|^2, \varphi) + (2|D^2 u|^2, \varphi) &= (|u|^{p-1} u, \varphi), \\ \text{or } (u_t, \varphi) + (\nabla u, \nabla \varphi) + (\Delta u, \Delta \varphi) + (2\Delta u \nabla u, \nabla \varphi) + (|\nabla u|^2, \Delta \varphi) &= (|u|^{p-1} u, \varphi). \end{aligned}$$

Moreover, there is the relationship for the energy of the weak solutions,

$$(1.6) \quad \int_0^t \|u_\tau\|_2^2 \, d\tau + J(u) = J(u_0) \quad \text{for a.e. } t \in (0, T).$$

Table 1: Complete classification of initial energy.

Initial energy	$J(u_0) - d$	$I(u_0)$	p	$\ u_0\ _2$	Solution	Main Theorems
Subcritical	-	+	$p > 2$		G.E.	Theorem 3.1
Subcritical	$-, J(u_0) \neq 0$	-	$p > 2$		B.U.	Theorem 3.2
Subcritical	$J(u_0) < 0$		$p > 2$		B.U.	Theorem 6.1
Subcritical	$J(u_0) \leq 0$		$p < 1$	$\ u_0\ _2^{1-p} \geq D_4 D_3^{-1}$	N.E.	Theorem 7.2
Subcritical	$J(u_0) \leq 0$		$p = 1$ or $p \geq 2$		N.E.	Theorem 7.2
Critical	0	+ or 0	$p > 2$		G.E.	Theorem 4.1
Critical	0	-	$p > 2$		B.U.	Theorem 4.2
Supercritical	+	+	$p > 2$	$\ u_0\ _2 \leq \lambda_{J(u_0)}$	G.E.	Theorem 5.1(i)
Supercritical	+	0	$p < 2, N = 1$	$\ u_0\ _2^{1-p} \geq D_2 D_1^{-1}$	E.	Corollary 7.2
Supercritical	+	-	$p > 2$	$\ u_0\ _2 \geq \Lambda_{J(u_0)}$	B.U.	Theorem 5.1(ii)

Define the Nehari manifold $\mathcal{N} := \{u \in H_0^2(\Omega) \mid I(u) = 0, \|\Delta u\|_2 \neq 0\}$. The potential well and its corresponding sets are defined by

$$\mathcal{W} := \{u \in H_0^2(\Omega) \mid I(u) > 0, J(u) < d\} \cup \{0\}, \quad \mathcal{V} := \{u \in H_0^2(\Omega) \mid I(u) < 0, J(u) < d\},$$

$$\mathcal{N}_+ := \{u \in H_0^2(\Omega) \mid I(u) > 0\}, \quad \mathcal{N}_- := \{u \in H_0^2(\Omega) \mid I(u) < 0\},$$

where $d := \inf_{u \in H_0^2(\Omega) \setminus \{0\}} \sup_{\lambda \geq 0} J(\lambda u) = \inf_{u \in \mathcal{N}} J(u)$ is the so-called depth of the potential well \mathcal{W} .

For any $\delta > 0$, we further define the modified functional and the Nehari manifold as

$$I_\delta(u) := \delta \|\Delta u\|_2^2 + \|\nabla u\|_2^2 + 3 \int_\Omega |\nabla u|^2 \Delta u dx - \|u\|_{p+1}^{p+1},$$

$$\mathcal{N}_\delta := \{u \in H_0^2(\Omega) \mid I_\delta(u) = 0, \|\Delta u\|_2 \neq 0\}.$$

The modified potential wells and their corresponding sets are defined respectively by

$$\mathcal{W}_\delta := \{u \in H_0^2(\Omega) \mid I_\delta(u) > 0, J(u) < d(\delta)\} \cup \{0\},$$

$$\mathcal{V}_\delta := \{u \in H_0^2(\Omega) \mid I_\delta(u) < 0, J(u) < d(\delta)\}.$$

Here, $d(\delta) := \inf_{u \in \mathcal{N}_\delta} J(u) > 0$ is the potential depth of \mathcal{W}_δ . We also define the open sublevels of J , $J^s := \{u \in H_0^2(\Omega) \mid J(u) < s\}$. Furthermore, by the definitions of $J(u)$, \mathcal{N} and J^s , we see that $\mathcal{N}^s := \mathcal{N} \cap J^s \neq \emptyset$ for $\forall s > d$. For any $s > d$, we define

$$(1.7) \quad \lambda_s := \inf \{\|u\|_2 \mid u \in \mathcal{N}^s\}, \quad \Lambda_s := \sup \{\|u\|_2 \mid u \in \mathcal{N}^s\}.$$

It is clear that λ_s is nonincreasing and Λ_s is nondecreasing with respect to s , respectively.

We summarize the main results through the following table. The abbreviations “N.E.,” “E.,” “B.U.,” and “G.E.” denote non-extinction, extinction, blow-up, and global existence of weak solutions of (1.1) or (1.2), respectively. It could be checked that if $J(u_0) < d$, then $I(u_0) \neq 0$.

This paper is arranged as follows. In the next section, we give some important lemmas. Sections 3, 4, and 5 are devoted to the subcritical, the critical, and the

supercritical energy cases, respectively. Section 6 gives the upper and the lower bounds of blow-up time of weak solutions. In Section 7, we show some results about non-extinction or extinction of weak solutions.

2 Preliminary Lemmas

In this section, we give ten lemmas which play important roles in the proof of the main results.

Lemma 2.1 *Let $p > 2$. For any $u \in H_0^2(\Omega)$ with $\|\Delta u\|_2 \neq 0$, we have*

- (i) $\lim_{\lambda \rightarrow 0^+} J(\lambda u) = 0, \lim_{\lambda \rightarrow +\infty} J(\lambda u) = -\infty.$
- (ii) *There exists an unique constant $\lambda^* = \lambda^*(u) > 0$ such that $\frac{d}{d\lambda} J(\lambda u)|_{\lambda=\lambda^*} = 0$. $J(\lambda u)$ is increasing for $0 < \lambda < \lambda^*$, is decreasing for $\lambda^* < \lambda < +\infty$, and takes its maximum at $\lambda = \lambda^*$.*
- (iii) $I(\lambda u) > 0$ for $0 < \lambda < \lambda^*$, $I(\lambda u) < 0$ for $\lambda^* < \lambda < +\infty$, and $I(\lambda^* u) = 0$.

Proof (i) Define the function $j : \lambda \mapsto J(\lambda u)$ for $\lambda > 0$. Then

$$j(\lambda) := J(\lambda u) = \frac{\lambda^2}{2} \|\nabla u\|_2^2 + \frac{\lambda^2}{2} \|\Delta u\|_2^2 + \lambda^3 \int_{\Omega} |\nabla u|^2 \Delta u \, dx - \frac{\lambda^{p+1}}{p+1} \|u\|_{p+1}^{p+1}.$$

We obtain $\lim_{\lambda \rightarrow 0^+} J(\lambda u) = 0$ and $\lim_{\lambda \rightarrow +\infty} J(\lambda u) = -\infty$.

(ii) Elementary calculations imply that

$$j'(\lambda) = \lambda \|\nabla u\|_2^2 + \lambda \|\Delta u\|_2^2 + 3\lambda^2 \int_{\Omega} |\nabla u|^2 \Delta u \, dx - \lambda^p \|u\|_{p+1}^{p+1}.$$

Let $k(\lambda) := \lambda^{-2} j'(\lambda)$. After direct calculation, we have

$$k'(\lambda) = -\lambda^{-2} \|\nabla u\|_2^2 - \lambda^{-2} \|\Delta u\|_2^2 - (p-2)\lambda^{p-3} \|u\|_{p+1}^{p+1} < 0.$$

Since $\lim_{\lambda \rightarrow 0^+} k(\lambda) = +\infty, \lim_{\lambda \rightarrow +\infty} k(\lambda) = -\infty$, there exists a unique constant $\lambda^* > 0$ such that $k(\lambda) > 0$ for $0 < \lambda < \lambda^*, k(\lambda) < 0$ for $\lambda^* < \lambda < +\infty$, and $k(\lambda^*) = 0$. By $j'(\lambda) = \lambda^2 k(\lambda), I(\lambda u) = \lambda j'(\lambda)$, cases (ii) and (iii) hold. ■

Lemma 2.2 *Let $p > 2$. The depth d of the potential well \mathcal{W} is positive.*

Proof Employing Hölder’s inequality, we obtain

$$-\int_{\Omega} |\nabla u|^2 \Delta u \, dx \leq \left(\int_{\Omega} |\nabla u|^4 \, dx \right)^{\frac{1}{2}} \left(\int_{\Omega} |\Delta u|^2 \, dx \right)^{\frac{1}{2}} \leq B_2^2 \|\Delta u\|_2^3.$$

Fix $u \in \mathcal{N}$. Since $p > 2$ and by (1.4),

$$\|\Delta u\|_2^2 \leq \|u\|_{p+1}^{p+1} + 3B_2^2 \|\Delta u\|_2^3 \leq B_1^{p+1} \|\Delta u\|_2^{p+1} + 3B_2^2 \|\Delta u\|_2^3,$$

where B_1 is the optimal constant in the embedding $H_0^2(\Omega) \hookrightarrow L^{p+1}(\Omega)$, and B_2 is the optimal constant in the embedding $W_0^{1,2}(\Omega) \hookrightarrow L^4(\Omega)$. Let

$$(2.1) \quad B_0 := \inf \left\{ x \in (0, +\infty) \mid 1 \leq B_1^{p+1} x^{p-1} + 3B_2^2 x \right\}.$$

Then $J(u) = \frac{1}{6} \|\nabla u\|_2^2 + \frac{1}{6} \|\Delta u\|_2^2 + \frac{p-2}{3(p+1)} \|u\|_{p+1}^{p+1} + \frac{1}{3} I(u) \geq \frac{1}{6} \|\Delta u\|_2^2 \geq \frac{1}{6} B_0^2 > 0$. Therefore, $d = \inf_{u \in \mathcal{N}} J(u) > 0$. ■

Lemma 2.3 Suppose $p > 2$ and $u \in H_0^2(\Omega)$. Define $r(\delta) := \inf \{ x \in (0, +\infty) \mid \delta \leq B_1^{p+1} x^{p-1} + 3B_2^2 x \}$, where B_1, B_2 are defined in (2.1). These are the following results.

- (i) If $I_\delta(u) < 0$, then $\|\Delta u\|_2 > r(\delta)$. Specially, if $I(u) < 0$, then $\|\Delta u\|_2 > r(1)$.
- (ii) If $0 \leq \|\Delta u\|_2 \leq r(\delta)$, then $I_\delta(u) \geq 0$. Specially, if $0 \leq \|\Delta u\|_2 \leq r(1)$, then $I(u) \geq 0$.
- (iii) If $I_\delta(u) = 0$, then $\|\Delta u\|_2 = 0$ or $\|\Delta u\|_2 \geq r(\delta)$. Specially, if $I(u) = 0$, then $\|\Delta u\|_2 = 0$ or $\|\Delta u\|_2 \geq r(1)$.

Proof If $I_\delta(u) < 0$ and $\delta \|\Delta u\|_2^2 < \delta \|\Delta u\|_2^2 + \|\nabla u\|_2^2 \leq B_1^{p+1} \|\Delta u\|_2^{p+1} + 3B_2^2 \|\Delta u\|_2^3$, then $\|\Delta u\|_2 > r(\delta)$, and hence, cases (i) and (ii) hold. If $I_\delta(u) = 0$, we get $\|\Delta u\|_2 \geq r(\delta)$. If $\|\Delta u\|_2 = 0$, $I_\delta(u) = 0$. ■

Lemma 2.4 (Lemma 2.4 in [7]) $d(\delta)$ is increasing for $0 < \delta \leq 1$, is decreasing for $\delta \geq 1$, and takes its maximum $d = d(1)$ at $\delta = 1$.

Lemma 2.5 (Lemma 5 in [15]) Let $p > 2$. Assume $u \in H_0^2(\Omega)$, $0 < J(u) < d$, and $\delta_1 < 1 < \delta_2$ is the two roots of the equation $d(\delta) = J(u)$. Then the sign of $I_\delta(u)$ does not change for $\delta_1 < \delta < \delta_2$.

Lemma 2.6 (Lemma 8 in [15]) Let $p > 2$ and assume that u is a weak solution of problem (1.1) in $\Omega \times [0, T)$ with $0 < J(u_0) < d$. Let $\delta_1 < 1 < \delta_2$ be the two roots of the equation $d(\delta) = J(u_0)$.

- (i) If $I(u_0) > 0$, then $u \in \mathcal{W}_\delta$ for $\delta_1 < \delta < \delta_2$ and $0 < t < T$.
- (ii) If $I(u_0) < 0$, then $u \in \mathcal{V}_\delta$ for $\delta_1 < \delta < \delta_2$ and $0 < t < T$.

Lemma 2.7 Let $p > 2$. $\text{dist}(0, \mathcal{N}) > 0$ and $\text{dist}(0, \mathcal{N}_-) > 0$.

Proof For any $u \in \mathcal{N}$, by the definition of d , we obtain

$$d = \inf_{u \in \mathcal{N}} J(u) \leq \frac{1}{6} B_3^2 \|\Delta u\|_2^2 + \frac{1}{6} \|\Delta u\|_2^2 + \frac{p-2}{3(p+1)} B_1^{p+1} \|\Delta u\|_2^{p+1},$$

which indicates that $\frac{1}{6} B_3^2 \|\Delta u\|_2^2 + \frac{1}{6} \|\Delta u\|_2^2 \geq \frac{d}{2}$ or $\frac{p-2}{3(p+1)} B_1^{p+1} \|\Delta u\|_2^{p+1} \geq \frac{d}{2}$. Then $\|\Delta u\|_2 \geq \left(\frac{3d}{B_3^2}\right)^{\frac{1}{2}}$, or $\|\Delta u\|_2 \geq \left(\frac{3d(p+1)}{2(p-2)B_1^{p+1}}\right)^{\frac{1}{p+1}}$, where B_3 is the optimal constant in $W_0^{1,2}(\Omega) \hookrightarrow L^2(\Omega)$. Let

$$(2.2) \quad C_0 := \min \left\{ \left(\frac{3d}{B_3^2 + 1} \right)^{\frac{1}{2}}, \left[\frac{3d(p+1)}{2(p-2)B_1^{p+1}} \right]^{\frac{1}{p+1}} \right\}.$$

Then $\text{dist}(0, \mathcal{N}) = \inf_{u \in \mathcal{N}} \|\Delta u\|_2 \geq C_0 > 0$. For any $u \in \mathcal{N}_-$, we have $\|\Delta u\|_2^2 \leq B_1^{p+1} \|\Delta u\|_2^{p+1} + 3B_2^2 \|\Delta u\|_2^3$, which implies $\|\Delta u\|_2 \geq B_0$. Here, B_0 is given in (2.1). Then $\text{dist}(0, \mathcal{N}_-) = \inf_{u \in \mathcal{N}_-} \|\Delta u\|_2 \geq B_0 > 0$. ■

Lemma 2.8 Let $p > 2$. For any $s > d$, $u \in J^s \cap \mathcal{N}_+$, $\|\Delta u\|_2 < C_3 := (6s)^{\frac{1}{2}}$.

Proof For any $s > d$ and $p > 2$, $u \in J^s \cap \mathcal{N}_+$, we have

$$s > J(u) = \frac{1}{6} \|\nabla u\|_2^2 + \frac{1}{6} \|\Delta u\|_2^2 + \frac{p-2}{3(p+1)} \|u\|_{p+1}^{p+1} + \frac{1}{3} I(u) > \frac{1}{6} \|\Delta u\|_2^2,$$

which yields $\|\Delta u\|_2 < C_3$. ■

For suitable u and $p > 2$, by using the Gagliardo-Nirenberg inequality, we have

$$(2.3) \quad \|\nabla u\|_4 \leq C_1 \|\Delta u\|_2^a \|u\|_2^{1-a}, \quad \|u\|_{p+1} \leq C_2 \|\Delta u\|_2^b \|u\|_2^{1-b},$$

where $a := \frac{N+4}{8} \in (0, 1)$ and $b := \left(\frac{1}{2} - \frac{1}{p+1}\right) \cdot \frac{N}{2} \in (0, 1)$.

Lemma 2.9 Let $p > 2$. For any $s > d$, λ_s and Λ_s in (1.7) satisfy $0 < K_4 \leq \lambda_s \leq \Lambda_s \leq K_1 < +\infty$, where C_0 is defined in (2.2) and the constants a, b, C_1, C_2 are defined in (2.3); $K_1 := B_4 C_3$,

$$K_2 := \begin{cases} \left[\frac{C_0^{2-b(p+1)}}{2C_2^{p+1}} \right]^{\frac{1}{(1-b)(p+1)}}, & p < \frac{8+N}{N}, \\ \left[\frac{C_3^{2-b(p+1)}}{2C_2^{p+1}} \right]^{\frac{1}{(1-b)(p+1)}}, & p \geq \frac{8+N}{N}, \end{cases}$$

$$K_3 := \left(\frac{C_3^{1-2a}}{6C_1^2} \right)^{\frac{1}{2(1-a)}}, \quad K_4 := \min \{K_2, K_3\}.$$

Proof For $u \in \mathcal{N}^s$, we have $\frac{1}{6} B_4^{-2} \|u\|_2^2 \leq \frac{1}{6} \|\Delta u\|_2^2 \leq J(u) < s$, where B_4 is the optimal constant in the embedding $H_0^2(\Omega) \hookrightarrow L^2(\Omega)$. Then $\|u\|_2 \leq K_1$. By (2.3), we have

$$\begin{aligned} \|\Delta u\|_2^2 &\leq \|u\|_{p+1}^{p+1} + 3\|\nabla u\|_4^2 \|\Delta u\|_2 \\ &\leq C_2^{p+1} \|\Delta u\|_2^{b(p+1)} \|u\|_2^{(1-b)(p+1)} + 3C_1^2 \|\Delta u\|_2^{2a+1} \|u\|_2^{2(1-a)}, \end{aligned}$$

and hence,

$$(2.4) \quad C_2^{p+1} \|\Delta u\|_2^{b(p+1)} \|u\|_2^{(1-b)(p+1)} \geq \frac{1}{2} \|\Delta u\|_2^2, \quad \text{or} \quad 3C_1^2 \|\Delta u\|_2^{2a+1} \|u\|_2^{2(1-a)} \geq \frac{1}{2} \|\Delta u\|_2^2.$$

By the similar proof of Lemma 2.8, we get $\|\Delta u\|_2 < C_3$. Combining with (2.4), we have $\|u\|_2 \geq K_2$ or $\|u\|_2 \geq K_3$. Then $\|u\|_2 \geq K_4 > 0$; hence, $\lambda_s > 0$. ■

Lemma 2.10 (Lemma 2.1 in [8]) *Suppose that a positive, twice-differentiable function $\theta(t)$ satisfies the inequality $\theta''(t)\theta(t) - (1 + \beta)\theta'(t)^2 \geq 0$, $t > 0$, where $\beta > 0$ is a constant. If $\theta(0) > 0$ and $\theta'(0) > 0$, then there exists $0 < t_1 < \frac{\theta(0)}{\beta\theta'(0)}$ such that $\theta(t)$ tends to ∞ as $t \rightarrow t_1$.*

3 The subcritical case

This section is devoted to the property of weak solution of (1.1) or (1.2) under the case $J(u_0) < d$.

Theorem 3.1 *Let $p > 2$. If $J(u_0) < d$ and $I(u_0) > 0$, then problem (1.1) admits a global weak solution $u \in L^\infty(0, \infty; H_0^2(\Omega))$ with $u_t \in L^2(0, \infty; L^2(\Omega))$ and $u(t) \in \mathcal{W}$ for $0 \leq t < \infty$. Moreover, there exists a constant $\hat{C} > 0$ such that $\|u\|_2^2 \leq \|u_0\|_2^2 e^{-\hat{C}t}$.*

Proof The proof is divided into two steps.

Step 1. Global existence. We would use the Galerkin's approximation with some priori estimates. Let $\{\omega_i(x)\}$ be the orthogonal basis of $H_0^2(\Omega)$. Construct the approximate solutions $u_m(x, t)$ of (1.1), $u_m(x, t) := \sum_{i=1}^m a_{mi}(t)\omega_i(x)$, $m = 1, 2, \dots$, $i = 1, 2, \dots, m$, which satisfy

$$(3.1) \quad \begin{aligned} & (u'_m, \omega_i) + (\nabla u_m, \nabla \omega_i) + (\Delta u_m, \Delta \omega_i) - (2|\Delta u_m|^2, \omega_i) + (2|D^2 u_m|^2, \omega_i) \\ & = (|u_m|^{p-1} u_m, \omega_i), \end{aligned}$$

and $u_{0m} := \sum_{i=1}^m b_{mi}(t)\omega_i(x) \rightarrow u_0(x)$ in $H_0^2(\Omega)$ as $m \rightarrow +\infty$.

By the standard theory of ODEs (e.g., the Peanos theorem), we deduce that the existence of a local solution to (3.1). Multiplying (3.1) by $a'_{mi}(t)$, summing over i from 1 to m and integrating with respect to t , we have

$$(3.2) \quad \int_0^t \|u'_m\|_2^2 d\tau + J(u_m(x, t)) = J(u_m(x, 0)), \quad 0 \leq t \leq T.$$

Due to the convergence of $u_{0m} \rightarrow u_0(x)$ in $H_0^2(\Omega)$, one has $J(u_m(x, 0)) \rightarrow J(u_0(x)) < d$, $I(u_m(x, 0)) \rightarrow I(u_0(x)) > 0$. Therefore, for sufficiently large m and any $0 \leq t < +\infty$, we obtain

$$(3.3) \quad \int_0^t \|u'_m\|_2^2 d\tau + J(u_m) = J(u_m(x, 0)) < d, \quad I(u_m(x, 0)) > 0,$$

which implies that $u_m(x, 0) \in \mathcal{W}$ for sufficiently large m .

By applying the similar discussion of Theorem 8 in [15], one could show from (3.3) that $u_m(x, t) \in \mathcal{W}$ for large m and $0 \leq t < +\infty$. Thus, $I(u_m(x, t)) > 0$, $J(u_m(x, t)) < d$ for all $t \in [0, T]$. Then

$$(3.4) \quad \frac{1}{6} \|\nabla u_m\|_2^2 + \frac{1}{6} \|\Delta u_m\|_2^2 + \frac{p-2}{3(p+1)} \|u_m\|_{p+1}^{p+1} < J(u_m(t)) < d.$$

In addition, by using (3.2–3.4), we get for some positive constant C ,

$$(3.5) \quad \|u'_m\|_{L^2(0,T;L^2(\Omega))} \leq C,$$

$$(3.6) \quad \|u_m\|_{L^\infty(0,T;H_0^2(\Omega))} \leq C,$$

$$(3.7) \quad \|u_m\|_{L^\infty(0,T;W_0^{1,2}(\Omega))} \leq C,$$

$$(3.8) \quad \|u_m\|_{L^\infty(0,T;L^{p+1}(\Omega))} \leq C.$$

By the uniform estimates (3.5–3.8), it was seen that the local solutions can be extended globally. Thus, by the standard diagonal method and the Aubin-Lions-Simon theorem, we know there exists a function u and a sequence of $\{u_m\}$ (still by $\{u_m\}$) such that for each $T > 0$, one could obtain

$$(3.9) \quad u'_m \rightharpoonup u', \quad \text{weakly in } L^2(0, T; L^2(\Omega)),$$

$$(3.10) \quad u_m \rightharpoonup u, \quad \text{weakly in } L^\infty(0, T; W_0^{1,2}(\Omega)),$$

$$(3.11) \quad u_m \rightharpoonup u, \quad \text{weakly in } L^\infty(0, T; H_0^2(\Omega)),$$

$$(3.12) \quad |u_m|^{p-1}u_m \rightarrow |u|^{p-1}u, \quad \text{strongly in } L^{\frac{p+1}{p}}(\Omega \times (0, T)),$$

$$(3.13) \quad |\Delta u_m|^2 \rightharpoonup |\Delta u|^2, \quad \text{weakly star in } L^\infty(0, T; L^1(\Omega)),$$

$$(3.14) \quad |D^2 u_m|^2 \rightharpoonup |D^2 u|^2, \quad \text{weakly star in } L^\infty(0, T; L^1(\Omega)),$$

as $m \rightarrow +\infty$. Fix $k \in \mathbb{N}$. In order to show the limit function u in (3.9–3.14) is a weak solution of (1.1), we choose a function $v \in C^1([0, T]; H_0^2(\Omega))$ defined as $v(x, t) := \sum_{i=1}^k l_i(t)\omega_i(x)$, where $\{l_i(t)\}_{i=1}^k$ are arbitrarily given C^1 functions. Taking $m \geq k$ in (3.1), multiplying (3.1) by $l_i(t)$, summing for i from 1 to k , and integrating with respect to t , we obtain

$$(3.15) \quad \int_0^T (u'_m, v) + (\nabla u_m, \nabla v) + (\Delta u_m, \Delta v) - (2|\Delta u_m|^2, v) + (2|D^2 u_m|^2, v) dt = \int_0^T (|u_m|^{p-1}u_m, v) dt.$$

Taking $m \rightarrow +\infty$ in (3.15) and recalling the convergence yield that

$$(3.16) \quad \int_0^T (u', v) + (\nabla u, \nabla v) + (\Delta u, \Delta v) - (2|\Delta u|^2, v) + (2|D^2 u|^2, v) dt = \int_0^T (|u|^{p-1}u, v) dt.$$

Since functions of the form in (3.15) are dense in $L^2(0, T; H_0^2(\Omega))$, (3.16) also holds for all $v \in L^2(0, T; H_0^2(\Omega))$. By the arbitrariness of $T > 0$, we know

$$(u', v) + (\nabla u, \nabla v) + (\Delta u, \Delta v) - (2|\Delta u|^2, v) + (2|D^2 u|^2, v) = (|u|^{p-1}u, v), \quad \text{a.e. } t > 0.$$

Then u is a global weak solution to problem (1.1). To prove (1.6), we first assume that u was smooth enough that $u_t \in L^2(0, T; H_0^2(\Omega))$. Taking $v = u_t$ in (3.16), it is seen that (1.6) is true. By the density of $L^2(0, T; H_0^2(\Omega))$ in $L^2(\Omega \times (0, T))$, we know (1.6) also holds for weak solutions to (1.1). The existence of global solutions to (1.1) is obtained.

Step 2. Decay rate. Taking $\varphi := u$ in (1.5), we get $\frac{d}{dt} \|u\|_2^2 = 2(u_t, u) = -2I(u)$. From Lemma 2.6, we know that $u(x, t) \in \mathcal{W}_\delta$ for $\delta_1 < \delta < \delta_2$ and $0 < t < \infty$ under the condition $J(u_0) < d$ and $I(u_0) > 0$. Thus, $I_{\delta_1}(u) \geq 0$ for $0 < t < \infty$. Therefore,

$$\frac{d}{dt} \frac{\|u\|_2^2}{2} = -I(u) = (\delta_1 - 1)\|\Delta u\|_2^2 - I_{\delta_1}(u) \leq (\delta_1 - 1)B_4^{-2}\|u\|_2^2,$$

where $B_4 > 0$ is the best embedding constant from $H_0^2(\Omega)$ to $L^2(\Omega)$ (i.e., $\|u\|_2 \leq B_4\|\Delta u\|_2$ for $\forall u \in H_0^2(\Omega)$). Consequently, $\|u\|_2^2 \leq \|u_0\|_2^2 e^{-\hat{C}t}$ with $\hat{C} := 2B_4^{-2}(1 - \delta_1) > 0$. ■

Theorem 3.2 *Let $p > 2$ and u be a weak solution of (1.2). If $J(u_0) < d$, $J(u_0) \neq 0$ and $I(u_0) < 0$, then u blows up at some finite time T in the sense of $\lim_{t \rightarrow T} \int_0^t \|u\|_2^2 d\tau = +\infty$.*

Proof We employ the concavity method. Assume on the contrary that u was a global weak solution to (1.2) with $J(u_0) < d$, $J(u_0) \neq 0$, $I(u_0) < 0$ and define $F(t) := \int_0^t \|u\|_2^2 d\tau$, $t \geq 0$. Then $F'(t) = \|u\|_2^2$,

$$(3.17) \quad F''(t) = 2(u, u_t) = -2I(u).$$

By (1.6), (1.4), and (3.17), we have

$$F''(t) \geq -6J(u) + \|\Delta u\|_2^2 = -6J(u_0) + 6 \int_0^t \|u'\|_2^2 d\tau + \|\Delta u\|_2^2.$$

Noticing that $\frac{1}{4}(F'(t) - F'(0))^2 = (\int_0^t \int_\Omega uu' dx d\tau)^2 \leq \int_0^t \|u\|_2^2 d\tau \int_0^t \|u'\|_2^2 d\tau$, we have

$$(3.18) \quad F''(t)F(t) - \frac{3}{2}(F'(t))^2 \geq B_4^{-2}F(t)F'(t) - 6J(u_0)F(t) - 3F'(0)F'(t).$$

In the forthcoming proof, the cases that $J(u_0) < 0$ and $0 < J(u_0) < d$ will be discussed separately.

(i) If $J(u_0) < 0$, then (3.18) implies $F''(t)F(t) - \frac{3}{2}(F'(t))^2 \geq B_4^{-2}F(t)F'(t) - 3F'(0)F'(t)$. Now we will prove that $I(u) < 0$ for all $t > 0$. Otherwise, there must be a constant $t_0 > 0$ such that $I(u(t_0)) = 0$ and $I(u) < 0$ for $0 \leq t < t_0$. From Lemma 2.3 (i, iii), $\|\Delta u\|_2 > r(1)$ for $0 \leq t < t_0$, and $\|\Delta u(t_0)\|_2 \geq r(1)$, where $\|\Delta u(t_0)\|_2 \neq 0$. In fact, using (1.6) and $J(u_0) < 0$, we have $J(u(t_0)) < 0$. Since $0 = F''(t_0) \geq -6J(u(t_0)) + \|\Delta u(t_0)\|_2^2$, we have $\|\Delta u(t_0)\|_2 \neq 0$. Hence, we have $J(u(t_0)) \geq d$, which contradicts to (1.6). By (3.17), we get $F''(t) > 0$ for $t \geq 0$. Since $F'(0) = \|u_0\|_2^2 \geq 0$, $F'(t) > 0$, for large t , $F(t) > 3B_4^2\|u_0\|_2^2$, we have

$$(3.19) \quad F''(t)F(t) - \frac{3}{2}(F'(t))^2 > 0.$$

(ii) If $0 < J(u_0) < d$, then by Lemma 2.6 (ii), we know that $u(x, t) \in \mathcal{V}_\delta$ for $t \geq 0$ and $\delta_1 < \delta < \delta_2$, where $\delta_1 < 1 < \delta_2$ are the two roots of $d(\delta) = J(u_0)$. Hence,

$I_{\delta_2}(u) \leq 0$ and $\|\Delta u\|_2 \geq r(\delta_2)$ for $t \geq 0$. It follows from (3.18) that for $t \geq 0$, $F''(t) \geq 2(\delta_2 - 1)r^2(\delta_2)$, which shows for all $t \geq 0$ that $F'(t) \geq 2(\delta_2 - 1)r^2(\delta_2)t$ and $F(t) \geq (\delta_2 - 1)r^2(\delta_2)t^2$. Therefore, for sufficiently large t , we have

$$B_4^{-2}F'(t) > 12J(u_0), \quad B_4^{-2}F(t) > 6F'(0).$$

Then we obtain (3.19). Due to $(F^{-\frac{1}{2}}(t))'' = -\frac{1}{2}F^{-\frac{5}{2}}(t)[F(t)F''(t) - \frac{3}{2}(F'(t))^2] < 0$, $F^{-\frac{1}{2}}(t)$ is concave in $(0, +\infty)$. So there exists a positive constant t_0 such that $F(t_0) > 0$, $F'(t_0) > 0$ and $(F^{-\frac{1}{2}}(t_0))' < 0$. Since $F(t) > 0$, $F''(t) > 0$ for $t \geq t_0$ and $F'(t_0) > 0$, one can find $(F^{-1/2})'(t) < 0$ for $t \geq t_0$, and hence, there is a constant $T > t_0$ such that $\lim_{t \rightarrow T} F^{-\frac{1}{2}}(t) = 0$; that is, $\lim_{t \rightarrow T} \int_0^t \|u(x, \tau)\|_2^2 d\tau = +\infty$. ■

4 The critical case

For $J(u_0) = d$, the invariance of \mathcal{W}_δ could not be proved in general. By using approximation, we could prove the global existence of weak solutions.

Theorem 4.1 *Let $p > 2$. If $J(u_0) = d$ and $I(u_0) \geq 0$, then problem (1.2) admits a global weak solution $u \in L^\infty(0, \infty; H_0^2(\Omega))$ with $u_t \in L^2(0, \infty; L^2(\Omega))$ and $u \in \overline{\mathcal{W}} = \mathcal{W} \cup \partial\mathcal{W}$ for $0 \leq t < \infty$. Moreover, if $I(u(x, t)) > 0$ for all $t > 0$, $u(x, t)$ does not vanish and there exist positive constants \hat{C}_1 and \hat{C}_2 such that $\|u\|_2^2 \leq \hat{C}_1 e^{-\hat{C}_2 t}$. If not, problem (1.2) admits a solution that vanishes in finite time.*

Proof Let $\lambda_k = 1 - \frac{1}{k}$, $k = 2, 3, \dots$. Consider the approximation problems

$$(4.1) \quad \begin{cases} u_t - \Delta u + \Delta^2 u - \operatorname{div}(2\nabla u \Delta u) + \Delta|\nabla u|^2 = |u|^{p-1}u, & (x, t) \in \Omega \times (0, T), \\ u = 0, \quad \frac{\partial u}{\partial \eta} = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = u_0^k(x) := \lambda_k u_0(x), & x \in \Omega. \end{cases}$$

Noticing that $I(u_0) \geq 0$, by Lemma 2.1 (iii), we could deduce that there exists a unique constant $\lambda^* = \lambda^*(u_0) \geq 1$ such that $I(\lambda^* u_0) = 0$. By $\lambda_k < 1 \leq \lambda^*$, we get $I(u_0^k) = I(\lambda_k u_0) > 0$ and $J(u_0^k) = J(\lambda_k u_0) < J(u_0) = d$. In view of Theorem 3.1, for each k , problem (4.1) admits a global weak solution $u^k(x, t) \in L^\infty(0, \infty; H_0^2(\Omega)) \cap \mathcal{W}$ with $u_t^k \in L^2(0, \infty; L^2(\Omega))$ for $0 \leq t < \infty$ satisfying $\int_0^t \|u_\tau^k\|_2^2 d\tau + J(u^k) = J(u_0^k) < d$. Applying the similar discussion in Theorem 3.1, there exist a subsequence of $\{u^k\}$ and a function u such that u is a weak solution of (1.2) with $I(u) \geq 0$ and $J(u) \leq d$ for $0 \leq t < \infty$.

Let us discuss the decay rate of $\|u\|_2^2$. First, suppose that $I(u) > 0$ for $0 < t < \infty$; then u does not vanish in finite time. Combining with (3.17), we have $u_t \neq 0$. By (1.6), for small $t_0 > 0$, we have $0 < J(u(t_0)) < J(u_0) = d$. Taking $t = t_0$ as the initial time and by Lemma 2.6, we get $u \in \mathcal{W}_\delta$ for $\delta_1 < \delta < \delta_2$ and $t_0 < t < \infty$. Hence, $I_{\delta_1}(u) \geq 0$ for $t_0 < t < \infty$ and

$$\frac{1}{2} \frac{d}{dt} \|u\|_2^2 = -I(u) = (\delta_1 - 1)\|\Delta u\|_2^2 - I_{\delta_1}(u) \leq B_4^{-2}(\delta_1 - 1)\|u\|_2^2,$$

which implies $\|u\|_2^2 \leq \|u(t_0)\|_2^2 e^{-2B_4^{-2}(1-\delta_1)(t-t_0)}$. The decay rate holds with $\hat{C}_2 := 2B_4^{-2}(1-\delta_1)$ and $\hat{C}_1 := \|u(t_0)\|_2^2 e^{2B_4^{-2}(1-\delta_1)t_0}$.

Next, suppose there is a positive constant t_1 such that $I(u) > 0$ for $0 < t < t_1$ and $I(u(x, t_1)) = 0$. Obviously, $u_t \neq 0$ for $0 < t < t_1$ and $\int_0^{t_1} \|u_\tau\|_2^2 d\tau > 0$. Applying (1.6) again, we have $J(u(t_1)) = J(u_0) - \int_0^{t_1} \|u_\tau\|_2^2 d\tau < J(u_0) = d$. By the definition of d , we know $\|u(t_1)\|_2 = 0$ and $\|u\|_2 \equiv 0$ for all $t \geq t_1$. It is seen that such a function u is a weak solution of (1.2) which vanishes in finite time. ■

Theorem 4.2 *Let $p > 2$. If $J(u_0) = d$ and $I(u_0) < 0$, then there exists a finite time T such that the solution u blows up at that time T in the sense of $\lim_{t \rightarrow T} \int_0^t \|u\|_2^2 d\tau = +\infty$.*

Proof Similarly to the proof of Theorem 4.1, we could get

$$F''(t)F(t) - \frac{3}{2} (F'(t))^2 \geq B_4^{-2}F'(t)F(t) - 6J(u_0)F(t) - 3F'(t)F'(0).$$

Since $J(u_0) = d$, $I(u_0) < 0$, by the continuity of $J(u)$ and $I(u)$ with respect to t , there exists a constant $t_0 > 0$ such that $J(u(x, t)) > 0$ and $I(u(x, t)) < 0$ for $0 < t \leq t_0$. From $(u_t, u) = -I(u)$, we have $u_t \neq 0$ for $0 < t \leq t_0$. Furthermore, we have $J(u(t_0)) = d - \int_0^{t_0} \|u_\tau\|_2^2 d\tau < d$. Taking $t = t_0$ as the initial time and by Lemma 2.6, we know that $u(x, t) \in \mathcal{V}_\delta$ for $\delta_1 < \delta < \delta_2$ and $t > t_0$, where $\delta_1 < 1 < \delta_2$ are the two roots of the equation $d(\delta) = J(u_0)$. Thus, $I_{\delta_2}(u) \leq 0$ for $t > t_0$. The rest of the proof is similar to that of Theorem 3.2. ■

5 The supercritical case

We obtain some results for arbitrarily high initial energy.

Theorem 5.1 *Let $p > 2$ and $J(u_0) > d$.*

- (i) *If $u_0 \in \mathcal{N}_+$ and $\|u_0\|_2 \leq \lambda_{J(u_0)}$, then the weak solution u of (1.1) exists globally and $u \rightarrow 0$ as $t \rightarrow +\infty$.*
- (ii) *If $u_0 \in \mathcal{N}_-$ and $\|u_0\|_2 \geq \Lambda_{J(u_0)}$, then the weak solution u of (1.1) blows up in finite time.*

Proof Let T_{\max} be the maximal existence time of (1.1). If $T_{\max} = \infty$, we denote the ω -limit set of u_0 as $\omega(u_0) = \bigcap_{t \geq 0} \overline{\{u(\tau) : \tau \geq t\}}^{H_0^2(\Omega)}$.

(i) Assume that $u_0 \in \mathcal{N}_+$ with $\|u_0\|_2 \leq \lambda_{J(u_0)}$. We first claim that $u(t) \in \mathcal{N}_+$ for all $t \in [0, T_{\max})$. If not, there would exist a constant $t_0 \in (0, T_{\max})$ such that $u(t) \in \mathcal{N}_+$ for $0 \leq t < t_0$ and $I(u(t_0)) = 0$. It follows from $I(u(t)) = -\int_\Omega u_t u dx$ that $u_t \neq 0$ for $(x, t) \in \Omega \times [0, t_0)$. Using (1.6), we have $J(u(t_0)) < J(u_0)$, which deduces $u(t_0) \in J^{I(u_0)}$. Therefore, $u(t_0) \in J^{I(u_0)} \cap \mathcal{N} = \mathcal{N}^{J(u_0)}$. By the definition of $\lambda_{J(u_0)}$, we have

$$(5.1) \quad \lambda_{J(u_0)} \leq \|u(t_0)\|_2.$$

Noticing $I(u(t)) > 0$ for $t \in [0, t_0)$, $\|u(t_0)\|_2 < \|u_0\|_2 \leq \lambda_{J(u_0)}$, a contradiction to (5.1). So $u(t) \in \mathcal{N}_+$ and $u(t) \in J^{I(u_0)} \cap \mathcal{N}_+$ for all $t \in [0, T_{\max})$. Lemma 2.8 shows

$\|\Delta u\|_2 < C_3$, $t \in [0, T_{\max})$, so that $T_{\max} = \infty$. Let ω be an arbitrary element in $\omega(u_0)$. We have $J(\omega) < J(u_0)$, $\|\omega\|_2 < \lambda_{J(u_0)}$, which implies $\omega \notin \mathcal{N}^{J(u_0)}$. Recalling the definition of $\lambda_{J(u_0)}$, $\omega(u_0) \cap \mathcal{N} = \emptyset$. Hence, $\omega(u_0) = \{0\}$. Therefore, the weak solution u of problem (1.1) exists globally and $u \rightarrow 0$ as $t \rightarrow +\infty$.

(ii) Assume that $u_0 \in \mathcal{N}_-$ with $\|u_0\|_2 \geq \Lambda_{J(u_0)}$. We claim that $u(t) \in \mathcal{N}_-$ for all $t \in [0, T_{\max})$. If not, there would be a constant $t_0 \in (0, T_{\max})$ such that $u(t) \in \mathcal{N}_-$ for $0 \leq t < t_0$ and $I(u(t_0)) = 0$. Similarly to case (i), one has $J(u(t_0)) < J(u_0)$, which implies $u(t_0) \in J^{J(u_0)}$, and $u(t_0) \in \mathcal{N}^{J(u_0)}$. By the definition of $\Lambda_{J(u_0)}$, we have

$$(5.2) \quad \Lambda_{J(u_0)} \geq \|u(t_0)\|_2.$$

However, $I(u(t)) < 0$ for $t \in [0, t_0)$, and we get $\|u(t_0)\|_2 > \|u_0\|_2 \geq \Lambda_{J(u_0)}$, a contradiction with (5.2). So $u(t) \in \mathcal{N}_-$, $t \in [0, T_{\max})$. Suppose $T_{\max} = \infty$; for every $\omega \in \omega(u_0)$, we get $\|\omega\|_2 > \Lambda_{J(u_0)}$, $J(\omega) < J(u_0)$, and hence, $\omega \in J^{J(u_0)}$ and $\omega \notin \mathcal{N}^{J(u_0)}$. Recalling the definition of $\Lambda_{J(u_0)}$, we obtain $\omega(u_0) \cap \mathcal{N} = \emptyset$. Thus, $\omega(u_0) = \{0\}$, a contradiction to Lemma 2.7. Hence, $T_{\max} < +\infty$. ■

Remark 5.1 For any $M > d$, there exists $u_M \in \mathcal{N}_-$ such that $J(u_M) \geq M$ and the solution of (1.1) blows up in finite time. In fact, by using Theorem 5.1, if the initial data satisfy the following inequality

$$\frac{3(p+1)}{p-2} J(u_0) |\Omega|^{\frac{p-1}{2}} \leq \|u_0\|_2^{p+1} \leq \|u_0\|_{p+1}^{p+1} |\Omega|^{\frac{p-1}{2}},$$

then the solution of (1.1) blows up in finite time.

6 Blow-up time estimates

Theorem 6.1 For $p > 2$ and $J(u_0) < 0$, the solution $u(x, t)$ blows up at finite time

$$(6.1) \quad T_* \leq \frac{\|u_0\|_2^2}{3|J(u_0)|}.$$

The upper bound of blow-up rate is given as $\|u\|_2 \leq [3 \cdot 2^{-\frac{3}{2}} \eta]^{-1} (T_* - t)^{-1}$, where $\eta := |J(u_0)| L^{-\frac{3}{2}}(0)$ and $L(t) := \frac{1}{2} \|u\|_2^2$.

Proof Let $K(t) := -J(u(t))$. Then $L(0) > 0$ and $K(0) > 0$. By (1.6), we get $K'(t) = \|u'\|_2^2 \geq 0$, which implies $K(t) \geq K(0) > 0$ for all $t \in [0, T)$. By (1.4), we have

$$(6.2) \quad L'(t) = -I(u) \geq -3J(u) = 3K(t).$$

Combining (6.2) with the Cauchy inequality, we obtain

$$(6.3) \quad L(t)K'(t) = \frac{1}{2} \|u\|_2^2 \|u'\|_2^2 \geq \frac{1}{2} (u, u')^2 = \frac{1}{2} (L'(t))^2 \geq \frac{3}{2} L'(t)K(t).$$

According to (6.3), $(K(t)L^{-\frac{3}{2}}(t))' = L^{-\frac{5}{2}}(t)(K'(t)L(t) - \frac{3}{2}K(t)L'(t)) \geq 0$. Therefore,

$$(6.4) \quad 0 < K(0)L^{-\frac{3}{2}}(0) \leq K(t)L^{-\frac{3}{2}}(t) \leq \frac{1}{3} L'(t)L^{-\frac{3}{2}}(t) = -\frac{2}{3} (L^{-\frac{1}{2}}(t))'.$$

Integrating (6.4) with respect to t , we have $\eta t \leq -\frac{2}{3}(L^{-\frac{1}{2}}(t) - L^{-\frac{1}{2}}(0))$, where $\eta := K(0)L^{-\frac{3}{2}}(0)$, and hence, we obtain that there exists a constant $T_* < +\infty$ such that $\lim_{t \rightarrow T_*} L(t) = \infty$ (i.e., the weak solution blows up). Hence, (6.1) holds. Similarly, integrating (6.4) from t to T_* , we have $L(t) \leq [3 \cdot 2^{-1}\eta]^{-2} (T_* - t)^{-2}$. ■

Theorem 6.2 For $p > 2$ and $0 \leq J(u_0) < \frac{1}{6}B_4^{-2}\|u_0\|_2^2$, the solution u blows up at finite time $T_* \leq \frac{16\|u_0\|_2^2}{B_4^{-2}\|u_0\|_2^2 - 6J(u_0)}$.

Proof We assert that for any $t \in [0, T)$, $I(u) < 0$. From (1.4), $I(u_0) \leq 3J(u_0) - \frac{1}{2}\|\Delta u_0\|_2^2 < 0$. If the assertion was not true, there would be a constant $t_0 \in (0, T)$ such that $I(u) < 0$ and $I(u(t_0)) = 0$. From (6.2), we know $\|u\|_2^2$ is strictly increasing with $t \in [0, t_0)$. Therefore,

$$(6.5) \quad J(u(t_0)) \leq J(u_0) < \frac{1}{6}\|\Delta u_0\|_2^2 \leq \frac{1}{6}\|\Delta u(t_0)\|_2^2.$$

From the definition of $J(u)$ and $I(u(t_0)) = 0$, we have $J(u(t_0)) > \frac{1}{6}\|\Delta u(t_0)\|_2^2$. It contradicts to (6.5). Therefore, we have $I(u) < 0$ for any $t \in [0, T)$, and $\|u\|_2^2$ is strictly increasing with respect to t . For any $T \in (0, T^*)$, $\rho > 0$, $\xi > 0$, we define an auxiliary function

$$G(t) := \int_0^t \|u\|_2^2 d\tau + (T - t)\|u_0\|_2^2 + \rho(t + \xi)^2, \quad t \in [0, T^*].$$

By direct calculation, we have $G'(t) = \|u\|_2^2 - \|u_0\|_2^2 + 2\rho(t + \xi) = \int_0^t \frac{d}{d\tau} \|u\|_2^2 d\tau + 2\rho(t + \xi) = 2 \int_0^t (u, u_\tau) d\tau + 2\rho(t + \xi)$, and

$$G''(t) \geq -6J(u) + \|\Delta u\|_2^2 + 2\rho \geq -6J(u_0) + 6 \int_0^t \|u_\tau\|_2^2 d\tau + \|\Delta u\|_2^2 + 2\rho.$$

It is obvious that $G''(t) > 0$. Then $G'(t)$ is increasing on $[0, T]$ and $G'(t) \geq G'(0) > 0$. This indicates that $G(t)$ is increasing on $[0, T]$. Define

$$f(t) := \left[\int_0^t \|u\|_2^2 d\tau + \rho(t + \xi)^2 \right] \left(\int_0^t \|u_\tau\|_2^2 d\tau + \rho \right) - \left[\int_0^t (u, u_\tau) d\tau + \rho(t + \xi) \right]^2.$$

By Cauchy-Schwarz inequality, we have $f(t) \geq 0$. For any $t \in [0, T]$,

$$G(t)G''(t) - \frac{3}{2}(G'(t))^2 \geq G(t) [-6J(u_0) + B_4^{-2}\|u_0\|_2^2 - 4\rho] \geq 0,$$

where $0 < \rho \leq -\frac{3}{2}J(u_0) + \frac{1}{4}B_4^{-2}\|u_0\|_2^2 := \hat{\rho}$. From Lemma 2.10, $T \leq \frac{T\|u_0\|_2^2 + \rho\xi^2}{\rho\xi}$. Fixing $\rho_0 \in (0, \hat{\rho})$, we have $0 < \frac{\|u_0\|_2^2}{\rho_0\xi} < 1$ for any $\xi \in (\frac{\|u_0\|_2^2}{\rho_0}, +\infty)$. This indicates $T \leq \frac{\rho_0\xi^2}{\rho_0\xi - \|u_0\|_2^2}$. The right side of the above formula takes the minimum value at $\xi := \frac{2\|u_0\|_2^2}{\rho_0}$. Then $T \leq \frac{4\|u_0\|_2^2}{\rho_0}$. For $\rho_0 \in (0, \hat{\rho})$, $T < T^* \leq \frac{16\|u_0\|_2^2}{B_4^{-2}\|u_0\|_2^2 - 6J(u_0)}$. ■

Theorem 6.3 *Let $N = 1$. If u blows up in finite time in its $H_0^2(\Omega)$ norm, there exists a positive constant such that*

$$(6.6) \quad T_0 := \int_{M(0)}^{+\infty} \frac{dM}{\tilde{\alpha}M^p + \tilde{\beta}M} \leq T,$$

where $M(0) := \|\Delta u_0\|_2^2$, $\tilde{\alpha} := \gamma^{2p}/\mu$, $\tilde{\beta} := 1/\lambda$, and γ is the embedding constant from $H_0^2(\Omega)$ to $L^{2p}(\Omega)$.

Proof Let $M(t) := \|\Delta u\|_2^2$ and $M(0) > 0$. When $N = 1$, problem (1.1) can be simplified to

$$(6.7) \quad \begin{cases} u_t - \Delta u + \Delta^2 u = |u|^{p-1}u, & (x, t) \in \Omega \times (0, T), \\ u = 0, \quad \frac{\partial u}{\partial \eta} = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = u_0(x), & x \in \Omega. \end{cases}$$

By computation, we have

$$(6.8) \quad M'(t) = 2 \int_{\Omega} (\Delta u - \Delta^2 u + |u|^{p-1}u)\Delta^2 u dx.$$

Let us estimate the integrals in (6.8). We have

$$(6.9) \quad 2 \int_{\Omega} \Delta^2 u \Delta u dx \leq \frac{1}{\lambda} \int_{\Omega} |\Delta u|^2 dx + \lambda \int_{\Omega} (\Delta^2 u)^2 dx,$$

$$(6.10) \quad 2 \int_{\Omega} |u|^{p-1}u \Delta^2 u dx \leq \frac{1}{\mu} \int_{\Omega} |u|^{2p} dx + \mu \int_{\Omega} (\Delta^2 u)^2 dx,$$

with two arbitrary positive constants λ, μ . By replacing (6.9) and (6.10) in (6.8), we obtain

$$(6.11) \quad M'(t) \leq \frac{1}{\mu} \int_{\Omega} |u|^{2p} dx + \frac{1}{\lambda} \int_{\Omega} |\Delta u|^2 dx + (\mu + \lambda - 2) \int_{\Omega} (\Delta^2 u)^2 dx.$$

Choosing $\lambda + \mu \leq 2$, inequality (6.11) reduces to $M'(t) \leq \tilde{\alpha}M^p(t) + \tilde{\beta}M(t)$. Then we get (6.6). ■

7 Extinction and non-extinction

Theorem 7.1 *If $p < 1$ and $N = 1$, then the weak solution u of (1.1) becomes extinct in finite time if the initial data satisfies $\|u_0\|_2^{1-p} \geq D_2 D_1^{-1}$. An upper bound of extinction rate for u is*

$$(7.1) \quad \begin{cases} \|u\|_2 \leq \left[\|u_0\|_2^{1-p} + \frac{1-p}{2} (D_2 - D_1 \|u_0\|_2^{1-p}) t \right]^{\frac{1}{1-p}}, & 0 < t < T_1, \\ \|u\|_2 = 0, & t \in [T_1, \infty), \end{cases}$$

where $T_1 := \frac{2\|u_0\|_2^{1-p}}{(1-p)(D_1\|u_0\|_2^{1-p} - D_2)}$, $D_1 := B_3^{-2} + B_4^{-2}$, and $D_2 := B_5^{p+1}$.

Proof When $N = 1$, we get (6.7). First, we give the upper bound of extinction rate. Multiplying the equation in (6.7) by u and integrating it over $\Omega \times (t, t+h)$ with $h > 0$ and then dividing the result by h yields that

$$(7.2) \quad \frac{1}{h} \int_t^{t+h} \int_{\Omega} u_{\tau} u dx d\tau + \frac{1}{h} \int_t^{t+h} \|\nabla u\|_2^2 d\tau + \frac{1}{h} \int_t^{t+h} \|\Delta u\|_2^2 d\tau = \frac{1}{h} \int_t^{t+h} \|u\|_{p+1}^{p+1} d\tau.$$

Let $h \rightarrow 0^+$ in (7.2). Using the Lebesgue differentiation theorem in [2], one could obtain

$$(7.3) \quad H'(t) + 2\|\nabla u\|_2^2 + 2\|\Delta u\|_2^2 = 2\|u\|_{p+1}^{p+1},$$

where $H(t) := \|u\|_2^2$. According to the embedding relationship $W_0^{1,2}(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow L^{p+1}(\Omega)$ and $H_0^2(\Omega) \hookrightarrow L^2(\Omega)$, we have $\|u\|_2 \leq B_3 \|\nabla u\|_2$, $\|u\|_2 \leq B_4 \|\Delta u\|_2$, $\|u\|_{p+1} \leq B_5 \|u\|_2$. Then (7.3) can be written $H'(t) + 2D_1 H(t) \leq 2D_2 H^{\frac{p+1}{2}}(t)$, where $D_1 := B_3^{-2} + B_4^{-2}$, $D_2 := B_5^{p+1}$. Defining $\varphi(t) := H^{\frac{1-p}{2}}(t)$, we get the following differential inequality:

$$(7.4) \quad \varphi'(t) \leq (1-p)(D_2 - D_1\varphi(t)) := \zeta(t).$$

It is clear from (7.4) that $\zeta(0) < 0$. Recalling the continuity of $\zeta(t)$, there exists a sufficiently small $T_0 > 0$ such that $\zeta(t) < \frac{\zeta(0)}{2} < 0$ for $0 < t \leq T_0$. Then $\varphi'(t) \leq \zeta(0)/2$, which implies that

$$\begin{cases} \varphi(t) \leq \varphi(0) + \frac{\zeta(0)t}{2}, & 0 < t < T_1, \\ \varphi(t) = 0, & t \geq T_1. \end{cases}$$

Obviously, by the definition of $\zeta(t)$, (7.1) holds. ■

Corollary 7.1 *If $p < 1$, $N = 1$, and $J(u_0) \leq 0$, then the weak solution u of (1.1) becomes extinct in finite time with the initial data satisfying $D_2 D_1^{-1} \leq \|u_0\|_2^{1-p} \leq a D_2 D_1^{-1}$. The results of Theorem 7.1 hold, and a lower bound of extinction rate for u is*

$$(7.5) \quad \begin{cases} \|u\|_2 \geq \left[(\|u_0\|_2^{1-p} - D_4 D_3^{-1}) e^{D_3 t} + D_4 D_3^{-1} \right]^{\frac{1}{1-p}}, & 0 < t < T_2, \\ \|u\|_2 = 0, & t \in [T_2, \infty), \end{cases}$$

where $T_2 := \frac{1}{D_3} \log \frac{D_4 D_3^{-1}}{D_4 D_3^{-1} - \|u_0\|_2^{1-p}}$, $D_3 := \frac{1-p}{2} D_1 > 0$, $D_4 := \frac{(2-p)(1-p)}{p+1} D_2 > 0$, and $a := \frac{4-2p}{p+1}$.

Proof The proof of an upper bound of extinction rate for u is the same as Theorem 7.1. We only give the proof of a lower bound of extinction rate. Let $H(t) := \|u\|_2^2$. We have

$$(7.6) \quad \begin{aligned} H'(t) &\geq -6J(u_0) + \|\nabla u\|_2^2 + \|\Delta u\|_2^2 + \frac{2(p-2)}{p+1} \|u\|_{p+1}^{p+1} \\ &\geq (B_3^{-2} + B_4^{-2})H(t) + \frac{2(p-2)}{p+1} B_5^{p+1} H^{\frac{p+1}{2}}(t). \end{aligned}$$

By the definition of $\varphi(t)$ in Theorem 7.1, we have $\varphi'(t) \geq D_3\varphi(t) - D_4$; hence, we get (7.5). ■

Corollary 7.2 Let $p < 1$, $N = 1$, and $\|u_0\|_2^{1-p} \geq D_2 D_1^{-1}$. If $J(u_0) > d$ and $I(u_0) = 0$, then the results of Theorem 7.1 hold.

In the following, we give a result on non-extinction of weak solutions.

Theorem 7.2 Let $J(u_0) \leq 0$. If one of the following conditions holds: (i) $p < 1$ and $\|u_0\|_2^{1-p} > D_4 D_3^{-1}$; (ii) $p = 1$; (iii) $p \geq 2$, then the solution u of (1.1) does not vanish in finite time.

Proof Let $H(t) := \|u\|_2^2$. In the forthcoming proof, the cases $p < 1$, $p = 1$, and $p \geq 2$ will be discussed separately.

(i) $p < 1$, $\|u_0\|_2^{1-p} > D_4 D_3^{-1}$. By the similar proof of Corollary 7.1, we have

$$\|u\|_2 \geq \left[\left(\|u_0\|_2^{1-p} - D_4 D_3^{-1} \right) e^{D_3 t} + D_4 D_3^{-1} \right]^{\frac{1}{1-p}}.$$

(ii) $p = 1$. By (7.6), we obtain $H'(t) \geq D_5 H(t)$, where $D_5 := B_3^{-2} + B_4^{-2} - B_5^2$. Hence, $\|u\|_2 \geq \|u_0\|_2^2 e^{\frac{D_5}{2} t}$.

(iii) $p \geq 2$. By (7.6), we obtain $H'(t) \geq D_6 H(t)$, where $D_6 := B_3^{-2} + B_4^{-2}$. Hence, $\|u\|_2 \geq \|u_0\|_2^2 e^{\frac{D_6}{2} t}$. ■

Acknowledgements The authors would like to express their sincere thanks to the editor and the reviewers for the constructive comments to improve this paper.

Funding This paper is supported by Shandong Provincial Natural Science Foundation of China.

Competing interest The authors declare no competing interests.

Data availability statement All data generated or analyzed during this study are included in this article.

Use of AI tools declaration The authors declare they have not used artificial intelligence (AI) tools in the creation of this article.

References

- [1] L. Agélas, *Global regularity of solutions of equation modeling epitaxy thin film growth in \mathbb{R}^d* , $d = 1, 2$. J. Evol. Equ. 15(2015), 89–106.
- [2] S. N. Antontsev and S. I. Shmarev, *Evolution PDEs with nonstandard growth conditions: Existence, uniqueness, localization, blow-up*, Atlantis Studies in Differential Equations, Vol. 4, Atlantis Press, 2015.
- [3] D. Blömker, C. Gugg, *On the existence of solutions for amorphous molecular beam epitaxy*, Nonlinear Anal. Real World Appl. 3(2002), 61–73.
- [4] D. Blömker, C. Gugg and M. Raible, *Thin-film-growth models: Roughness and correlation functions* Eur. J. Appl. Math. 13(2002), 385–402.

- [5] S. Das Sarma and G. V. Ghalas, *Solid-on-solid rules and models for nonequilibrium growth in 2 + 1 dimensions*. Phys. Rev. Lett. 69(1992), 3762–3765.
- [6] C. Escudero, *Explicit blowing up solutions for a higher order parabolic equation with Hessian nonlinearity*. J. Dyn. Diff. Equat. 35(2023), 2939–2949.
- [7] Y. Z. Han, *A class of fourth-order parabolic equation with arbitrary initial energy*, Nonlinear Anal. Real World Appl. 43(2018), 451–466.
- [8] A. Khelghati and K. Baghaei, *Blow-up phenomena for a nonlocal semilinear parabolic equation with positive initial energy*. Comput. Math. Appl. 70(2015), 896–902.
- [9] J. M. Kim and S. Das Sarma, *Discrete models for conserved growth equations*. Phys. Rev. Lett. 72(1994), 2903–2906.
- [10] R. V. Kohn and X. Yan, *Upper bound on the coarsening rate for an epitaxial growth model*. Comm. Pure Appl. Math. 56(2003), 1549–1564.
- [11] Y. Liu and W. K. Li, *A class of fourth-order nonlinear parabolic equations modeling the epitaxial growth of thin films*. Discrete Contin. Dyn. Syst. (Ser. S) 14(2021), 4367–4381.
- [12] W. W. Mullins, *Theory of thermal grooving*. J. Appl. Phys. 28(1957), 333–339.
- [13] M. Ortiz, E. A. Repetto and H. Si, *A continuum model of kinetic roughening and coarsening in thin films*. J. Mech. Phys. Solids. 47(1999), 697–730.
- [14] M. Winkler, *Global solutions in higher dimensions to a fourth-order parabolic equation modeling epitaxial thin-film growth*. Z. Angew. Math. Phys. 62(2011), 575–608.
- [15] R. Z. Xu and J. Su, *Global existence and finite time blow-up for a class of semilinear pseudo-parabolic equations*. J. Funct. Anal. 264(2013), 2732–2763.
- [16] G. Y. Xu and J. Zhou, *Global existence and blow-up for a fourth order parabolic equation involving the Hessian*. Nonlinear Differ. Equ. Appl. 24(2017), 41.
- [17] A. Zangwill, *Some causes and a consequence of epitaxial roughening*. J. Crystal Growth. 163(1996), 8–21.
- [18] J. Zhou, *L^2 -norm blow-up of solutions to a fourth order parabolic PDE involving the Hessian*, J. Differential Equations 265(2018), 4632–4641.

College of Science, China University of Petroleum, Qingdao 266580, P.R. China

e-mail: bcniu@upc.edu.cn