

CO (2-1) OBSERVATIONS OF MAFFEI 2

Anneila I. Sargent, E.C. Sutton, C.R. Masson, T.G. Phillips
and K.-Y. Lo
Owens Valley Radio Observatory, California Institute of
Technology, Pasadena, CA U.S.A.

Maffei 2 is a highly obscured galaxy, probably of type Sbc, at a distance of 5 Mpc (Allen and Raimond 1972; Spinrad *et al.* 1973). Since it lies close to the Galactic plane, there is considerable confusion in infrared and 21-cm HI observations due to Galactic emission, but investigations of its structure can be carried out at millimeter wavelengths where the Galaxy contribution is confined to a limited velocity range. The high resolution (30") of our CO J=2-1 observations permits both a detailed examination of Maffei 2 and a study of the nature of the gas in its nucleus, through comparison with the CO J=1-0 observations.

The CO (2-1) observations were made with one of the 10.4 meter telescopes at the Owens Valley Radio Observatory in January, 1983, using an SIS receiver (Sutton 1983) and a 512-channel acousto-optical spectrometer (*cf.* Masson 1982). The frequency width of each channel was 1.03 MHz, 1.34 km s⁻¹ at 230 GHz. The receiver temperature, T_r was 550 K SSB. Spectra, taken at spacings of 0.3' on a grid oriented along the major axis of the galaxy (PA = 30°), are shown in Figure 1. The central position, marked by




Figure 1

an asterisk, is α (1950) = $02^{\text{h}}38^{\text{m}}08^{\text{s}}.5$, δ (1950) = $+59^{\circ}23'24''$. At -7 km s^{-1} and -56 km s^{-1} there is some evidence of Galactic emission in the reference position (see the dip near zero velocity in Figure 2). Assuming the source to be just resolved, a beam coupling efficiency 0.52 was adopted in our determination of the corrected antenna temperature, $T_{\text{A}}^{*(2-1)}$. While variations in peak velocity are consistent with the rotation curve, CO(2-1) emission is much more confined than CO(1-0) and 21-cm HI radiation (Rickard *et al.* 1977; Bottinelli *et al.* 1972).




Figure 2

In Figure 2, our central CO(2-1) profile, the CO(1-0) profile from Rickard *et al.* (1977) and the velocity extent of the 21-cm HI emission from Bottinelli *et al.* (1972) are reproduced. Published values of $T_{\text{A}}^{*(1-0)}$ were corrected assuming a beam efficiency of 0.65 (Ulich and Haas 1976) and a source size $1''$ at CO(1-0). Peak $T_{\text{A}}^{*(2-1)} = 1.6\text{K}$, significantly higher than peak $T_{\text{A}}^{*(1-0)} = 0.4\text{K}$. If the nuclear source is ultra-compact, much of this discrepancy could be the effect of beam dilution, but it seems more likely that gas in the nucleus of Maffei 2 is at least partially optically thin, as in the active galaxy, M82 (Sutton *et al.* 1983).

Nuclear activity is often evident in galaxies with nearby companions. Maffei 1, a giant elliptical, is only $40''$ from Maffei 2 on the plane of the sky but, assuming a mass-to-light ratio of 30, is at a distance of 1 Mpc (Spinrad *et al.* 1971). Adopting instead $M/L = 6$ (Sargent *et al.* 1978) puts Maffei 1, like Maffei 2, at about 5 Mpc. Thus Maffei 1 and Maffei 2 may be companion galaxies and interaction between them may have influenced star formation in Maffei 2.

REFERENCES

- Allen, R.J., and Raimond, E.: 1972, *Astron. Astrophys.* 19, 317
 Bottinelli, L., *et al.*: 1972, *Astron. Astrophys.* 12, 264
 Masson, C.R.: 1982, *Astron. Astrophys.* 114, 270
 Rickard, L.J., Turner, B.E., and Palmer, P.: 1977, *Astrophys. J. (Letters)*, 218, L51
 Sargent, W.L.W., Young, P.J., Boksenberg, A., Shortridge, K., Lynds, C.R., and Hartwick, F.D.A.: 1978, *Astrophys. J.* 221, 731
 Spinrad, H. *et al.*: 1971, *Astrophys. J. (Letters)*, 163, L25
 Spinrad, H. *et al.*: 1973, *Astrophys. J.* 180, 351
 Sutton, E.C.: 1983, *IEEE Trans. on MTT*, in press
 Sutton, E.C., Masson, C.R., and Phillips, T.G.: 1983, *Astrophys. J.*, submitted
 Ulich, B.M., and Haas, R.W.: 1976, *Astrophys. J. (Supp.)*, 30, 247