

Descriptive analysis is used for quantitative questions, while thematic analysis covers qualitative questions.

Results: Data collection is currently ongoing and will finish in December 2022. Preliminary results will be presented at the conference.

Conclusions: Conclusions will be derived from the results. It is anticipated that the findings will be helpful in further developing the LTP programme and similar CFT programmes for psychosis.

Disclosure of Interest: None Declared

EPV1001

Alternative initiations with 6-monthly paliperidone palmitate. A descriptive study

O. C. Sobrino^{1*} and B. Sergio²

¹Rey Juan Carlos Hospital, Madrid and ²Hospital Infanta Elena, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2295

Introduction: 6-monthly paliperidone palmitate features an initiation regimen through 1-monthly paliperidone palmitate or 3-monthly paliperidone palmitate. Some patients don't have sufficient adherence to treatment and it is necessary at the clinical level to start directly with 6-monthly paliperidone palmitate. There is little clinical experience with these alternative initiations and through this work those that have been carried out for 12 months at the Rey Juan Carlos Hospital are exposed.

Objectives: The main objective of the study is to describe the alternative initiations performed with 6-monthly paliperidone palmitate in routine clinical practice, having opted for a regimen different from the standard for clinical reasons.

Methods: A retrospective selection of patients will be made through non-probabilistic consecutive sampling, including all patients who have been administered 6-monthly paliperidone palmitate with a standard different from the standard during the last 6 months. To do this, the electronic medical record will be used, first selecting the patients who have started 6-monthly paliperidone palmitate through the anonymized digital records and, later, including in the study only those who have followed an alternative initiation pattern. The variables studied will be the following: age, sex, diagnosis, dose of paliperidone palmitate, initiation regimen, consumption of toxic substances, absenteeism from 6-monthly paliperidone palmitate, visits to the emergency room and admissions.

Results: The study included a total of 5 patients (n=5). 80% of the patients were male and 20% were female. The mean age was 39.7 years. 80% of the patients had an associated substance use disorder. The following alternate starting schedules were with biannual paliperidone palmitate: monthly paliperidone palmitate 150 mg together with semi-annual paliperidone palmitate both on day 1 (n=2) or monthly paliperidone palmitate 150 mg on day 1, monthly paliperidone palmitate 100 mg on day 5 and semi-annual paliperidone palmitate on day 12 (n=3).

A total of 0 visits to the emergency department and 0 admissions were observed after the 6-monthly paliperidone palmitate regimen.

Conclusions: Alternative initiations with 6-monthly paliperidone palmitate may be a useful and safe clinical alternative in patients with very low adherence who, due to clinical needs, require starting 6-monthly paliperidone palmitate earlier in order to guarantee adherence.

Disclosure of Interest: None Declared

EPV1002

Suicidal risk in Latino patients with schizophrenia

O. Olive*, J. A. Ontiveros, A. A. Perez and M. F. Serna

Psychiatry, University Hospital "Dr José E. González" Monterrey, Nuevo Leon, Mexico, Monterrey, Mexico

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2296

Introduction: Schizophrenia is a highly debilitating disorder afflicting more than 24 million individuals worldwide. In Mexico, the Ministry of Health estimates that it affects more than 1 million people. Suicide is one of the main causes of death among people diagnosed with schizophrenia, their risk is 12 times higher than in the general population

Objectives: To evaluate the clinical characteristics of schizophrenic patients at risk of suicide in the Latino population.

Methods: We included 130 patients recruited from genetics studies in Latino patients with schizophrenia from the outpatient and inpatient psychiatric ward of the University Hospital "Dr José E. González" in Monterrey, Nuevo Leon, Mexico. Beck Depression Inventory (BDI-II), the Convergent Functional Information for Suicidality (CFI-S) were applied to all participants. We compared the sociodemographic and clinical characteristics of patients with suicidal risk (measured by history of suicidal attempt or current suicidal risk) and present depressive episode.

Results: Of the 130 participants, 66.9% were male, the median age was 38 years. We found 11(14.3%) patients with suicidal risk and 119 (91.5%) without suicidal risk. Sociodemographic and clinical characteristics of the study population at risk of suicide are described in graphic 1. Patients with a history of suicide attempt scored higher on the CFI-S scale with a median of 0.5 (q1=0.45; q3=0.54) vs. 0.31 (q1=0.22; q3=0.45) (p=0.004)(Graphic 2). Based on the BDI-II we found 2.30% patients showed a mild depression, 20.0% moderate depression and 4.61% severe depression (graphic 3). Schizophrenic patients with a previous suicide attempt and depressive episode had higher score range in CFI-S, median .65 (q1=.65; q2=.59, p=0.000). Also, 63.60% were severely depressive (p=0.000) when they compared with patients with low risk of suicide. Schizophrenic patients with suicidal risk were characterized by: age >=60 years old, unemployment, no children, single, without religion, family history of suicide, previous suicide attempt, depressive episodes, substance abuse, auditory hallucinations and referential delusions.