



Weeds

VOLUME 9
JANUARY 1961
NUMBER 1

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

C. J. WILLARD, *Editor*, Department of Agronomy, The Ohio State University, 1827 Neil Ave., Columbus 10, Ohio

W. C. JACOB, *Business Manager*, Department of Agronomy, University of Illinois, Urbana, Illinois

EDITORIAL COMMITTEE

O. C. LEE, Department of Botany and Plant Pathology, Purdue University, Lafayette, Indiana

R. A. PETERS, Department of Plant Science, University of Connecticut, Storrs, Connecticut

W. C. ROBOCKER, Crops Research Division, ARS, USDA, State College of Washington, Pullman, Washington

E. G. RODGERS, Department of Agronomy, University of Florida, Gainesville, Florida

D. W. STANFORTH, Department of Botany and Plant Pathology, Iowa State University, Ames, Iowa

WEEDS is a quarterly journal published by the Weed Society of America. The subscription price is \$10.00 yearly for four issues; single copies \$2.50. Address all communications regarding subscriptions, advertising, and reprints to W. C. Jacob, Department of Agronomy, University of Illinois. All checks, money orders, and other remittances should be made payable to the Weed Society of America. The editorial office is located at Department of Agronomy, The Ohio State University, 1827 Neil Avenue, Columbus 10, Ohio. Inquiries concerning manuscripts and other material for publication should be addressed to the editorial office. Printing is by the W. F. Humphrey Press, Inc., Geneva, New York.

Entered as second-class matter at the post office at Urbana, Illinois with additional entry at Geneva, New York.

Table of Contents

	<i>Page</i>
Uptake and Distribution of Simazine by Oat and Cotton Seedlings. T. J. Sheets.	1
Chemical Weed Control in Seedlings Alfalfa. III. Effect of Some Herbicides on Five Varieties. M. K. McCarty and Paul F. Sand.	14
Control of Bamboo in Puerto Rico by Herbicides. H. J. Cruzado, T. J. Muzik, and W. C. Kennard.	20
Soil-Moisture Retention and Snow-Holding Capacity as Affected by the Chemical Control of Big Sagebrush. (<i>Artemisia tridentata</i> Nutt.). Leslie W. Sonder and Harold P. Alley.	27
A Device for Subsurface Application of Herbicides. O. B. Wooten and C. G. McWhorter.	36
The Use of Fluorescent Tracers to Study Distribution of Soil-Applied Herbicides. C. G. McWhorter and O. B. Wooten.	42
Control of Blackjack Oak by Basal Spraying with 2,4,5-T. Fred A. Peevy. . .	50
The Role of Surfactants in the Foliar Absorption of Indole-3-Acetic Acid (IAA). R. E. Hughes and V. H. Freed.	54
Recent Advances in the Chemical Control of Wild Oats. G. W. Selleck. . . .	60
Degradation of C ¹⁴ Labeled Simazine by Plants and Soil Microorganisms. Mohamed Tawfik H. Ragab and John P. McCollum.	72
Substituted Urea Herbicides in the Control of Undesirable Hardwoods. Robert G. Merrifield and Thomas Hansbrough.	85
Thistle Control with Cultivation, Cropping and Chemicals. Lyle A. Derscheid, Russel L. Nash and Gail A. Wicks.	90
Acetylenic Derivatives as Herbicides. J. K. Leasure and M. J. Josephs. . . .	103
Confining Fumigants in the Soil with Surface Seals. Stanley J. Pieczarka and G. F. Warren.	106
Photochemical Degradation of Diuron and Monuron. L. W. Weldon and F. L. Timmons.	111
Pre-Emergence Weed Control in Soybeans with Mixtures of Two Herbicides. J. R. Hay.	117
The Medusahead Problem in Idaho. Paul J. Torell, Lambert C. Erickson, and Robert H. Haas.	124
Responses of Corn Hybrids to Yellow Foxtail Competition. David W. Stanforth.	132
Pre-Emergence Herbicides for Crabgrass Control and Their Effects on Germination of Turfgrass Species. Felix V. Juska.	137
Phytotoxicity of Simazine. O. C. Burnside and R. Behrens.	145
News and Notes.	158
Sustaining Members.	159
Committee Membership of Weed Society of America.	160
Bibliography of Weed Investigations, April to June, 1960.	161


Advertisers Index

Spraying Systems Co.	ii
duPont de Nemours & Co.	iii
Stauffer Chemical Co.	iv
Chipman Chemical Co.	v


SPRAYING SYSTEMS
TeeJet
 SPRAY NOZZLES

the precision
 nozzle for
 effective
 spraying


Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.



TEEJET
 SPRAY
 NOZZLE
 male pipe
 connection



TEEJET
 SPRAY
 NOZZLE
 female pipe
 connection



INTER-
 CHANGE-
 ABLE
 ORIFICE TIPS
 flat and cone
 spray types

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single boom, that are not accessible with a boom.

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

SPRAYING SYSTEMS CO.
Engineers and Manufacturers

3296 RANDOLPH STREET

BELLWOOD, ILLINOIS

For Agriculture and Industry . . .

Du Pont WEED KILLERS

***offer new economies
and efficiency in killing
weeds, grass and brush***

KARMEX® diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KARMEX® DL for pre-emergence weed control in cotton.

TELVAR® monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KLOBEN® neburon for weed and grass control in nursery plantings and tomatoes.

DYBAR® fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

TRYSBEN® 200, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broadleaf weeds; and certain woody vines and brush.

AMMATE® X for long-term, low-cost brush control.



Better Things for Better Living . . . *through Chemistry*

Eptam[®]

HERBICIDE

CONTROLS NUTGRASS AND OTHER GRASSES IN POTATOES

For further information contact one of
these conveniently located
STAUFFER CHEMICAL COMPANY offices.

Glendale, Ariz.
P. O. Box 505
Phone: YE 9-1428

North Little Rock, Ark.
P. O. Box 509
Phone: FR 4-7445

Los Angeles, Calif.
500 S. Virgil Ave.
Phone: MA 7-5771

San Francisco, Calif.
636 California Street
Phone: SU 1-5350

Tampa, Florida
P. O. Box 5388
Phone: 4-3127

Harvey, La.
P. O. Box 127
Phone: FL 1-4208

So. Omaha, Nebr.
P. O. Box 222
Phone: OR 3200

New York, N. Y.
380 Madison Ave.
Phone: OX 7-0600

North Portland, Ore.
Suttle Road
Phone: AV 6-3675

Houston, Texas
Heights Station
P. O. Box 7222
Phone: UN 9-3333

Lubbock, Texas
P. O. Box 592
Phone: PO 2-8133

Weslaco, Texas
P. O. Box 495
Phone: WO 8-2112



©Stauffer's Reg. T. M.
for an herbicide.



ATLACIDE: Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water.

ATLACIDE-2,4-D: Offers dual killing action of sodium chlorate and 2,4-D.

CHLOREA: A combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray.

CHLOREA GRANULAR Similar to Chlorea, but a "pelletized" material. No mixing...easy to apply by hand or with mechanical spreader.

CHLORAX "40": A composition of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys vegetation and algae in ponds and lakes. Controls certain turf weeds. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

ATLAS "A" SP: A spray powder containing 94% sodium arsenite. Used for the same purposes as Atlas "A". Applied as a water-mixed spray.

CHIP-CAL: Special low-lime calcium arsenate formulation in *granular* and *spray powder* forms. Used as crabgrass preventative for established turf. Applied before crabgrass seed starts to germinate. Treatment lasts for several years.

2,4-D & 2,4,5-T: 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

METHOXONE: Contains MCP sodium salt. Used for control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

BUTOXONE: A liquid containing 4-(2,4-DB). For selective control of many broadleaf weeds in established alfalfa and certain other legumes.

Write for Weed Control Bulletins

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

Bound Brook, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.

Bessemer, Ala.

Manufacturers of Weed Killers Since 1912