



Integral mean estimates for univalent and locally univalent harmonic mappings

Suman Das and Anbareeswaran Sairam Kaliraj

Abstract. We verify a long-standing conjecture on the membership of univalent harmonic mappings in the Hardy space, whenever the functions have a “nice” analytic part. We also produce a coefficient estimate for these functions, which is in a sense best possible. The problem is then explored in a new direction, without the additional hypothesis. Interestingly, our ideas extend to certain classes of locally univalent harmonic mappings. Finally, we prove a Baernstein-type extremal result for the function $\log(h' + cg')$, when $f = h + \bar{g}$ is a close-to-convex harmonic function, and c is a constant. This leads to a sharp coefficient inequality for these functions.

1 Introduction

A central problem pertaining to the growth of univalent harmonic mappings is to determine the exact range of $p > 0$ so that these functions belong to the harmonic Hardy space h^p . The early developments in this direction were due to Abu-Muhanna and Lyzzaik [1], which were later improved by Nowak [17]. She proved sharp results for the classes of convex and close-to-convex harmonic functions, and conjectured an analogous range of p for the whole class S_H of normalized univalent harmonic functions. The aim of this paper is to verify this conjecture for functions $f \in S_H$ with an additional property. The main theorems and their implications are presented in Section 2, while the proofs are given in Section 4. Section 3 contains results from the literature that are useful for our purpose.

A function f analytic in the open unit disk $\mathbb{D} = \{z : |z| < 1\}$ is of class H^p ($0 < p \leq \infty$) if the integral means

$$M_p(r, f) = \begin{cases} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{\frac{1}{p}}, & 0 < p < \infty, \\ \sup_{|z|=r} |f(z)|, & p = \infty \end{cases}$$

remain bounded as $r \rightarrow 1^-$. The norm of a function $f \in H^p$ is defined as $\|f\|_p = \lim_{r \rightarrow 1^-} M_p(r, f)$. Integral means and H^p spaces play a fundamental role

Received by the editors September 20, 2023; revised January 7, 2024; accepted January 9, 2024.

Published online on Cambridge Core January 15, 2024.

The second author was partly supported by the Core Research Grant (CRG/2022/008920) from the Science and Engineering Research Board (SERB), India.

AMS subject classification: 31A05, 30C45, 30H10.

Keywords: Integral means, univalent harmonic functions, growth problems, Hardy space, Baernstein-type inequalities.



in studies concerning the growth of functions; we refer to the books [9, 14, 18] for a detailed survey.

A harmonic function f in the unit disk has a unique representation $f = h + \bar{g}$, where h, g are analytic functions in \mathbb{D} with $g(0) = 0$. The function f is locally univalent and sense-preserving if, and only if, the Jacobian $J_f(z) = |h'(z)|^2 - |g'(z)|^2$ is positive for all $z \in \mathbb{D}$. Let S_H be the class of all sense-preserving univalent harmonic functions f in \mathbb{D} , normalized by the conditions $h(0) = g(0) = h'(0) - 1 = 0$. Denote by K_H and C_H the subclasses of S_H consisting of harmonic mappings onto convex and close-to-convex regions, respectively. Let $S_H^0 = \{f = h + \bar{g} \in S_H : g'(0) = 0\}$, $K_H^0 = K_H \cap S_H^0$, and $C_H^0 = C_H \cap S_H^0$. Two leading examples of univalent harmonic functions are the harmonic Koebe function

$$K(z) = H(z) + \overline{G(z)} = \left(\frac{z - \frac{1}{2}z^2 + \frac{1}{6}z^3}{(1-z)^3} \right) + \overline{\left(\frac{\frac{1}{2}z^2 + \frac{1}{6}z^3}{(1-z)^3} \right)}$$

which maps \mathbb{D} onto $\mathbb{C} \setminus (-\infty, -1/6]$, and the function

$$L(z) = H_1(z) + \overline{G_1(z)} = \left(\frac{z - \frac{1}{2}z^2}{(1-z)^2} \right) + \overline{\left(\frac{-\frac{1}{2}z^2}{(1-z)^2} \right)}$$

which maps \mathbb{D} onto the half-plane $\operatorname{Re}\{w\} > -1/2$. It is obvious that $K \in C_H^0$ and $L \in K_H^0$. More details on univalent harmonic functions can be found in [6, 10].

A harmonic function f is said to be of class h^p ($0 < p \leq \infty$) if $\|f\|_p < \infty$. Let us give an account of the problem considered in this paper. Every function $f = h + \bar{g} \in S_H$ admits the representation

$$(1.1) \quad h(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=1}^{\infty} b_n z^n.$$

Let us define $\alpha = \sup_{f \in S_H} |a_2|$. Then α has crucial influence in the growth of functions in S_H (see [21] for an exposition). Interest in the boundary behavior of functions $f \in S_H$ was initiated by Abu-Muhanna and Lyzzaik [1], who proved that $f \in h^p$ for $p < 1/(2\alpha + 2)^2$. Bshouty and Hengartner [5] proposed to find the exact range of $p > 0$ for which $f \in h^p$. In [17], Nowak improved the range to $p < 1/\alpha^2$, and obtained the sharp results that $f \in h^p$ for $p < 1/2$ (resp. $p < 1/3$) whenever f is a convex (resp. close-to-convex) harmonic function. These observations led her to conjecture that if $f \in S_H$, then $f \in h^p$ for $p < 1/\alpha$. The conjecture seems challenging, and in the relatively recent development [20], the authors verified it by confining interest to harmonic quasiconformal mappings.

In this paper, we first give a relation between $M_p(r, f)$ and $M_p(r, h')$, which naturally allows us to check the boundedness of $\|f\|_p$ whenever h' behaves “nicely.” As it turns out, this can be achieved by placing the simple restriction that h' takes no value infinitely often. An analytic function φ in \mathbb{D} has *valency* m if φ takes no value more than m times. More generally, let for a function φ analytic in \mathbb{D} , $W(R)$ denote the area (regions covered multiply being counted multiply) of the image of \mathbb{D} under φ that lies in the disk $|w| \leq R$. If $W(R) \leq m\pi R^2$ for all $R > 0$, where m is a positive number, then we say that φ has *mean valency* m . This notion is due to Spencer, who proved the following inequality on the integral means of such functions.

Theorem A [22] If f has mean valency $m \geq 1$, $f(0) = 0$, and $p > 0$, then

$$M_p^p(r, f) \leq K \int_0^r \frac{M_\infty^p(s, f)}{s} ds,$$

where $K = K(m, p)$ is independent of f .

This result was initially proved by Prawitz (see [19, Theorem 5.1]) for univalent functions.

Over the years, extremal problems on the growth of univalent functions have been widely studied. Let \mathcal{S} denote the class of univalent analytic functions f in \mathbb{D} with $f(0) = f'(0) - 1 = 0$. A seminal result in this direction is the following inequality of Baernstein.

Theorem B [2] If $f \in \mathcal{S}$ and $0 < p < \infty$, then

$$M_p(r, f) \leq M_p(r, k),$$

where $k(z) = z/(1-z)^2$ is the Koebe function.

Baernstein's theorem was extended to derivatives by Leung [15] and Brown [4] for certain subclasses of \mathcal{S} . We refer to [3, 12, 13, 16] for more problems of this type. In [12], Girela obtained similar results for the functions $\log(f(z)/z)$. These functions appear in the definition of *logarithmic coefficients* γ_n of a function $f \in \mathcal{S}$:

$$\log \frac{f(z)}{z} = 2 \sum_{n=1}^{\infty} \gamma_n z^n.$$

The logarithmic coefficients were instrumental in de Branges' proof of the Bieberbach conjecture (see [8]). Girela's work readily led to the sharp inequality

$$\sum_{n=1}^{\infty} |\gamma_n|^2 \leq \frac{\pi^2}{6},$$

an important estimate earlier obtained by Duren and Leung [11]. Interestingly, Girela proved the following extremal result for close-to-convex functions.

Theorem C [12] Let $f \in \mathcal{S}$ be close-to-convex, and let $0 < p \leq 2$. Then

$$M_p(r, \log f') \leq M_p(r, \log k'),$$

where k is the Koebe function.

On the other hand, similar problems for harmonic functions remained unexplored until very recently, when the present authors initiated the study of Baernstein-type inequalities for harmonic functions.

Theorem D [7] Let $0 < p < \infty$. If $f = h + \bar{g} \in C_H^0$, then

$$M_p(r, h') \leq M_p(r, H') \quad \text{and} \quad M_p(r, g') \leq M_p(r, G').$$

For $f = h + \bar{g} \in K_H^0$, we have

$$M_p(r, h') \leq M_p(r, H_1') \quad \text{and} \quad M_p(r, g') \leq M_p(r, G_1').$$

The functions H, G and H_1, G_1 come from the harmonic functions K and L , respectively.

To explore the logarithmic coefficients in the setting of a harmonic mapping $f = h + \bar{g}$, it is not feasible to consider $f(z)/z$, as this function need not be harmonic, neither is the logarithm of a harmonic function defined in the literature. One cannot consider the functions $(h(z) + cg(z))/z$ (c constant) either, since $h(z) + cg(z)$ may have zeros at points other than the origin. Therefore, proceeding along the line of Theorem C, the functions $\log(h' + cg')$ seem to be the most natural choice.

Thus, we conclude the paper with a harmonic analogue of Girela's result: we prove that Theorem C remains true for the functions $\log(h' + cg')$, whenever $f = h + \bar{g}$ is a close-to-convex harmonic function and c is a constant. This takes forward the authors' earlier line of work in [7].

2 Main theorems and auxiliary results

As discussed, we start with the following simple but useful lemma, the proof of which is just a slight modification of standard techniques.

Lemma 1 Let $0 < p \leq 1$. Suppose $f = h + \bar{g}$ is a locally univalent, sense-preserving harmonic function in \mathbb{D} with $f(0) = 0$. Then

$$M_p^p(r, f) \leq C \int_0^r (r-s)^{p-1} M_p^p(s, h') ds,$$

where C is a constant independent of f .

This, together with Theorem D, immediately give integral mean estimates for convex and close-to-convex harmonic functions.

Corollary 1 Let $0 < p \leq 1$, then we have the inequalities

$$M_p(r, f) \leq C \int_0^r (r-s)^{p-1} M_p^p(s, H') ds, \quad \text{whenever } f \in C_H^0,$$

$$M_p(r, f) \leq C \int_0^r (r-s)^{p-1} M_p^p(s, H_1') ds, \quad \text{whenever } f \in K_H^0.$$

Remark 1 The estimates for $p > 1$ for these classes are already obtained in [7]. It is pertinent to mention that an elementary upper bound can be easily given in the case of convex harmonic functions. If $f = h + \bar{g}$ is convex, it is well known [6, Theorem 5.7] that h is close-to-convex, and $|g(z)| \leq |h(z)|$, $z \in \mathbb{D}$. Therefore, from [19, Theorem 5.1],

we have

$$M_p^p(r, f) \leq 2^p M_p^p(r, h) \leq 2^p p \int_0^r \frac{M_\infty^p(s, h)}{s} ds \leq 2^p p \int_0^r s^{p-1} (1-s)^{-2p} ds.$$

The last integral is the incomplete beta function $B(r; p, 1-2p)$.

As a consequence of Lemma 1, we verify Nowak's conjecture for certain functions in S_H . Indeed, the result is true for a more general class of functions. Let us recall that a family \mathcal{L} of harmonic functions in \mathbb{D} is said to be *linear invariant* (see [21]) if for every $f = h + \bar{g} \in \mathcal{L}$, the functions

$$T_\varphi(f(z)) = \frac{f(\varphi(z)) - f(\varphi(0))}{\varphi'(0)h'(\varphi(0))}, \quad \varphi \in \text{Aut}(\mathbb{D}),$$

belong to \mathcal{L} , where $\text{Aut}(\mathbb{D})$ denotes the set of analytic automorphisms of \mathbb{D} . Our result does not require univalence, and holds for any linear invariant class \mathcal{H} of locally univalent and sense-preserving harmonic functions (with usual normalizations), for which $\alpha(\mathcal{H}) = \sup_{f \in \mathcal{H}} |a_2|$ is finite. For the remainder of this paper, we preserve the notation \mathcal{H} to mean any such class of locally univalent harmonic functions.

Theorem 1 Let $f = h + \bar{g} \in S_H$ be such that h' has finite mean valency. Then $f \in h^p$ for $p < 1/\alpha$. If $f \in \mathcal{H}$ and h' has finite mean valency, then $f \in h^p$ for $p < 1/\alpha(\mathcal{H})$.

Lemma 1 also leads us to the following coefficient bound for these functions.

Theorem 2 Suppose $f = h + \bar{g} \in S_H$ has series representation (1.1), and h' has finite mean valency. Then $|a_n|$ and $|b_n|$ are $O(n^{\alpha-1})$, $n = 2, 3, 4, \dots$. For $f \in \mathcal{H}$ with h' having finite mean valency, $|a_n|$ and $|b_n|$ are $O(n^{\alpha(\mathcal{H})-1})$.

Remark 2 This coefficient estimate for $f \in S_H$ is in a sense best possible. The conjectured value of α is 3. Given this, Theorem 2 asserts that $|a_n|$ and $|b_n|$ are $O(n^2)$, which is the same order as in the harmonic analogue of the Bieberbach conjecture [6].

The problem, even without the assumption of finite mean valency, can be explored in another direction to produce a very interesting result. Since S_H is known to be linear invariant, for $f = h + \bar{g} \in S_H$ and any $\zeta \in \mathbb{D}$, the function

$$T(z) = \frac{f\left(\frac{z+\zeta}{1+\bar{\zeta}z}\right) - f(\zeta)}{(1-|\zeta|^2)h'(\zeta)} = A(z) + \overline{B(z)} \quad (z \in \mathbb{D})$$

again lies in S_H . Let us write

$$A(z) = z + a_2(\zeta)z^2 + a_3(\zeta)z^3 + \dots$$

A customary computation gives

$$a_2(\zeta) = \frac{1}{2} \left\{ (1-|\zeta|^2) \frac{h''(\zeta)}{h'(\zeta)} - 2\bar{\zeta} \right\}.$$

Since $|a_2(\zeta)| \leq \alpha$, we find that

$$\left| \frac{h''(\zeta)}{h'(\zeta)} \right| \leq \frac{C}{1-|\zeta|} \quad (\zeta \in \mathbb{D}),$$

for some positive constant C . However, this is the extreme bound on h''/h' that a function $f = h + \bar{g} \in S_H$ can possess. In general, it is reasonable to expect a large subclass of S_H to have slightly restricted growth, or more precisely, to exhibit the bound

$$\left| \frac{h''(\zeta)}{h'(\zeta)} \right| \leq \frac{C}{(1-|\zeta|)^\beta} \quad (0 \leq \beta < 1).$$

The expression h''/h' is of special interest in the theory of univalent functions. For example, it appears in the definition of the Schwarzian derivative, as well as in characterization results for certain geometric subclasses (e.g., convex and close-to-convex). The growth condition on h''/h' leads us to the following result on the membership of univalent and locally univalent harmonic functions in the Hardy space.

Theorem 3 *Let $f = h + \bar{g} \in S_H$ be such that*

$$(2.1) \quad \left| \frac{h''(z)}{h'(z)} \right| \leq \frac{C}{(1-|z|)^\beta},$$

for some β with $0 \leq \beta < 1$. Then $f \in h^p$ for $p < 2(1-\beta)/\alpha$. Analogously, if $f = h + \bar{g} \in \mathcal{H}$ satisfies the growth estimate (2.1), then $f \in h^p$ for $p < 2(1-\beta)/\alpha(\mathcal{H})$.

Finally, we prove Girela's result (Theorem C) in the setting of harmonic functions.

Theorem 4 *Suppose $0 < p \leq 2$ and $f = h + \bar{g} \in C_H^0$. Then, for any constant $c \in \mathbb{D}$, we have*

$$M_p(r, \log(h' + cg')) \leq M_p(r, \log(H' + G')).$$

The bound is sharp.

Like logarithmic coefficients in the case of analytic functions, it is interesting to study the power series coefficients of $\log(h'(z) + cg'(z))$. Suppose $\log(h'(z) + cg'(z)) = \sum_{n=1}^{\infty} \lambda_n z^n$. Then Theorem 4 has the following implication.

Corollary 2 *For $f = h + \bar{g} \in C_H^0$, we have the sharp inequality*

$$\sum_{n=1}^{\infty} |\lambda_n|^2 \leq \frac{14\pi^2}{3}.$$

3 Preliminaries

Definition 1 [2] For a real-valued function $g(x)$ integrable over $[-\pi, \pi]$, the Baernstein star-function is defined as

$$g^*(\theta) = \sup_{|E|=2\theta} \int_E g(x) dx \quad (0 \leq \theta \leq \pi),$$

where $|E|$ is the Lebesgue measure of the set $E \subseteq [-\pi, \pi]$.

The following properties of the star-function are due to Leung [15].

Lemma A For $g, h \in L^1[-\pi, \pi]$,

$$[g(\theta) + h(\theta)]^* \leq g^*(\theta) + h^*(\theta).$$

Equality holds if g, h are both symmetric in $[-\pi, \pi]$ and nonincreasing in $[0, \pi]$.

Lemma B If g, h are subharmonic functions in \mathbb{D} and g is subordinate to h , then for each r in $(0, 1)$,

$$g^*(re^{i\theta}) \leq h^*(re^{i\theta}), \quad 0 \leq \theta \leq \pi.$$

Lemma C If $p(z) = e^{i\beta} + p_1z + \dots$ is analytic and of positive real part in \mathbb{D} , then

$$(\log |p(re^{i\theta})|)^* \leq \left(\log \left| \frac{1 + re^{i\theta}}{1 - re^{i\theta}} \right| \right)^*, \quad 0 \leq \theta \leq \pi.$$

An important feature in the proof of Lemma C is that a rotation factor does not affect the star-function. This observation will be suitably deployed in our work.

Definition 2 [12] A domain D in \mathbb{C} is said to be *Steiner symmetric* if its intersection with each vertical line is either empty or a segment placed symmetrically with respect to the real axis.

The next result by Girela is crucial in the proof of Theorem 4.

Lemma D [12] Let F and \mathcal{F} be analytic in $\overline{\mathbb{D}}$ and satisfy:

- (i) $F(0) = \mathcal{F}(0) = 0$,
- (ii) $(\operatorname{Re} F)^* \leq (\operatorname{Re} \mathcal{F})^*$ in $\mathbb{D}^+ = \{z \in \mathbb{D} : \operatorname{Im} z > 0\}$,
- (iii) $\min_{z \in \mathbb{D}} \operatorname{Re} \mathcal{F}(z) \leq \min_{z \in \mathbb{D}} \operatorname{Re} F(z) \leq \max_{z \in \mathbb{D}} \operatorname{Re} F(z) \leq \max_{z \in \mathbb{D}} \operatorname{Re} \mathcal{F}(z)$,
- (iv) \mathcal{F} is univalent and $\mathcal{F}(\mathbb{D})$ is a Steiner symmetric domain.

Then, for $0 < p \leq 2$,

$$\int_{-\pi}^{\pi} |F(e^{i\theta})|^p d\theta \leq \int_{-\pi}^{\pi} |\mathcal{F}(e^{i\theta})|^p d\theta.$$

4 Proofs

4.1 Proof of Lemma 1

In what follows, C will denote a positive constant that is not necessarily the same at each occurrence. Let $|\nabla f| = (|h'|^2 + |g'|^2)^{1/2}$. For $0 \leq r_1 < r_2 < 1$, we have

$$\begin{aligned} |f(r_2 e^{i\theta}) - f(r_1 e^{i\theta})| &= \left| \int_{r_1}^{r_2} \frac{d}{dt} f(te^{i\theta}) dt \right| \\ &\leq \int_{r_1}^{r_2} \left| e^{i\theta} h'(te^{i\theta}) + \overline{e^{i\theta}} g'(te^{i\theta}) \right| dt \\ &\leq \sqrt{2} \int_{r_1}^{r_2} (|h'(te^{i\theta})|^2 + |g'(te^{i\theta})|^2)^{1/2} dt \\ &= \sqrt{2} \int_{r_1}^{r_2} |\nabla f(te^{i\theta})| dt \\ &\leq \sqrt{2} (r_2 - r_1) \sup_{r_1 \leq t \leq r_2} |\nabla f(te^{i\theta})|. \end{aligned}$$

Since f is sense-preserving, i.e., $|g'(z)| < |h'(z)|$ for every $z \in \mathbb{D}$, we find that

$$|\nabla f(te^{i\theta})| \leq |h'(te^{i\theta})| + |g'(te^{i\theta})| < 2|h'(te^{i\theta})|.$$

Therefore,

$$\begin{aligned} M_p^p(r_2, f) - M_p^p(r_1, f) &\leq \frac{1}{2\pi} \int_0^{2\pi} |f(r_2 e^{i\theta}) - f(r_1 e^{i\theta})|^p d\theta \\ &\leq 2^{3p/2} (r_2 - r_1)^p \frac{1}{2\pi} \int_0^{2\pi} \left(\sup_{r_1 \leq t \leq r_2} |h'(te^{i\theta})| \right)^p d\theta. \end{aligned}$$

An appeal to the Hardy–Littlewood maximal theorem gives

$$\frac{1}{2\pi} \int_0^{2\pi} \left(\sup_{r_1 \leq t \leq r_2} |h'(te^{i\theta})| \right)^p d\theta \leq C M_p^p(r_2, h'),$$

so that

$$(4.1) \quad M_p^p(r_2, f) - M_p^p(r_1, f) \leq C (r_2 - r_1)^p M_p^p(r_2, h').$$

Let $0 < r < 1$ be arbitrary, and let $r_n = r(1 - 2^{-n})$, $n = 0, 1, 2, \dots$. Clearly, $M_p(0, f) = 0$ as $f(0) = 0$. Using (4.1), we find that

$$\begin{aligned} M_p^p(r_{n+1}, f) &= \sum_{k=1}^{n+1} [M_p^p(r_k, f) - M_p^p(r_{k-1}, f)] \\ &\leq C \sum_{k=1}^{n+1} (r_k - r_{k-1})^p M_p^p(r_k, h') \\ &= C \sum_{k=1}^{n+1} (r_k - r_{k-1}) (r - r_k)^{p-1} M_p^p(r_k, h'), \end{aligned}$$

since $r_k - r_{k-1} = 2^{-k}r = r - r_k$. We now let $n \rightarrow \infty$ and obtain

$$M_p^p(r, f) \leq C \int_0^r (r-s)^{p-1} M_p^p(s, h') ds$$

by means of Riemann integration.

4.2 Proof of Theorem 1

Let $1/(\alpha+1) < p \leq 1$ and $f \in S_H$. We may choose $r_n = 1 - 2^{-n}$ in the proof of Lemma 1 to obtain

$$(4.2) \quad \|f\|_p^p \leq C \int_0^1 (1-s)^{p-1} M_p^p(s, h') ds,$$

whenever the integral is finite. We break the integral in two parts, to separately deal with possible complications around 0 and 1. For example, let us write

$$(4.3) \quad \|f\|_p^p \leq C \left[\int_0^{1/4} (1-s)^{p-1} M_p^p(s, h') ds + \int_{1/4}^1 (1-s)^{p-1} M_p^p(s, h') ds \right].$$

Throughout our computations, the constants will be denoted by C , K , etc., and they need not be the same at each occurrence. We do this for convenience, as constants do not affect our purpose.

We appeal to Theorem A for an estimate of $M_p^p(s, h')$. Since h' is finitely mean valent, so is zh' . Therefore, we have

$$(4.4) \quad M_p^p(s, zh') \leq K \int_0^s \frac{M_\infty^p(r, zh')}{r} dr.$$

It is known (see [10, p. 98]) that

$$(4.5) \quad M_\infty(r, h') \leq \frac{(1+r)^{\alpha-1}}{(1-r)^{\alpha+1}}.$$

This, together with (4.4), implies

$$(4.6) \quad M_p^p(s, h') \leq \frac{K}{s^p} \int_0^s \frac{r^{p-1}}{(1-r)^{(\alpha+1)p}} dr.$$

For $s \leq 1/4$,

$$M_p^p(s, h') \leq \frac{K}{s^p} \int_0^{1/4} \frac{r^{p-1}}{(1-r)^{(\alpha+1)p}} dr \leq \frac{K_1}{s^p} \int_0^{1/4} r^{p-1} dr \leq \frac{K_2}{s^p} \quad (\text{as } p \leq 1).$$

For $s > 1/4$,

$$\begin{aligned} M_p^p(s, h') &\leq \frac{K}{s^p} \int_0^{1/4} \frac{r^{p-1}}{(1-r)^{(\alpha+1)p}} dr + \frac{K}{s^p} \int_{1/4}^s \frac{r^{p-1}}{(1-r)^{(\alpha+1)p}} dr \\ &\leq \frac{K_2}{s^p} + K_3 \int_{1/4}^s \frac{dr}{(1-r)^{(\alpha+1)p}} \end{aligned}$$

$$\begin{aligned}
&= \frac{K_2}{s^p} + \frac{K_3}{(\alpha+1)p-1} \left[\frac{1}{(1-s)^{(\alpha+1)p-1}} - K_4 \right] \quad (\text{as } p > 1/(\alpha+1)) \\
&\leq \frac{K_2}{s^p} + \frac{K_5}{(1-s)^{(\alpha+1)p-1}}.
\end{aligned}$$

Substituting these bounds in (4.3), we see that

$$\begin{aligned}
\|f\|_p^p &\leq C_1 \int_0^{1/4} s^{-p}(1-s)^{p-1} ds + C_1 \int_{1/4}^1 s^{-p}(1-s)^{p-1} ds + C_2 \int_{1/4}^1 \frac{ds}{(1-s)^{p\alpha}} \\
&= C_1 \int_0^1 s^{-p}(1-s)^{p-1} ds + C_2 \int_{1/4}^1 \frac{ds}{(1-s)^{p\alpha}}.
\end{aligned}$$

The first integral is the beta function $B(1-p, p)$ and converges for every $p \in (0, 1)$. The second integral is finite for $p < 1/\alpha$. Therefore, $f \in h^p$ for $p < 1/\alpha$.

To prove the result for $f \in \mathcal{H}$, we just need to establish the bound

$$M_\infty(r, h') \leq \frac{(1+r)^{\alpha(\mathcal{H})-1}}{(1-r)^{\alpha(\mathcal{H})+1}}.$$

The argument presented here is well known (see, for example, [10, p. 98]), and will be useful in the later results. Since \mathcal{H} is linear invariant, for any $\zeta \in \mathbb{D}$, the function

$$T(z) = \frac{f\left(\frac{z+\zeta}{1+\bar{\zeta}z}\right) - f(\zeta)}{(1-|\zeta|^2)h'(\zeta)} = A(z) + \overline{B(z)} \quad (z \in \mathbb{D})$$

is in \mathcal{H} . We write

$$A(z) = z + a_2(\zeta)z^2 + a_3(\zeta)z^3 + \dots,$$

so that

$$a_2(\zeta) = \frac{1}{2} \left\{ (1-|\zeta|^2) \frac{h''(\zeta)}{h'(\zeta)} - 2\bar{\zeta} \right\}.$$

Since $|a_2(\zeta)| \leq \alpha(\mathcal{H})$, we find that

$$\frac{2r^2 - 2r\alpha(\mathcal{H})}{1-r^2} \leq \operatorname{Re} \left\{ \frac{zh''(z)}{h'(z)} \right\} \leq \frac{2r^2 + 2r\alpha(\mathcal{H})}{1-r^2} \quad (|z| = r),$$

which is equivalent to

$$\frac{2r - 2\alpha(\mathcal{H})}{1-r^2} \leq \frac{\partial}{\partial r} \{ \log |h'(re^{i\theta})| \} \leq \frac{2r + 2\alpha(\mathcal{H})}{1-r^2}.$$

Now we integrate from 0 to r to reach the estimate

$$(4.7) \quad \frac{(1-r)^{\alpha(\mathcal{H})-1}}{(1+r)^{\alpha(\mathcal{H})+1}} \leq |h'(z)| \leq \frac{(1+r)^{\alpha(\mathcal{H})-1}}{(1-r)^{\alpha(\mathcal{H})+1}}.$$

The rest of the proof follows through an identical argument, and the details are omitted.

4.3 Proof of Theorem 2

We see that

$$(n+1)|a_{n+1}| = \left| \frac{1}{2\pi i} \int_{|z|=r} \frac{h'(z)}{z^{n+1}} dz \right| \leq r^{-n} M_1(r, h').$$

We see from (4.6), for $p = 1$, that

$$M_1(r, h') \leq \frac{K}{r} \int_0^r \frac{ds}{(1-s)^{\alpha+1}} \leq \frac{K}{r(1-r)^\alpha},$$

for some absolute constant K which varies through occurrences. Therefore,

$$(n+1)|a_{n+1}| \leq \frac{K}{r^{n+1}(1-r)^\alpha}.$$

The function on the right-hand side attains a minimum at $r = (n+1)/(n+1+\alpha)$. With this choice of r , we obtain

$$(n+1)|a_{n+1}| \leq K \left(1 + \frac{\alpha}{n+1}\right)^{n+1} (n+1+\alpha)^\alpha \leq K(n+1)^\alpha.$$

Therefore, replacing $n+1$ by n ,

$$|a_n| \leq Kn^{\alpha-1}.$$

Using a similar argument, one can show that $|b_n| \leq Kn^{\alpha-1}$. The proof for the second part of the theorem is identical, except for α suitably replaced by $\alpha(\mathcal{H})$.

4.4 Proof of Theorem 3

The Hardy–Stein identity (see [19, p. 126]) for the function h' implies that

$$\frac{d}{dr} \left[r \frac{d}{dr} M_p^p(r, h') \right] = \frac{p^2 r}{2\pi} \int_0^{2\pi} |h'(re^{i\theta})|^{p-2} |h''(re^{i\theta})|^2 d\theta.$$

Since $M_p^p(r, h')$ is a (strictly) increasing function of r , we have

$$\frac{d}{dr} M_p^p(r, h') > 0.$$

Therefore,

$$\begin{aligned} \frac{d^2}{dr^2} M_p^p(r, h') &\leq \frac{p^2}{2\pi} \int_0^{2\pi} |h'(re^{i\theta})|^{p-2} |h''(re^{i\theta})|^2 d\theta \\ &= \frac{p^2}{2\pi} \int_0^{2\pi} |h'(re^{i\theta})|^p \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right|^2 d\theta \\ &\leq \frac{p^2}{2\pi} \int_0^{2\pi} \frac{(1+r)^{(\alpha-1)p}}{(1-r)^{(\alpha+1)p}} \cdot \frac{C^2}{(1-r)^{2\beta}} d\theta \quad (\text{by (4.5)}) \\ &\leq \frac{K}{(1-r)^{(\alpha+1)p+2\beta}}, \end{aligned}$$

for some positive constant K , which is not the same in subsequent occurrences. Integrating twice from 0 to s ($s < 1$), we arrive at the estimate

$$M_p^p(s, h') \leq \frac{K}{(1-s)^{(\alpha+1)p+2\beta-2}}.$$

Thus, an appeal to (4.2) gives

$$\|f\|_p^p \leq C \int_0^1 (1-s)^{p-1} M_p^p(s, h') ds \leq C \int_0^1 \frac{ds}{(1-s)^{\alpha p+2\beta-1}}.$$

The last integral converges for $\alpha p + 2\beta - 1 < 1$, or equivalently, $p < 2(1-\beta)/\alpha$. Therefore, $f \in h^p$ for $p < 2(1-\beta)/\alpha$.

The proof for $f \in \mathcal{H}$ is similar, one only needs to replace α by $\alpha(\mathcal{H})$, wherever applicable.

4.5 Proof of Theorem 4

Let $0 < r < 1$ and write

$$F(z) = \log(h'(rz) + cg'(rz)), \quad \mathcal{F}(z) = \log(H'(rz) + G'(rz)).$$

Clearly, $F(0) = \mathcal{F}(0) = 0$. Lemma 4 of [23] implies that there exist real numbers μ, θ_0 and an analytic function $Q(z)$ with positive real part, such that

$$\operatorname{Re} \left\{ Q(z) \left[ie^{i\theta_0} (1-z^2) \left(e^{-i\mu} h'(e^{i\theta_0} z) + e^{i\mu} g'(e^{i\theta_0} z) \right) \right] \right\} > 0, \quad z \in \mathbb{D}.$$

Let

$$P(z) = Q(z) \left[ie^{i\theta_0} (1-z^2) \left(e^{-i\mu} h'(e^{i\theta_0} z) + e^{i\mu} g'(e^{i\theta_0} z) \right) \right].$$

Without any loss of generality, we may assume $|Q(0)| = 1$, so that $|P(0)| = 1$. Since $Q(z)$ has positive real part, so does $1/Q(z)$. The dilatation $w(z) = g'(z)/h'(z)$ satisfies $w(0) = 0$ and $|w(z)| < 1$ for all $z \in \mathbb{D}$. We see that

$$\begin{aligned} \log |h'(e^{i\theta_0} z) + cg'(e^{i\theta_0} z)| &= \log |P(z)| + \log \left| \frac{1}{Q(z)} \right| + \log \left| \frac{1}{1-z^2} \right| \\ &\quad + \log \left| \frac{1}{1+e^{2i\mu} w(e^{i\theta_0} z)} \right| + \log |1+cw(e^{i\theta_0} z)|. \end{aligned}$$

In view of Lemmas A–C, we have for $z \in \mathbb{D}^+$,

$$\begin{aligned} (\log |h'(z) + cg'(z)|)^* &\leq \left(\log \left| \frac{1+z}{1-z} \right| \right)^* + \left(\log \left| \frac{1+z}{1-z} \right| \right)^* \\ &\quad + \left(\log \left| \frac{1}{1-z^2} \right| \right)^* + \left(\log \left| \frac{1}{1-z} \right| \right)^* + (\log |1+z|)^*, \end{aligned}$$

which implies

$$(\log |h'(z) + cg'(z)|)^* \leq \left(\log \left| \frac{(1+z)^2}{(1-z)^4} \right| \right)^* = (\log |H'(z) + G'(z)|)^*,$$

i.e., $(\operatorname{Re} F)^* \leq (\operatorname{Re} \mathcal{F})^*$. For $f = h + \bar{g} \in C_H^0$, the function $f + c\bar{f} \in C_H$ for every constant $c \in \mathbb{D}$. Also, it is known that C_H is linear invariant and $\alpha(C_H) = 3$. Therefore, (4.7) leads to the inequalities

$$\frac{(1-r)^2}{(1+r)^4} \leq |h'(re^{i\theta}) + cg'(re^{i\theta})| \leq \frac{(1+r)^2}{(1-r)^4},$$

so that

$$\min_{z \in \mathbb{D}} \operatorname{Re} \mathcal{F}(z) \leq \min_{z \in \mathbb{D}} \operatorname{Re} F(z) \leq \max_{z \in \mathbb{D}} \operatorname{Re} F(z) \leq \max_{z \in \mathbb{D}} \operatorname{Re} \mathcal{F}(z).$$

That \mathcal{F} is univalent and $\mathcal{F}(\mathbb{D})$ is a Steiner symmetric domain can be proved using an argument similar to the one presented in [12, Lemma 1], we include the details below for the convenience of the reader. Therefore, the proof of the theorem is completed through an appeal to Lemma D. Since the harmonic Koebe function $K = H + \bar{G} \in C_H^0$, the sharpness can be seen by letting $c \rightarrow 1^-$.

Proposition 1 *Let $\mathcal{G}(z) = \log(H'(z) + G'(z))$. Then \mathcal{G} is univalent and $\mathcal{G}(\mathbb{D})$ is a Steiner symmetric domain.*

Proof For $0 < r < 1$, we see that

$$\operatorname{Re} \mathcal{G}(re^{i\theta}) = \log \frac{1}{|1 - re^{i\theta}|^2} + 2 \log \left| \frac{1 + re^{i\theta}}{1 - re^{i\theta}} \right|.$$

Thus, $\operatorname{Re} \mathcal{G}(re^{i\theta})$ is symmetric with respect to θ on $[-\pi, \pi]$, and strictly decreases on $[0, \pi]$. It is easy to see that $\operatorname{Im} \mathcal{G}(re^{i\theta}) > 0$ in $0 < \theta < \pi$, because the same holds for $\log(1/(1-z)^2)$ and $\log((1+z)/(1-z))$. Also, $\mathcal{G}(re^{-i\theta}) = \overline{\mathcal{G}(re^{i\theta})}$. Therefore, \mathcal{G} is injective on $|z| = r$ and hence the argument principle implies that \mathcal{G} is univalent in $|z| \leq r$. Finally, \mathcal{G} maps $\{|z| = r\}$ onto a simple closed curve, which is symmetric with respect to the real axis, with decreasing real part for θ increasing from 0 to π . This shows that $\mathcal{G}(|z| < r)$ is a Steiner symmetric domain. As this is true for every $r \in (0, 1)$, the desired conclusion follows. ■

Remark 3 The restriction $0 < p \leq 2$ in Theorem 4 is imposed by Lemma D. In other words, we do not know if Theorem 4 remains valid for $p > 2$.

4.6 Proof of Corollary 2

Let $\log(H'(z) + G'(z)) = \sum_{n=1}^{\infty} c_n z^n$. Through a routine computation, we see that

$$c_n = \frac{2(2 - (-1)^n)}{n}.$$

For $p = 2$, Theorem 4 gives

$$\sum_{n=1}^{\infty} |\lambda_n|^2 r^{2n} \leq \sum_{n=1}^{\infty} |c_n|^2 r^{2n}.$$

Letting $r \rightarrow 1$, we obtain

$$\sum_{n=1}^{\infty} |\lambda_n|^2 \leq \sum_{n=1}^{\infty} |c_n|^2 = \frac{14\pi^2}{3}.$$

The sharpness follows from Theorem 4.

Acknowledgments The authors thank Prof. S. Ponnusamy and Prof. D. Bshouty for useful suggestions on an earlier version of this paper. The first author thanks Mr. Kapil Jaglan for helpful discussions during the course of this work.

Competing interests The authors have no competing interests to declare that are relevant to the content of this paper.

References

- [1] Y. Abu-Muhanna and A. Lyzzaik, *The boundary behaviour of harmonic univalent maps*. Pacific J. Math. 141(1990), no. 1, 1–20.
- [2] A. Baernstein, *Integral means, univalent functions and circular symmetrization*. Acta Math. 133(1974), 139–169.
- [3] A. Baernstein, *Some sharp inequalities for conjugate functions*. Indiana Univ. Math. J. 27(1978), 833–852.
- [4] J. E. Brown, *Derivatives of close-to-convex functions, integral means and bounded mean oscillation*. Math. Z. 178(1981), 353–358.
- [5] D. Bshouty and W. Hengartner, *Univalent harmonic mappings in the plane*. Ann. Univ. Mariae Curie-Skłodowska Sect. A 48(1994), 12–42.
- [6] J. Clunie and T. Sheil-Small, *Harmonic univalent functions*. Ann. Acad. Sci. Fenn. Ser. A I Math. 9(1984), 3–25.
- [7] S. Das and A. Sairam Kaliraj, *Growth of harmonic mappings and Baernstein type inequalities*. Potential Anal. (2023), 17 pp. <https://doi.org/10.1007/s11118-023-10081-w>
- [8] L. de Branges, *A proof of the Bieberbach conjecture*. Acta Math. 154(1985), 137–152.
- [9] P. L. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics, 38, Academic Press, New York, 1970.
- [10] P. L. Duren, *Harmonic mappings in the plane*, Cambridge Tracts in Mathematics, 156, Cambridge University Press, Cambridge, 2004.
- [11] P. L. Duren and Y. J. Leung, *Logarithmic coefficients of univalent functions*. J. Analyse Math. 36(1979), 36–43.
- [12] D. Girela, *Integral means and BMOA-norms of logarithms of univalent functions*. J. Lond. Math. Soc. 33(1986), no. 2, 117–132.
- [13] D. Girela, *Integral means, bounded mean oscillation, and Gelfer functions*. Proc. Amer. Math. Soc. 113(1991), 365–370.
- [14] P. Koosis, *Introduction to H^p spaces*, 2nd ed., Cambridge Tracts in Mathematics, 115, Cambridge University Press, Cambridge, 1998.
- [15] Y. J. Leung, *Integral means of the derivatives of some univalent functions*. Bull. Lond. Math. Soc. 11(1979), 289–294.
- [16] M. Nowak, *Some inequalities for BMOA functions*. Complex Variables Theory Appl. 16(1991), 81–86.
- [17] M. Nowak, *Integral means of univalent harmonic maps*. Ann. Univ. Mariae Curie-Skłodowska Sect. A 50(1996), 155–162.
- [18] M. Pavlović, *Introduction to function spaces on the disk*, Biblioteka Posebna Izdanja, 20, Matematički Institut SANU, Belgrade, 2004.
- [19] C. Pommerenke, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [20] S. Ponnusamy, J. Qiao, and X. Wang, *Uniformly locally univalent harmonic mappings*. Proc. Indian Acad. Sci. Math. Sci. 128(2018), Article no. 32, 14 pp.
- [21] T. Sheil-Small, *Constants for planar harmonic mappings*. J. Lond. Math. Soc. 42(1990), no. 2, 237–248.

- [22] D. C. Spencer, *On finitely mean valent functions*. Proc. Lond. Math. Soc. 47(1941), no. 2, 201–211.
- [23] X.-T. Wang, X.-Q. Liang, and Y.-L. Zhang, *Precise coefficient estimates for close-to-convex harmonic univalent mappings*. J. Math. Anal. Appl. 263(2001), 501–509.

Department of Mathematics, Indian Institute of Technology Ropar, Rupnagar, Punjab 140001, India,
e-mail: sumandas471@gmail.com sairam@iitrpr.ac.in