

Comparison of neural substrates of temporal discounting between youth with autism spectrum disorder and with obsessive-compulsive disorder

C. O. Carlisi^{1*†}, L. Norman^{1†}, C. M. Murphy^{1,2,3}, A. Christakou⁴, K. Chantiluke¹, V. Giampietro⁵, A. Simmons^{5,6,7}, M. Brammer⁵, D. G. Murphy^{2,3}, MRC AIMS consortium[‡], D. Mataix-Cols⁸ and K. Rubia¹

¹Department of Child and Adolescent Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK

²Department of Forensic and Neurodevelopmental Sciences, Sackler Institute for Translational Neurodevelopmental Sciences, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK

³Behavioural Genetics Clinic, Adult Autism Service, Behavioural and Developmental Psychiatry Clinical Academic Group, South London and Maudsley Foundation NHS Trust, London, UK

⁴Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK

⁵Department of Neuroimaging, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK

⁶National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) for Mental Health at South London and Maudsley NHS Foundation Trust and Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

⁷Department of Neurobiology, Care Sciences and Society, Center for Alzheimer Research, Division of Clinical Geriatrics, Karolinska Institutet, Stockholm, Sweden

⁸Department of Clinical Neuroscience, Centre for Psychiatry Research, Karolinska Institutet, Stockholm, Sweden

Background. Autism spectrum disorder (ASD) and obsessive-compulsive disorder (OCD) share abnormalities in hot executive functions such as reward-based decision-making, as measured in the temporal discounting task (TD). No studies, however, have directly compared these disorders to investigate common/distinct neural profiles underlying such abnormalities. We wanted to test whether reward-based decision-making is a shared transdiagnostic feature of both disorders with similar neurofunctional substrates or whether it is a shared phenotype with disorder-differential neurofunctional underpinnings.

Methods. Age and IQ-matched boys with ASD ($N=20$), with OCD ($N=20$) and 20 healthy controls, performed an individually-adjusted functional magnetic resonance imaging (fMRI) TD task. Brain activation and performance were compared between groups.

Results. Boys with ASD showed greater choice-impulsivity than OCD and control boys. Whole-brain between-group comparison revealed shared reductions in ASD and OCD relative to control boys for delayed-immediate choices in right ventromedial/lateral orbitofrontal cortex extending into medial/inferior prefrontal cortex, and in cerebellum, posterior cingulate and precuneus. For immediate-delayed choices, patients relative to controls showed reduced activation in anterior cingulate/ventromedial prefrontal cortex reaching into left caudate, which, at a trend level, was more decreased in ASD than OCD patients, and in bilateral temporal and inferior parietal regions.

Conclusions. This first fMRI comparison between youth with ASD and with OCD, using a reward-based decision-making task, shows predominantly shared neurofunctional abnormalities during TD in key ventromedial, orbital- and inferior fronto-striatal, temporo-parietal and cerebellar regions of temporal foresight and reward processing, suggesting trans-diagnostic neurofunctional deficits.

Received 14 September 2016; Revised 10 March 2017; Accepted 29 March 2017; First published online 24 April 2017

Key words: ASD, fMRI, OCD, temporal discounting.

* Address for correspondence: C. Carlisi, BA, Department of Child and Adolescent Psychiatry/MRC Social, Genetic and Developmental Psychiatry (SGDP) Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, 16 DeCrespigny Park, London, SE5 8AF, UK.

(Email: carlisi.christina@gmail.com)

† CC and LN contributed equally to this work.

‡ The MRC AIMS Consortium is a collaboration of Autism research centres in the UK including the Institute of Psychiatry, Psychology & Neuroscience, London, the Autism Research Centre, University of Cambridge, and the Autism Research Group, University of Oxford. It is funded by the MRC UK and headed by the Department of Forensic and Developmental Sciences, Institute of Psychiatry, Psychology & Neuroscience. The Consortium members are in alphabetical order: Bailey A.J., Baron-Cohen S., Bolton P.F., Bullmore E.T., Carrington S., Chakrabarti B., Daly E.M., Deoni S.C., Ecker C., Happe F., Henty J., Jezzard P., Johnston P., Jones D.K., Lombardo M., Madden A., Mullins D., Murphy C.M., Murphy D.G., Pasco G., Sadek S., Spain D., Steward R., Suckling J., Wheelwright S., Williams S.C.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Autism Spectrum Disorder (ASD) is characterized by social communication difficulties and stereotyped repetitive behaviours (American Psychiatric Association, 2013) with a prevalence of 0.6–2%, predominantly in males (Blumberg *et al.* 2013). Obsessive-Compulsive Disorder (OCD) involves recurrent, intrusive and distressing thoughts (obsessions) and repetitive rituals (compulsions) (American Psychiatric Association, 2013), affecting 1–3% of the population with a higher male prevalence in children (Ruscio *et al.* 2010). These disorders are highly comorbid, with rates exceeding 30% (Simonoff *et al.* 2008) and can sometimes be clinically difficult to separate (Doshi-Velez *et al.* 2014).

The allowance of co-diagnosis of OCD with ASD in DSM-5 questions whether phenotypes common to both disorders are mediated by shared or disorder-specific mechanisms. Characteristic behaviours observed in ASD are wide-ranging and heterogeneous but can include physical rocking, tapping, counting and behavioural inflexibility (e.g. insistence on performing actions in a certain order). Similarly, behaviours in OCD vary widely, but compulsions often include hand-washing, checking, and, sometimes seemingly similar to ASD, counting and behavioural inflexibility surrounding order and symmetry. It has been hypothesized that in both cases, these behaviours may relate to abnormalities in fronto-striatal circuitry that is also important in reward-based decision-making (Langen *et al.* 2011). In ASD, repetitive behaviours are often considered soothing and rewarding, while in OCD, compulsions are performed to reduce anxiety and are often debilitating. However, despite this distinction, converging evidence suggests repetitive behaviours in ASD and OCD may be mediated by shared mechanisms including behavioural disinhibition or motivation control (Hollander *et al.* 2007). Such impairments may maintain diminished control over repetitive behaviours in ASD and compulsions in OCD and involve goal-directed reward-based decision-making. A meta-analysis of structural and functional neuroimaging studies comparing ASD and OCD found shared reduced structure and function during cognitive control in medial prefrontal regions but that OCD had disorder-specific increased function and structure in basal ganglia and insula while ASD had disorder-specific functional reduction in DLPFC and reduced PCC deactivation, presumably reflecting disorder-specific fronto-striato-insular dysregulation in OCD but fronto-striato-insular maldevelopment in ASD, both underpinned by shared reduced prefrontal control (Carlisi *et al.* 2016b).

Both disorders also share deficits in motivated 'hot' executive functions (EF) (Zelazo & Müller,

2007) including reward-based decision-making measured by choice-impulsivity tasks of gambling and temporal discounting (TD) (Hill, 2004; Sanders *et al.* 2008; Abramovitch *et al.* 2013; Chen *et al.* 2016). TD requires choosing between small immediate rewards and larger later rewards, assessing the extent to which a reward is subjectively discounted when delayed in time (Rubia *et al.* 2009). The ability to inhibit immediate reward choices and wait for larger rewards depends on well-developed frontal lobe-mediated motivation control and temporal foresight and is a key for mature decision-making. A TD function is typically hyperbolic, with steeper rates reflecting more impulsive choice behaviour (Richards *et al.* 1999) (see online Supplement). TD matures with age (Christakou *et al.* 2011; Steinbeis *et al.* 2016) and varies among individuals (Odum, 2011), with steeper TD observed in younger people and individuals with attention deficit hyperactivity disorder (ADHD) and related impulsive disorders (Rubia *et al.* 2009; Noreika *et al.* 2013). Functional magnetic resonance imaging (fMRI) studies of TD in healthy adults and children implicate ventromedial-fronto-limbic networks of reward-based decision-making and dorsolateral and inferior-fronto-insula-striato-parietal networks of temporal foresight (Christakou *et al.* 2011; Chantiluke *et al.* 2014b; Wesley & Bickel, 2014).

People with ASD have been shown to have deficits in reward-motivated and forward-thinking behaviour including reward processing and reversal learning (Scott-Van Zeeland *et al.* 2010; Chantiluke *et al.* 2015a), incentive processing (Dichter *et al.* 2012), planning (Ozonoff & Jensen, 1999; Geurts *et al.* 2004; Hill, 2004) and TD (Chantiluke *et al.* 2014b). However, there have also been negative findings (Antrop *et al.* 2006; Demurie *et al.* 2013). ASD is characterized by fronto-temporo-limbic abnormalities mediating socio-emotional processes (Via *et al.* 2011; Philip *et al.* 2012; Carlisi *et al.* 2016b), and in ventromedial/fronto-limbic brain regions involved in TD (Christakou *et al.* 2011; Peters & Büchel, 2011) during reward-related and planning tasks (Just *et al.* 2007; Schmitz *et al.* 2008; Dichter *et al.* 2012; Kohls *et al.* 2013). However, only one fMRI study has been published investigating the neural correlates of TD in adolescents with ASD, which found a weaker relationship between task-performance and bilateral superior temporal and right insular activation relative to controls (Chantiluke *et al.* 2014b).

Patients with OCD show deficits during planning (van den Heuvel *et al.* 2011; Shin *et al.* 2014), goal-directed learning (Gillan & Robbins, 2014; Voon *et al.* 2015), reward-based decision-making, gambling (Grassi *et al.* 2015; Figgie *et al.* 2016), and incentive processing (Figgie *et al.* 2011). Despite evidence that heightened impulsivity is a phenotype associated with OCD

(Benatti *et al.* 2014), only one (Sohn *et al.* 2014) of three TD studies in OCD (Vloet *et al.* 2010; Pinto *et al.* 2014; Sohn *et al.* 2014) found performance deficits.

Neuroimaging studies show that OCD is characterized by structural and functional abnormalities in medial and orbitofronto-striato-thalamo-cortical networks mediating EF (Menzies *et al.* 2008; Radua *et al.* 2010; Carlisi *et al.* 2016b; Norman *et al.* 2016). No fMRI studies, however, have investigated TD in OCD. Studies using other decision-making tasks in OCD have found hyperactivity in ventral-affective regions including ventromedial prefrontal, orbitofrontal and rostral anterior cingulate cortex (rACC) projecting to ventral striatum and mediodorsal thalamus, and hypoactivity in dorsal-cognitive cortico-striato-thalamic regions including dorsolateral prefrontal (DLPFC), temporal and parietal association cortex projecting to the dorsal striatum and caudate in patients relative to controls (Menzies *et al.* 2008; Brem *et al.* 2012). Hypoactivation in DLPFC and caudate has furthermore been shown in OCD patients during planning (van den Heuvel *et al.* 2005, 2011).

This suggests that ASD and OCD have abnormalities during planning and 'hot' EF tasks including reward-based decision-making, and that this may be underpinned by ventromedial and dorsolateral prefronto-striato-limbic abnormalities. However, it is unclear whether reward-based decision-making problems in both disorders are underpinned by shared trans-diagnostic mechanisms or by disorder-specific underlying abnormalities.

We hypothesized that adolescents with ASD would be more impaired on TD relative to adolescents with OCD and controls (Scott-Van Zeeland *et al.* 2010; Chantiluke *et al.* 2014b; Chen *et al.* 2016) and that both clinical groups compared with healthy controls would show underactivation in underlying ventromedial prefrontal, limbic and striatal regions mediating TD (Fineberg *et al.* 2009), reflecting a trans-diagnostic neuro-functional phenotype (Chantiluke *et al.* 2015a; Grassi *et al.* 2015; Chen *et al.* 2016). However, we hypothesized that people with OCD would show disorder-specific (ventro)medial and dorsolateral-prefrontal dysfunction (Menzies *et al.* 2008; Carlisi *et al.* 2016b; Norman *et al.* 2016) while ASD adolescents would show disorder-specific insular and temporo-parietal dysfunction compared to controls (Di Martino *et al.* 2009; Chantiluke *et al.* 2014b; Carlisi *et al.* 2016b).

Methods

Participants

Sixty-nine right-handed (Oldfield, 1971) boys (20 controls, 29 boys with ASD, 20 boys with OCD), 11–17 years, IQ \geq 70 (Wechsler, 1999) participated. Medication-

naïve boys with high-functioning ASD were recruited from local clinics and support-groups. ASD diagnosis was made by a consultant psychiatrist using ICD-10 research diagnostic criteria (WHO, 1992) and confirmed with the Autism Diagnostic Interview-Revised [ADI-R; (Lord *et al.* 1994)]. The ADI-R and the Autism Diagnostic Observation Schedule [ADOS; (Lord *et al.* 2000)] were completed for all ASD boys; all 29 reached autism cut-offs on all ADI-R (social/communication/restricted/stereotyped) and ADOS (communication/social) domains. ASD participants either fulfilled ICD-10 research diagnostic criteria for autism ($N=7$) or fulfilled these criteria but had no history of language delay and therefore were subtyped with Asperger's syndrome ($N=22$). Parents of ASD boys completed the Social Communication Questionnaire [SCQ; (Rutter *et al.* 2003)] and the Strengths and Difficulties Questionnaire [SDQ; (Goodman & Scott, 1999)] (see online Supplement). ASD participants had a physical examination to exclude comorbid medical disorders and biochemical, haematological and chromosomal abnormalities associated with ASD. None of the ASD individuals had a comorbid diagnosis of OCD or any psychiatric disorder, and none of the OCD patients had comorbid ASD.

OCD boys were recruited from National and Specialist OCD clinics. Diagnosis was made by a consultant psychiatrist using ICD-10 criteria and confirmed by the Children's Yale-Brown Obsessive-Compulsive Scale [CY-BOCS; (Goodman *et al.* 1989)]. Parents of OCD patients completed the SDQ. Patients with comorbid psychiatric or neurological disorders, including ASD, were not included in the OCD sample, although OCD patients were not specifically assessed for ASD. Four boys were prescribed stable doses of antidepressants (see online Supplement).

Twenty age and handedness-matched healthy controls were recruited locally by advertisement. Controls scored below clinical threshold on the SDQ and SCQ for any disorder and did not have any psychiatric condition.

Exclusion criteria for all participants included comorbid psychiatric or medical disorders affecting brain development (e.g. epilepsy/psychosis), drug/alcohol dependency, head injury, genetic conditions associated with ASD, abnormal structural brain scan and MRI contraindications. All controls also participated in previously published studies testing fluoxetine effects on TD in ADHD (Carlisi *et al.* 2016a) and neurofunctional maturation of TD in healthy adults and adolescents (Christakou *et al.* 2011); all but four ASD boys participated in our fMRI TD study comparing ASD and ADHD (Chantiluke *et al.* 2014b). Most ASD and control participants also participated in other fMRI tasks during their visit, published elsewhere (Christakou *et al.* 2013a, 2013b; Chantiluke *et al.* 2014a, 2015a, b; Murphy *et al.* 2014).




Fig. 1. Schematic of the temporal discounting fMRI paradigm. Subjects are asked to indicate whether they would prefer a small, variable amount of money immediately (immediate reward), or whether they would rather wait for a larger delay (up to £100) later (delayed reward). An algorithm adjusts the amount of the immediate reward offered based on the choices of the participant, so as to determine the lowest immediate reward they would tolerate before instead choosing to wait for the larger delayed reward. Three hypothetical delays are presented in random order: 1 week, 1 month and 1 year. Each delay choice is presented 20 times. Trials start with the presentation of the choice display, which remains available for 4 s, within which the subject must choose between the immediate (always on left side) and delayed (always on right) rewards. Total trial duration is 12 s.

This study was conducted in accordance with the Declaration of Helsinki. Ethical approval was obtained from the local Research Ethics Committee (05/Q0706/275). Study details were explained to child and guardian, and written informed consent was obtained for all participants.

TD paradigm

Prior to scanning, subjects practiced the 12-min TD task (Rubia *et al.* 2009; Christakou *et al.* 2011; Chantiluke *et al.* 2014b) in a mock-scanner. Subjects chose by pressing a left/right button with right index/middle-finger between receiving a small amount of money immediately (£0-£100) or receiving £100 in 1 week, month or year (Fig. 1). Delays (20 trials each) were randomized, but the delayed option (£100) was consistently displayed on the right side of the screen, and variable immediate choices on the left, minimizing sensorimotor mapping effects. Choices were displayed for 4 s, followed by a blank screen of at least 8 s (inter-trial-interval:12 s). The immediate reward amount was adjusted through an algorithm based on previous choices and calculated separately for each delay. This narrows the range of values, converging on an indifference point where the immediate reward is subjectively

considered equivalent to the delayed amount for the given delay (Rubia *et al.* 2009), ensuring comparable numbers of immediate and delayed choices for analysis.

Analysis of performance data

To estimate TD steepness for each subject, indifference values between the immediate amount and delayed £100 for each delay were calculated, equal to the participant's subjective value of £100 after each delay and defined as the midpoint between the lowest chosen immediate reward and the next lowest immediate reward available (i.e. the value of the immediate reward offered at which point the subject began to choose the delayed reward) (Christakou *et al.* 2011).

TD was measured using area under the curve (AUC) (Myerson *et al.* 2001). Smaller AUC denotes steeper discounting rates (i.e. increased choice-impulsivity) (see online Supplement).

One-way between-group analysis of variance (ANOVA) was conducted with AUC as dependent measure to examine group-differences.

fMRI image acquisition

Gradient-echo echo-planar imaging (EPI) data were acquired at King's College London on a 3T-General

Electric SIGNA HDx MRI scanner (Milwaukee, WI) using the body coil for radio frequency transmission and a quadrature birdcage head coil for reception. See online Supplement for acquisition parameters. Total scan was 1.5 h during which subjects completed 2–3 additional fMRI tasks.

fMRI image analysis

Event-related data were acquired in randomized trial presentation and analysed using the non-parametric XBAM package (v4.1) [www.brainmap.co.uk; (Brammer *et al.* 1997)]. The individual and group-level analysis methods are described in detail elsewhere (Brammer *et al.* 1997; Bullmore *et al.* 1999b; Cubillo *et al.* 2014) and in the online Supplement.

Briefly, fMRI data were realigned to minimize motion-related artefacts and smoothed using a 7.2 mm full-width-at-half-maximum (FWHM) Gaussian filter (Bullmore *et al.* 1999a). Time-series analysis of individual activation was performed with a wavelet-based resampling method (Bullmore *et al.* 2001). The main experimental conditions were convolved with 2 Poisson model functions (peaking at 4 and 8 s). The weighted sum of these convolutions giving the best fit (least-squares) to the time series at each voxel was calculated. A goodness-of-fit statistic (SSQ ratio) was then computed at each voxel consisting of the ratio of the sum of squares of deviations from the mean intensity value due to the model (fitted time series) divided by that of the squares due to the residuals (original minus model time series). This statistic, the SSQ ratio, was used in further analyses. Individual maps were then normalised to Talairach space (Talairach & Tournoux, 1988), and a group activation map was produced for each group.

ANCOVA of between-group effects

One-way between-group analysis of covariance (ANCOVA) with age as covariate was conducted using randomization-based testing to investigate case-control differences (Bullmore *et al.* 1999b, 2001). For these comparisons, statistical thresholds of 0.05 (voxel-level)/0.015 (cluster-level) were selected to obtain <1 false-positive 3D cluster per map. Standardized blood-oxygenation level-dependent (BOLD) responses were extracted from significant clusters for each participant and plotted to determine effect direction. *Post-hoc* significance was determined among pairwise comparisons using a one-way ANOVA.

Influence of behaviour, symptoms and medication

To examine whether clusters showing significant group effects were related to TD performance or symptoms,

BOLD response from these clusters was extracted for each participant and Spearman correlations (two-tailed) were performed with AUC and symptom subscales within each group. FMRI analyses were also repeated including AUC as covariate.

Lastly, analyses were repeated excluding the four OCD participants prescribed medication.

Results

Participants

There were no significant group-differences in age and IQ (Table 1). Multivariate ANOVAs showed group-differences on SDQ scores; *Post-hoc* tests revealed that patients had higher total-scores than controls, with ASD being more impaired than OCD patients (all $p < 0.001$). On the emotional-distress subscale, both patient groups were more impaired than controls ($p < 0.001$) but did not differ from each other. On all other SDQ subscales, ASD patients were significantly more impaired than controls and OCD patients (all $p < 0.005$), who did not differ on any measure, with the exception of the conduct subscale where ASD patients differed from controls only ($p < 0.001$).

Performance

AUC correlated inversely with k (as measured by the square-root transform of these values: $r = -0.555$, $p < 0.001$), suggesting adequate congruency between these two metrics. AUC differed between groups [controls: 0.56 ± 0.13 ; ASD: 0.45 ± 0.24 ; OCD: 0.59 ± 0.15 ; $F(2,66) = 4.04$, $p = 0.02$]. *Post-hoc* comparisons showed that ASD patients had significantly smaller AUC compared with controls ($p < 0.05$) and OCD patients ($p < 0.01$), indicating ASD patients discounted rewards more steeply than the other groups, who did not differ from each other.

fMRI data

Movement

Multivariate ANOVA showed no group-differences in mean head rotation [$F(2,66) = 1.17$, $p = \text{n.s.}$] or translation [$F(2,66) = 2.59$, $p = \text{n.s.}$] in 3-dimensional Euclidian space.

Group maps of brain activation for delayed-immediate choices

See online Supplement for maps of brain activation within each group for the contrast of delayed-immediate choices (online Supplementary Fig. S1).

Table 1. Participant characteristics for healthy control boys and patients with OCD or ASD

Variables	HC (N = 20) Mean (S.D.)	ASD (N = 29) Mean (S.D.)	OCD (N = 20) Mean (S.D.)	F test (df)	p value
Age (years)	15.29 (1.8)	14.72 (1.8)	15.74 (1.4)	2.22 (2,66)	0.12
IQ	118.90 (11.9)	113.17 (13.1)	117.70 (13.4)	1.38 (2,66)	0.26
SCQ total score	2.32 (2.3)	18.66 (8.1)	–	76.98 (1,47)	<0.001
SDQ total score	5.58 (4.2)	19.66 (6.8)	12.45 (5.6)	35.56 (2,66)	<0.001
SDQ emotional distress subscale	0.93 (1.8)	4.38 (2.9)	4.35 (2.6)	13.12 (2,66)	<0.001
SDQ conduct subscale	0.86 (1.1)	2.69 (2.2)	1.85 (1.5)	6.55 (2,66)	0.003
SDQ peer relations subscale	1.53 (1.7)	6.59 (2.3)	3.30 (3.0)	28.72 (2,66)	<0.001
SDQ hyperactive impulsive/inattentive subscale	2.72 (2.4)	5.93 (2.6)	2.95 (2.7)	12.52 (2,66)	<0.001
SDQ prosocial behaviour subscale	8.38 (2.4)	4.41 (2.4)	7.65 (2.6)	18.61 (2,66)	<0.001
ADOS communication score	–	3.62 (1.2)	–	–	–
ADOS social interaction score	–	9.03 (2.3)	–	–	–
ADOS communication + social	–	12.66 (3.1)	–	–	–
ADOS stereotypy score	–	1.52 (1.5)	–	–	–
ADI communication score	–	16.59 (4.7)	–	–	–
ADI social interaction score	–	19.97 (5.3)	–	–	–
ADI repetitive behaviour score	–	6.45 (2.4)	–	–	–
CY-BOCS total score	–	–	22.33 (5.8)	–	–
CY-BOCS – obsessions	–	–	10.79 (3.6)	–	–
CY-BOCS – compulsions	–	–	12.01 (3.1)	–	–

ADI, autism diagnostic interview; ADOS, autism diagnostic observation schedule; ASD, autism spectrum disorder; CY-BOCS, Children's Yale-Brown obsessive-compulsive symptoms checklist; HC, healthy controls; OCD, obsessive-compulsive disorder; SCQ, social communication questionnaire; SDQ, strengths and difficulties questionnaire.

Group-effects on brain activation

One-way ANOVA showed a significant group-effect for delayed-immediate choices in right ventromedial orbitofrontal cortex (vmOFC) extending into MPFC/lateral OFC/inferior frontal cortex (IFC), in cerebellum extending into occipital lobe/posterior cingulate (PCC)/precuneus, in rACC/vmPFC extending into left caudate, in left superior/middle temporal lobe (STL/MTL)/inferior parietal lobe (IPL) and in right MTL/STL extending into posterior insula/postcentral gyrus/IPL (Fig. 2a; Table 2). ANCOVA including AUC as covariate showed that effects in rACC/vmPFC and PCC/precuneus were related to task performance.

Post-hoc analyses based on extracted SSQs showed that abnormalities in vmOFC/MPFC/IFC were shared between OCD and ASD patients, who had increased activation to immediate-delayed choices relative to controls (both $p < 0.001$), who had more activation to delayed choices. In cerebellum/occipital lobe/PCC/precuneus, ASD and OCD patients had reduced activation to delayed-immediate choices compared with controls (both $p < 0.001$). In rACC/vmPFC/caudate, both patient groups had decreased activation to immediate-delayed choices relative to controls (ASD: $p < 0.001$; OCD: $p < 0.05$), who had enhanced activation to immediate-delayed choices, but this effect was more pronounced

in ASD *v.* OCD patients at trend-level ($p < 0.1$). Findings in right MTL/STL/insula/postcentral gyrus/IPL (all $p < 0.005$) and left STL/MTL/IPL were due to shared abnormalities in ASD ($p < 0.001$) and OCD ($p < 0.005$) patients, who had less activation to immediate-delayed choices relative to controls who activated this region for immediate *v.* delayed choices (Fig. 2b). When the four OCD patients prescribed medication were excluded from analyses, main findings remained, suggesting medication did not influence task-related activation.

Correlations between differentially activated brain regions and performance

Correlations between areas that differed between groups and AUC showed that greater activation to delayed-immediate choices in cerebellum/occipital lobe/PCC/precuneus was correlated with less-steep TD in the ASD ($r = 0.66$, $p < 0.001$) and OCD groups ($r = 0.45$, $p < 0.05$). Greater activation to immediate-delayed choices in left STL/IPL correlated with less-steep TD performance in the ASD group ($r = -0.41$, $p < 0.05$). In right MTL/STL/insula/postcentral gyrus/IPL, it correlated with better TD performance in both ASD ($r = -0.39$, $p < 0.05$) and OCD ($r = -0.59$, $p < 0.005$).




Fig. 2. Between-group activation differences for delayed minus immediate choices. (a) Axial slices showing split-plot analysis of variance (ANOVA) effects of group on brain activation to delayed – immediate choices. Talairach Z coordinates are indicated for slice distance (in mm) from the intercommissural line. The right side of the image corresponds to the right side of the brain. (b) Extracted statistical measures of BOLD response are shown for each of the three groups for each of the brain regions that showed a significant group effect. Black asterisks indicate a significant difference between controls and patient group. Red asterisk indicates a difference between the two patient groups. (*) = significant at a trend level; * = significant at the $p < 0.05$ level; ** = significant at the $p \leq 0.005$ level; *** = significant at the $p \leq 0.001$ level.

Table 2. Between-group activation differences for delayed minus immediate choices

Brain regions of activation difference	Brodman area (BA)	Peak Talairach coordinates (x, y, z)	Voxels	Cluster <i>p</i> value
(A) HC > OCD, ASD				
R vmOFC/MPFC/lateral OFC/IFC	47/11/25/10/46	40, 56, -13	189	0.009
PCC/precuneus/occipital lobe/cerebellum	31/7/19/18/17	-14, -89, 4	1060	0.0003
(B) OCD, ASD > HC				
rACC/vmPFC/left caudate	10/32/24	0, 41, 4	137	0.01
L STL/MTL/IPL	22/39/40/7/19	-51, -56, 9	273	0.005
R MTL/STL/posterior insula/postcentral gyrus/IPL	22/39/19/5/3/1/2/4/40/7	61, -22, 9	654	0.001

ASD, autism spectrum disorder; HC, healthy controls; IFC, inferior frontal cortex; IPL, inferior parietal lobe; L, left; MTL, middle temporal lobe; OCD, obsessive-compulsive disorder; OFC, orbitofrontal cortex; R, right; STL, superior temporal lobe; rACC, rostral anterior cingulate cortex; vmOFC, ventromedial orbitofrontal cortex; vmPFC, ventromedial prefrontal cortex.

Correlations between differentially activated brain regions and symptoms

In ASD boys, greater activation to delayed *v.* immediate choices in right vmOFC/MPFC/lateral OFC/IFC correlated at trend-level with lower symptom severity on the repetitive behaviour subscale of the ADI-R ($r = -0.34$, $p = 0.07$). In bilateral STL/insula, lower repetitive behaviour symptom severity was related to increased activation to immediate-delayed choices in the ASD group (left: $r = 0.47$, $p < 0.01$; right: $r = 0.42$, $p < 0.05$). In the OCD group, increased activation to delayed *v.* immediate choices in cerebellum/occipital lobe/PCC/precuneus correlated with lower symptom severity on the CY-BOCS compulsions subscale ($r = -0.58$, $p < 0.01$). There were no correlations between activation and other subscales from the CY-BOCS in OCD or ADOS/ADI-R in ASD.

Discussion

This comparison between ASD and OCD adolescents on a 'hot' EF measure of decision-making showed disorder-specific impaired TD in ASD relative to OCD boys and controls. Despite this, patients had predominantly shared neurofunctional deficits in key TD areas including vmOFC/MPFC/IFC, bilateral temporoparietal and cerebellar regions, suggesting that the neural basis of TD is a trans-diagnostic feature of both disorders. In ACC/vmPFC extending into caudate, ASD boys had trend-level more severe underactivation relative to OCD and controls for immediate *v.* delayed choices.

Disorder-specific performance impairment in ASD relative to OCD boys extends previous findings of impairments in ASD during TD (Chantiluke et al. 2014b), although there have been negative findings (Demurie et al. 2012). The absence of performance differences between OCD boys and controls is in line with

previous studies (Vloet et al. 2010; Pinto et al. 2014) [but see (Sohn et al. 2014)]. Moreover, ASD boys had elevated scores on the hyperactive-impulsive/inattention subscale of the SDQ compared with OCD boys and controls. The disorder-specific performance impairment in the ASD group may relate to these elevated impulsivity symptoms observed in ASD but not OCD, given that ADHD patients are consistently impaired in TD (Jackson & MacKillop, 2016). This finding exclusive to ASD lends support to the distinction between impulsive and compulsive behaviours (Robbins et al. 2012), suggesting that while both disorders exhibit deficits in top-down cognitive control and related circuitry (Dalley et al. 2011), ASD individuals exhibit more impulsive decision-making during TD, as evidenced by disorder-specific impairments and possibly supported by trend-level disorder-specific abnormalities in ACC/vmPFC/caudate, while OCD patients are more habitually compulsive, supported by intact choice behaviour and no disorder-specific abnormalities.

Both patient groups had reduced activation relative to controls to delayed-immediate choices in ventromedial and ventrolateral OFC/IFC. Ventromedial and ventrolateral fronto-limbic regions are key temporal foresight areas (Christakou et al. 2011; Peters & Büchel, 2011) thought to support calculation of discounted reward value. Moreover, right IFC is a key region for working memory, attention to time and integration of external information with internal value representations, supporting goal-directed EF and mediation of temporal foresight (Wittmann et al. 2007; Rubia et al. 2009; Carlisi et al. 2016a) and has previously been shown to be abnormal during reward-related decision-making in both OCD (Bari & Robbins, 2013; Stern & Taylor, 2014) and ASD (Dichter et al. 2012; Kohls et al. 2013).

Both patient groups showed reduced activation in PCC/precuneus/occipital lobe/cerebellum to delayed-

immediate choices compared with controls. These areas are important parts of fronto-limbic-parieto-cerebellar networks involved in motivation, reward evaluation and reward response (Vogt *et al.* 1992; McCoy *et al.* 2003). The cerebellum is typically activated during delayed choices in healthy populations and has been associated with future outcome expectancy and temporal bridging (Smith *et al.* 2003; Wittmann *et al.* 2007, 2010; Rubia *et al.* 2009; Christakou *et al.* 2011; Peters & Büchel, 2011; Noreika *et al.* 2013). We previously found similar effects in ADHD patients relative to controls during the same task, suggesting that cerebellar underactivation maybe a trans-diagnostic feature of disorders that are challenged in TD (Rubia *et al.* 2009). Moreover, given the aforementioned role of fronto-limbic-parieto-temporo-cerebellar networks in motivation and reward evaluation, shared abnormalities in this network could possibly relate to neurofunctional similarities in the motivational and reward salience of e.g. performing repetitive behaviours in each disorder, in line with theories of shared impairments in motivation control underpinning these behaviours in each disorder (Hollander *et al.* 2007). This collectively provides first evidence for shared functional abnormalities in ventromedial and ventrolateral fronto-parieto-striato-cerebellar regions between ASD and OCD.

Conversely, relative to controls, both patient groups had reduced activation to immediate choices in the rACC/vmPFC reaching into caudate. However, these abnormalities were at trend-level more pronounced in ASD relative to OCD, possibly linking to ASD-specific performance impairments. rACC mediates decision conflict (Pochon *et al.* 2008) and typically is increased in activation with decision difficulty during intertemporal choice (Pine *et al.* 2009). Our recent meta-analysis of structural and functional MRI studies also found shared reductions in this region in ASD and OCD relative to controls both in volume and in activation during cognitive control (Carlisi *et al.*, 2016b). In this study, however, we find that this dysfunction was trend-wise more impaired in ASD, implying a gradual rather than dichotomic effect of more severe impairment in ASD.

Findings of shared reduced vmPFC, left caudate, posterior insula and STL/IPL activation during immediate *v.* delayed choices in patients relative to controls are in line with a wealth of evidence implicating these regions in temporal foresight and reward-based decision-making as well as possible abnormal maturation of networks mediating these processes in ASD and OCD. We showed previously that vmPFC activation to immediate choices during TD increases with age and AUC, indicating an increase in delay-tolerant behaviour linked to increased limbic-cortico-striatal

activation with age (Christakou *et al.* 2013a). In children and adults, steeper TD has been associated with an imbalance between reduced activation in ventromedial prefrontal and lateral frontal systems mediating evaluation of future reward and temporal foresight, and reduced top-down control over ventral-striatal and limbic systems, which respond to immediate reward (Christakou *et al.* 2011; Peters & Büchel, 2011; Chantiluke *et al.* 2014b). Moreover, tasks indexing vmPFC functioning have shown age-dependent increases in sensitivity to future consequences (Crone & van der Molen, 2004) and behavioural control during TD (Steinbeis *et al.* 2016).

The caudate is involved in time discrimination (Smith *et al.* 2003), has been linked to reward expectation and evaluation (Hinvest *et al.* 2011) and is activated during immediate choices in healthy individuals (Christakou *et al.* 2011). In OCD, OFC-caudate loops are proposed to drive impulsivity as well as compulsive behaviour (Fineberg *et al.* 2009; Dalley *et al.* 2011). Thus, results could suggest that adolescents with ASD and OCD both have problems with context-dependent decision-making but that this is more problematic for people with ASD, potentially relating to the findings of disorder-specific behavioural deficits in the ASD group. Moreover, the posterior insula is associated with decision-making in the context of prior risk (Xue *et al.* 2010) and is important for the integration of temporal-affective information (Elliott *et al.* 2000) and temporal encoding (Wittmann *et al.* 2010). While previous studies have found specifically anterior insula activation during TD in children (Rubia *et al.* 2009) and adults (Tanaka *et al.* 2004; Bickel *et al.* 2009; Hinvest *et al.* 2011), the present results highlight a differential abnormality in the posterior insula during reward presentation and internal state evaluation (Elliott *et al.* 2000) shared between ASD and OCD.

Findings of reduced activation to immediate-delayed choices in STL/IPL in ASD relative to controls are in line with evidence of weaker brain-behaviour correlations in this region in ASD relative to controls during TD (Chantiluke *et al.* 2014b) and extend these findings to OCD. These regions are important for temporal coding and reward selection (Cardinal, 2006; Christakou *et al.* 2011), suggesting deficits with planning, consistent with behavioural deficits in this domain in ASD (Hill, 2004) and OCD (Shin *et al.* 2014). IPL is specifically sensitive to delay (Rubia *et al.* 1998) and attention-allocation to time (Ortuno *et al.* 2002; Coull, 2004; Rubia, 2006), as well as duration encoding (Wittmann, 2009) and quantity representation, which may contribute to inter-temporal choices regarding the IPL's role in comparing time and value (Sandrini *et al.* 2004). Correlations between enhanced activation to immediate choices in the

patient groups and better TD performance suggest that in both groups, this upregulation is related to a shift in performance towards that of controls, providing possible mechanistic implications of this region in the context of TD behaviour. Moreover, increased activation bilaterally in this region in the ASD group correlated with lower levels of repetitive behaviours, linking performance improvement and symptom reduction to brain activation in these individuals, further highlighting the mechanistic implications of this region in the context of repetitive behaviours and decision-making.

Clinically, the fact that these disorders exhibit shared neural abnormalities during TD has implications for identification of common mechanisms, which may drive overlapping behaviours in each disorder. While symptoms such as compulsions in OCD can sometimes appear similar to repetitive behaviours in ASD at an observational level, less is known about the mechanistic underpinnings of these behaviours and related cognitive functions and whether they are shared or disorder-specific. Thus, this evidence sheds light on trans-diagnostic phenotypes that could aid in future treatment targets and work toward providing a biological explanation of commonalities and differences in clinical behaviour. This has similarly been shown in the case of inhibitory control and brain structure/function differences/similarities in a recent meta-analysis comparing ASD and OCD (Carlisi et al. 2016b), and this study extends this understanding to temporal foresight and decision-making.

This study's strengths include the thoroughness with which ASD individuals were assessed for the presence of ASD-related symptomatology and the exclusion of patients with psychiatric comorbidities. However, sub-threshold symptoms may have been present in the patient samples. The group of ASD patients tested in this study had a relatively high IQ, comparable with that of controls. While matching groups for IQ is important for fMRI studies to disentangle the effects of ASD from the effects of low IQ, this also means that the findings are not generalizable to other more typical ASD patients with low IQ (Charman et al. 2011; Crespi, 2016). The fact that most patients had high-functioning Asperger's syndrome further limits generalizability. Thus, it is possible that OCD-related symptoms were present in the ASD sample and could account for some of the neurobiological overlap in results. In addition, sub-clinical levels of ASD-related symptoms may have been present in the OCD sample, as reflected by shared impairments compared with controls on the emotional-distress SDQ subscale. It would also be interesting to examine the possible effects of puberty on any observed abnormalities. However, it has been shown that impulsive behaviour is independent of puberty in males

(Steinberg et al. 2008). Additionally, four OCD patients were prescribed antidepressant medication. While there is evidence for effects of serotonin on brain function (Murphy et al. 2008; Murphy, 2010), results remained when analyses were repeated excluding these patients. Lastly, it is a common finding that brain activation is more sensitive than performance to detect differences between groups in these patient groups (Fitzgerald et al. 2010; Duerden et al. 2013; Ambrosino et al. 2014; Marsh et al. 2014; Chantiluke et al. 2015b; Morein-Zamir et al. 2015). While the subject numbers have been shown to be sufficient for fMRI analyses (Thirion et al. 2007), the performance and correlation analyses, however, were underpowered.

Conclusions

This is the first study to compare brain function between these disorders and provides novel evidence to suggest that ASD and OCD share trans-diagnostic abnormalities during TD in ventromedial and ventrolateral fronto-striatal and fronto-temporo-parieto-cerebellar regions important for temporal foresight and reward-related decision-making. This may drive shared problems with reward-related behaviours and delaying repetitive actions.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717001088>.

Acknowledgements

This work was supported by grants from the Medical Research Council (MRC G0300155) to K. R. and the MRC UK Autism Imaging Multicentre Study (G0400061) to D. M. This paper represents independent research part-funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. A. C. was supported by a post-doctoral fellowship from MRC G0300155. K.C. and L.N. were supported by Ph.D. studentships from the Institute of Psychiatry, Psychology and Neuroscience, King's College London. C.C. was supported by a NIHR-BRC Ph.D. studentship.

Declaration Interest

K. R. has received funding from Lilly for another project and speaker's honoraria from Lilly, Shire, Novartis

and Medice. D. M. has received funding for another project from Lilly. M. B. has served as a consultant for PIVital. The other authors have no conflict of interests to declare.

Ethical standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Abramovitch A, Abramowitz JS, Mittelman A** (2013). The neuropsychology of adult obsessive-compulsive disorder: a meta-analysis. *Clinical Psychology Review* **33**, 1163–1171.
- Ambrosino S, Bos D, Van Raalten T, Kobussen N, Van Belle J, Oranje B, Durston S** (2014). Functional connectivity during cognitive control in children with autism spectrum disorder: an independent component analysis. *Journal of Neural Transmission* **121**, 1145–1155.
- American Psychiatric Association** (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th edn. American Psychiatric Publishing: Arlington, VA.
- Antrop I, Stock P, Verté S, Wiersema JR, Baeyens D, Roeyers H** (2006). ADHD and delay aversion: the influence of non-temporal stimulation on choice for delayed rewards. *Journal of Child Psychology and Psychiatry* **47**, 1152–1158.
- Bari A, Robbins TW** (2013). Inhibition and impulsivity: behavioral and neural basis of response control. *Progress in Neurobiology* **108**, 44–79.
- Benatti B, Dell'osso B, Arici C, Hollander E, Altamura AC** (2014). Characterizing impulsivity profile in patients with obsessive-compulsive disorder. *International Journal of Psychiatry in Clinical Practice* **18**, 156–160.
- Bickel WK, Pitcock JA, Yi R, Angtuaco EJC** (2009). Congruence of BOLD response across intertemporal choice conditions: fictive and real money gains and losses. *Journal of Neuroscience* **29**, 8839–8846.
- Blumberg SJ, Bramlett MD, Kogan MD, Schieve LA, Jones JR, Lu MC** (2013). Changes in prevalence of parent-reported autism spectrum disorder in school-aged US children: 2007 to 2011–2012. *National Health Statistics Reports* **65**, 1–7.
- Brammer MJ, Bullmore ET, Simmons A, Williams SCR, Grasby PM, Howard RJ, Woodruff PWR, Rabe-Hesketh S** (1997). Generic brain activation mapping in functional magnetic resonance imaging: a nonparametric approach. *Magnetic Resonance Imaging* **15**, 763–770.
- Brem S, Hauser TU, Iannaccone R, Brandeis D, Drechsler R, Walitza S** (2012). Neuroimaging of cognitive brain function in paediatric obsessive compulsive disorder: a review of literature and preliminary meta-analysis. *Journal of Neural Transmission* **119**, 1425–1448.
- Bullmore E, Brammer M, Rabe-Hesketh S, Curtis V, Morris R, Williams S, Sharma T, McGuire P** (1999a). Methods for diagnosis and treatment of stimulus-correlated motion in generic brain activation studies using fMRI. *Human Brain Mapping* **7**, 38–48.
- Bullmore E, Long C, Suckling J, Fadili J, Calvert G, Zelaya F, Carpenter TA, Brammer M** (2001). Colored noise and computational inference in neurophysiological (fMRI) time series analysis: resampling methods in time and wavelet domains. *Human Brain Mapping* **12**, 61–78.
- Bullmore ET, Suckling J, Overmeyer S, Rabe-Hesketh S, Taylor E, Brammer MJ** (1999b). Global, voxel, and cluster tests, by theory and permutation, for a difference between two groups of structural MR images of the brain. *IEEE Transactions on Medical Imaging* **18**, 32–42.
- Cardinal RN** (2006). Neural systems implicated in delayed and probabilistic reinforcement. *Neural Networks* **19**, 1277–1301.
- Carlisi CO, Chantiluke K, Norman L, Christakou A, Barrett N, Giampietro V, Brammer M, Simmons A, Rubia K** (2016a). The effects of acute fluoxetine administration on temporal discounting in youth with ADHD. *Psychological Medicine* **46**, 1197–1209.
- Carlisi CO, Norman LJ, Lukito SS, Radua J, Mataix-Cols D, Rubia K** (2016b). Comparative multimodal meta-analysis of structural and functional brain abnormalities in autism spectrum disorder and obsessive-compulsive disorder. *Biological Psychiatry*. doi: 10.1016/j.biopsych.2016.10.006.
- Chantiluke K, Barrett N, Giampietro V, Brammer M, Simmons A, Murphy DG, Rubia K** (2015a). Inverse effect of fluoxetine on medial prefrontal cortex activation during reward reversal in ADHD and autism. *Cerebral Cortex* **25**, 1757–1770.
- Chantiluke K, Barrett N, Giampietro V, Brammer M, Simmons A, Rubia K** (2014a). Disorder-dissociated effects of fluoxetine on brain function of working memory in attention deficit hyperactivity disorder and autism spectrum disorder. *Psychological Medicine* **45**, 1195–1205.
- Chantiluke K, Barrett N, Giampietro V, Santosh P, Brammer M, Simmons A, Murphy D, Rubia K** (2015b). Inverse fluoxetine effects on inhibitory brain activation in non-comorbid boys with ADHD and with ASD. *Psychopharmacology* **232**, 2071–2082.
- Chantiluke K, Christakou A, Murphy CM, Giampietro V, Daly EM, Brammer M, Murphy DG, Rubia K** (2014b). Disorder-specific functional abnormalities during temporal discounting in youth with attention deficit hyperactivity disorder (ADHD), Autism and comorbid ADHD and Autism. *Psychiatry Research: Neuroimaging* **223**, 113–120.
- Charman T, Pickles A, Simonoff E, Chandler S, Loucas T, Baird G** (2011). IQ in children with autism spectrum disorders: data from the special needs and autism project (SNAP). *Psychological Medicine* **41**, 619–627.
- Chen S, Chien Y, Wu C, Shang C, Wu Y, Gau S** (2016). Deficits in executive functions among youths with autism spectrum disorders: an age-stratified analysis. *Psychological Medicine* **46**, 1625–1638.
- Christakou A, Brammer M, Rubia K** (2011). Maturation of limbic corticostriatal activation and connectivity associated

- with developmental changes in temporal discounting. *NeuroImage* **54**, 1344–1354.
- Christakou A, Gershman SJ, Niv Y, Simmons A, Brammer M, Rubia K** (2013a). Neural and psychological maturation of decision-making in adolescence and young adulthood. *Journal of Cognitive Neuroscience* **25**, 1807–1823.
- Christakou A, Murphy C, Chantiluke K, Cubillo A, Smith A, Giampietro V, Daly E, Ecker C, Robertson D, Murphy D** (2013b). Disorder-specific functional abnormalities during sustained attention in youth with attention deficit hyperactivity disorder (ADHD) and with Autism. *Molecular Psychiatry* **18**, 236–244.
- Coull JT** (2004). fMRI studies of temporal attention: allocating attention within, or towards, time. *Cognitive Brain Research* **21**, 216–226.
- Crespi BJ** (2016). Autism as a disorder of high intelligence. *Frontiers in Neuroscience* **10**, 300. <http://doi.org/10.3389/fnins.2016.00300>
- Crone EA, Van Der Molen MW** (2004). Developmental changes in real life decision making: performance on a gambling task previously shown to depend on the ventromedial prefrontal cortex. *Developmental Neuropsychology* **25**, 251–279.
- Cubillo A, Smith AB, Barrett N, Giampietro V, Brammer MJ, Simmons A, Rubia K** (2014). Shared and drug-specific effects of atomoxetine and methylphenidate on inhibitory brain dysfunction in medication-naïve ADHD boys. *Cerebral Cortex* **24**, 174–185.
- Dalley JW, Everitt BJ, Robbins TW** (2011). Impulsivity, compulsivity, and top-down cognitive control. *Neuron* **69**, 680–694.
- Demurie E, Roeyers H, Baeyens D, Sonuga-Barke E** (2012). Temporal discounting of monetary rewards in children and adolescents with ADHD and autism spectrum disorders. *Developmental Science* **15**, 791–800.
- Demurie E, Roeyers H, Baeyens D, Sonuga-Barke E** (2013). Domain-general and domain-specific aspects of temporal discounting in children with ADHD and autism spectrum disorders (ASD): a proof of concept study. *Research in Developmental Disabilities* **34**, 1870–1880.
- Dichter GS, Richey JA, Rittenberg AM, Sabatino A, Bodfish JW** (2012). Reward circuitry function in autism during face anticipation and outcomes. *Journal of Autism and Developmental Disorders* **42**, 147–160.
- Di Martino A, Ross K, Uddin LQ, Sklar AB, Castellanos FX, Milham MP** (2009). Functional brain correlates of social and nonsocial processes in autism spectrum disorders: an activation likelihood estimation meta-analysis. *Biological Psychiatry* **65**, 63–74.
- Doshi-Velez F, Ge Y, Kohane I** (2014). Comorbidity clusters in autism spectrum disorders: an electronic health record time-series analysis. *Pediatrics* **133**, e54–e63.
- Duerden EG, Taylor MJ, Soorya LV, Wang T, Fan J, Anagnostou E** (2013). Neural correlates of inhibition of socially relevant stimuli in adults with autism spectrum disorder. *Brain Research* **1533**, 80–90.
- Elliott R, Friston KJ, Dolan RJ** (2000). Dissociable neural responses in human reward systems. *Journal of Neuroscience* **20**, 6159–6165.
- Fiege M, Pattij T, Willuhn I, Luigjes J, Van Den Brink W, Goudriaan A, Potenza MN, Robbins TW, Denys D** (2016). Compulsivity in obsessive-compulsive disorder and addictions. *European Neuropsychopharmacology* **26**, 856–868.
- Fiege M, Vink M, De Geus F, Vulink N, Veltman DJ, Westenberg H, Denys D** (2011). Dysfunctional reward circuitry in obsessive-compulsive disorder. *Biological Psychiatry* **69**, 867–874.
- Fineberg NA, Potenza MN, Chamberlain SR, Berlin HA, Menzies L, Bechara A, Sahakian BJ, Robbins TW, Bullmore ET, Hollander E** (2009). Probing compulsive and impulsive behaviors, from animal models to endophenotypes: a narrative review. *Neuropsychopharmacology* **35**, 591–604.
- Fitzgerald KD, Stern ER, Angstadt M, Nicholson-Muth KC, Maynor MR, Welsh RC, Hanna GL, Taylor SF** (2010). Altered function and connectivity of the medial frontal cortex in pediatric obsessive-compulsive disorder. *Biological Psychiatry* **68**, 1039–1047.
- Geurts HM, Verté S, Oosterlaan J, Roeyers H, Sergeant JA** (2004). How specific are executive functioning deficits in attention deficit hyperactivity disorder and autism? *Journal of Child Psychology and Psychiatry* **45**, 836–854.
- Gillan CM, Robbins TW** (2014). Goal-directed learning and obsessive-compulsive disorder. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**, 20130475. <http://doi.org/10.1098/rstb.2013.0475>.
- Goodman R, Scott S** (1999). Comparing the strengths and difficulties questionnaire and the child behavior checklist: is small beautiful? *Journal of Abnormal Child Psychology* **27**, 17–24.
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Delgado P, Heninger GR, Charney DS** (1989). The yale-brown obsessive compulsive scale: II. validity. *Archives of General Psychiatry* **46**, 1012–1016.
- Grassi G, Pallanti S, Righi L, Fiege M, Mantione M, Denys D, Piccagliani D, Rossi A, Stratta P** (2015). Think twice: impulsivity and decision making in obsessive-compulsive disorder. *Journal of Behavioral Addictions* **4**, 263–272.
- Hill EL** (2004). Executive dysfunction in autism. *Trends in Cognitive Sciences* **8**, 26–32.
- Hinvest NS, Elliott R, Mckie S, Anderson IM** (2011). Neural correlates of choice behavior related to impulsivity and venturesomeness. *Neuropsychologia* **49**, 2311–2320.
- Hollander E, Kim S, Khanna S, Pallanti S** (2007). Obsessive-compulsive disorder and obsessive-compulsive spectrum disorders: diagnostic and dimensional issues. *CNS Spectrums* **12**(S3), 5–13.
- Jackson JNS, Mackillop J** (2016). Attention-deficit/hyperactivity disorder and monetary delay discounting: a meta-analysis of case-control studies. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **1**, 316–325.
- Just MA, Cherkassky VL, Keller TA, Kana RK, Minshew NJ** (2007). Functional and anatomical cortical underconnectivity in autism: evidence from an fMRI study of an executive function task and corpus callosum morphometry. *Cerebral Cortex* **17**, 951–961.
- Kohls G, Schulte-Rüther M, Nehr Korn B, Müller K, Fink GR, Kamp-Becker I, Herpertz-Dahlmann B, Schultz RT,**

- Konrad K** (2013). Reward system dysfunction in autism spectrum disorders. *Social Cognitive and Affective Neuroscience* **8**, 565–572.
- Kriete T, Noelle DC** (2015). Dopamine and the development of executive dysfunction in autism spectrum disorders. *PLoS ONE* **10**, e0121605.
- Langen M, Durston S, Kas MJ, Van Engeland H, Staal WG** (2011). The neurobiology of repetitive behavior: . . . and men. *Neuroscience and Biobehavioral Reviews* **35**, 356–365.
- Lord C, Risi S, Lambrecht L, Cook Jr. E, Leventhal B, Dilavore P, Pickles A, Rutter M** (2000). The autism diagnostic observation schedule – generic: a standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders* **30**, 205–223.
- Lord C, Rutter M, Couteur A** (1994). Autism diagnostic interview-revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders* **24**, 659–685.
- Marsh R, Horga G, Parashar N, Wang Z, Peterson BS, Simpson HB** (2014). Altered activation in fronto-striatal circuits during sequential processing of conflict in unmedicated adults with obsessive-compulsive disorder. *Biological Psychiatry* **75**, 615–622.
- Mccoy AN, Crowley JC, Haghigian G, Dean HL, Platt ML** (2003). Saccade reward signals in posterior cingulate cortex. *Neuron* **40**, 1031–1040.
- Menon V, Uddin LQ** (2010). Saliency, switching, attention and control: a network model of insula function. *Brain Structure and Function* **214**, 655–667.
- Menzies L, Chamberlain SR, Laird AR, Thelen SM, Sahakian BJ, Bullmore ET** (2008). Integrating evidence from neuroimaging and neuropsychological studies of obsessive-compulsive disorder: the orbitofronto-striatal model revisited. *Neuroscience and Biobehavioral Reviews* **32**, 525–549.
- Mesulam M-M** (1999). Spatial attention and neglect: parietal, frontal and cingulate contributions to the mental representation and attentional targeting of salient extrapersonal events. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **354**, 1325–1346.
- Morein-Zamir S, Voon V, Dodds CM, Sule A, Van Niekkerk J, Sahakian BJ, Robbins TW** (2015). Divergent subcortical activity for distinct executive functions: stopping and shifting in obsessive compulsive disorder. *Psychological Medicine FirstView* 1–12.
- Murphy CM, Christakou A, Daly EM, Ecker C, Giampietro V, Brammer M, Smith AB, Johnston P, Robertson DM, Consortium MA, Murphy DG, Rubia K** (2014). Abnormal functional activation and maturation of fronto-striato-temporal and cerebellar regions during sustained attention in autism spectrum disorder. *American Journal of Psychiatry* **171**, 1107–1116.
- Murphy SE** (2010). Using functional neuroimaging to investigate the mechanisms of action of selective serotonin reuptake inhibitors (SSRIs). *Current Pharmaceutical Design* **16**, 1990–1997.
- Murphy SE, Longhitano C, Ayres RE, Cowen PJ, Harmer CJ, Rogers RD** (2008). The role of serotonin in nonnormative risky choice: the effects of tryptophan supplements on the ‘reflection effect’ in healthy adult volunteers. *Journal of Cognitive Neuroscience* **21**, 1709–1719.
- Myerson J, Green L, Warusawitharana M** (2001). Area under the curve as a measure of discounting. *Journal of the Experimental Analysis of Behavior* **76**, 235–243.
- Noreika V, Falter CM, Rubia K** (2013). Timing deficits in attention-deficit/hyperactivity disorder (ADHD): evidence from neurocognitive and neuroimaging studies. *Neuropsychologia* **51**, 235–266.
- Norman LJ, Carlisi C, Lukito S, Hart H, Mataix-Cols D, Radua J, Rubia K** (2016). Structural and functional brain abnormalities in attention-deficit/hyperactivity disorder and obsessive-compulsive disorder: a comparative meta-analysis. *JAMA Psychiatry* **73**, 815–825.
- Odum AL** (2011). Delay discounting: trait variable? *Behavioural Processes* **87**, 1–9.
- Oldfield RC** (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* **9**, 97–113.
- Ortuno F, Ojeda N, Arbizu J, Lopez P, Marti-Climent J, Penuelas I, Cervera S** (2002). Sustained attention in a counting task: normal performance and functional neuroanatomy. *NeuroImage* **17**, 411–420.
- Ozonoff S, Jensen J** (1999). Brief report: specific executive function profiles in three neurodevelopmental disorders. *Journal of Autism and Developmental Disorders* **29**, 171–177.
- Paulus MP, Rogalsky C, Simmons A, Feinstein JS, Stein MB** (2003). Increased activation in the right insula during risk-taking decision making is related to harm avoidance and neuroticism. *NeuroImage* **19**, 1439–1448.
- Peters J, Büchel C** (2010). Episodic future thinking reduces reward delay discounting through an enhancement of prefrontal-mediotemporal interactions. *Neuron* **66**, 138–148.
- Peters J, Büchel C** (2011). The neural mechanisms of inter-temporal decision-making: understanding variability. *Trends in Cognitive Sciences* **15**, 227–239.
- Philip RCM, Dauvermann MR, Whalley HC, Baynham K, Lawrie SM, Stanfield AC** (2012). A systematic review and meta-analysis of the fMRI investigation of autism spectrum disorders. *Neuroscience and Biobehavioral Reviews* **36**, 901–942.
- Pine A, Seymour B, Roiser JP, Bossaerts P, Friston KJ, Curran HV, Dolan RJ** (2009). Encoding of Marginal Utility across Time in the Human Brain. *Journal of Neuroscience* **29**, 9575–9581.
- Pinto A, Steinglass JE, Greene AL, Weber EU, Simpson HB** (2014). Capacity to delay reward differentiates obsessive-compulsive disorder and obsessive compulsive personality disorder. *Biological Psychiatry* **75**, 653–659.
- Pochon J-B, Riis J, Sanfey AG, Nystrom LE, Cohen JD** (2008). Functional imaging of decision conflict. *Journal of Neuroscience* **28**, 3468–3473.
- Radua J, Pozo NOD, Gómez J, Guillen-Grima F, Ortuno F** (2014). Meta-analysis of functional neuroimaging studies indicates that an increase of cognitive difficulty during executive tasks engages brain regions associated with time perception. *Neuropsychologia* **58**, 14–22.
- Radua J, Van Den Heuvel OA, Surguladze S, Mataix-Cols D** (2010). Meta-analytical comparison of voxel-based

- morphometry studies in obsessive-compulsive disorder vs other anxiety disorders. *Archives of General Psychiatry* **67**, 701–711.
- Richards JB, Zhang L, Mitchell SH, Wit H** (1999). Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *Journal of the Experimental Analysis of Behavior* **71**, 121–143.
- Robbins TW, Gillan CM, Smith DG, De Wit S, Ersche KD** (2012). Neurocognitive endophenotypes of impulsivity and compulsivity: towards dimensional psychiatry. *Trends in Cognitive Sciences* **16**, 81–91.
- Rubia K** (2006). The neural correlates of timing functions. In *Timing the Future: the Case for a Time-Based Prospective Memory* (ed. J. Glicksohn), pp. 213–238. World Scientific Publishing: Hackensack, NJ.
- Rubia K, Halari R, Christakou A, Taylor E** (2009). Impulsiveness as a timing disturbance: neurocognitive abnormalities in attention-deficit hyperactivity disorder during temporal processes and normalization with methylphenidate. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**, 1919–1931.
- Rubia K, Overmeyer S, Taylor E, Brammer M, Williams S, Simmons A, Andrew C, Bullmore E** (1998). Prefrontal involvement in 'temporal bridging' and timing movement. *Neuropsychologia* **36**, 1283–1293.
- Ruscio A, Stein D, Chiu W, Kessler R** (2010). The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Molecular Psychiatry* **15**, 53–63.
- Rutter M, Bailey A, Lord C** (2003). *The Social Communication Questionnaire: Manual*. Western Psychological Services: Los Angeles.
- Sanders J, Johnson KA, Garavan H, Gill M, Gallagher L** (2008). A review of neuropsychological and neuroimaging research in autistic spectrum disorders: attention, inhibition and cognitive flexibility. *Research in Autism Spectrum Disorders* **2**, 1–16.
- Sandrini M, Rossini PM, Miniussi C** (2004). The differential involvement of inferior parietal lobule in number comparison: a rTMS study. *Neuropsychologia* **42**, 1902–1909.
- Schmitz N, Rubia K, Van Amelsvoort T, Daly E, Smith A, Murphy DGM** (2008). Neural correlates of reward in autism. *British Journal of Psychiatry* **192**, 19–24.
- Scott-Van Zeeland AA, Dapretto M, Ghahremani DG, Poldrack RA, Bookheimer SY** (2010). Reward processing in autism. *Autism Research* **3**, 53–67.
- Shin NY, Lee TY, Kim E, Kwon JS** (2014). Cognitive functioning in obsessive-compulsive disorder: a meta-analysis. *Psychological Medicine* **44**, 1121–1130.
- Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, Baird G** (2008). Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child and Adolescent Psychiatry* **47**, 921–929.
- Singer T, Critchley HD, Preuschoff K** (2009). A common role of insula in feelings, empathy and uncertainty. *Trends in Cognitive Sciences* **13**, 334–340.
- Smith A, Taylor E, Lidzba K, Rubia K** (2003). A right hemispheric frontocerebellar network for time discrimination of several hundreds of milliseconds. *NeuroImage* **20**, 344–350.
- Sohn SY, Kang JI, Namkoong K, Kim SJ** (2014). Multidimensional measures of impulsivity in obsessive-compulsive disorder: cannot wait and stop. *PLoS ONE* **9**, e111739.
- Steinbeis N, Haushofer J, Fehr E, Singer T** (2016). Development of behavioral control and associated vmPFC–DLPFC connectivity explains children's increased resistance to temptation in intertemporal choice. *Cerebral Cortex* **26**, 32–42.
- Steinberg L, Albert D, Cauffman E, Banich M, Graham S, Woolard J** (2008). Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: evidence for a dual systems model. *Developmental Psychology* **44**, 1764–1778.
- Stern ER, Taylor SF** (2014). Cognitive neuroscience of obsessive-compulsive disorder. *Psychiatric Clinics of North America* **37**, 337–352.
- Talairach J, Tournoux P** (1988). *Coplanar Stereotaxic Atlas of the Human Brain, a 3-dimensional Proportional System: an Approach to Cerebral Imaging*. Thieme: New York.
- Tanaka SC, Doya K, Okada G, Ueda K, Okamoto Y, Yamawaki S** (2004). Prediction of immediate and future rewards differentially recruits cortico-basal ganglia loops. *Nature Neuroscience* **7**, 887–893.
- Thirion B, Pinel P, Mériaux S, Roche A, Dehaene S, Poline J-B** (2007). Analysis of a large fMRI cohort: statistical and methodological issues for group analyses. *NeuroImage* **35**, 105–120.
- Van Den Heuvel OA, Mataix-Cols D, Zwieter G, Cath DC, Van Der Werf YD, Groenewegen HJ, Van Balkom AJLM, Veltman DJ** (2011). Common limbic and frontal-striatal disturbances in patients with obsessive compulsive disorder, panic disorder and hypochondriasis. *Psychological Medicine* **41**, 2399–2410.
- van den Heuvel OA, Veltman DJ, Groenewegen HJ, Cath DC, van Balkom AJ, van Hartkamp J, Barkhof F, van Dyck R** (2005). Frontal-striatal dysfunction during planning in obsessive-compulsive disorder. *Archives of General Psychiatry* **62**, 301–309.
- Via E, Radua J, Cardoner N, Happé F, Mataix-Cols D** (2011). Meta-analysis of gray matter abnormalities in autism spectrum disorder: should asperger disorder be subsumed under a broader umbrella of autistic spectrum disorder? *Archives of General Psychiatry* **68**, 409–418.
- Vloet T, Marx I, Kahraman-Lanzerath B, Zepf F, Herpertz-Dahlmann B, Konrad K** (2010). Neurocognitive performance in children with ADHD and OCD. *Journal of Abnormal Child Psychology* **38**, 961–969.
- Vogt BA, Finch DM, Olson CR** (1992). Functional heterogeneity in cingulate cortex: the anterior executive and posterior evaluative regions. *Cerebral Cortex* **2**, 435–443.
- Voon V, Derbyshire K, Ruck C, Irvine MA, Worbe Y, Enander J, Schreiber LRN, Gillan C, Fineberg NA, Sahakian BJ, Robbins TW, Harrison NA, Wood J, Daw ND, Dayan P, Grant JE, Bullmore ET** (2015). Disorders of compulsivity: a common bias towards learning habits. *Molecular Psychiatry* **20**, 345–352.

- Wechsler D** (1999). *Wechsler Abbreviated Scale of Intelligence*. Psychological Corporation: San Antonio, TX.
- Wesley MJ, Bickel WK** (2014). Remember the future II: meta-analyses and functional overlap of working memory and delay discounting. *Biological Psychiatry* **75**, 435–448.
- WHO** (1992). *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. World Health Organization: Geneva.
- Wiener M, Turkeltaub P, Coslett HB** (2010). The image of time: a voxel-wise meta-analysis. *NeuroImage* **49**, 1728–1740.
- Wittmann M** (2009). The inner experience of time. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **364**, 1955–1967.
- Wittmann M, Leland DS, Paulus MP** (2007). Time and decision making: differential contribution of the posterior insular cortex and the striatum during a delay discounting task. *Experimental Brain Research* **179**, 643–653.
- Wittmann M, Simmons AN, Aron JL, Paulus MP** (2010). Accumulation of neural activity in the posterior insula encodes the passage of time. *Neuropsychologia* **48**, 3110–3120.
- Xue G, Lu Z, Levin IP, Bechara A** (2010). The impact of prior risk experiences on subsequent risky decision-making: the role of the insula. *NeuroImage* **50**, 709–716.
- Zelazo PD, Müller U** (2007). Executive function in typical and atypical development. In *Blackwell Handbook of Childhood Cognitive Development* (ed. U. Goswami), pp. 445–469. Blackwell Publishers Ltd: Malden, MA.