

A micronised, dispersible ferric pyrophosphate with high relative bioavailability in man

Meredith C. Fidler¹, Thomas Walczyk¹, Lena Davidsson^{1*}, Christophe Zeder¹, Noboru Sakaguchi², Lekh R. Juneja² and Richard F. Hurrell¹

¹Laboratory for Human Nutrition, Institute of Food Science and Nutrition, Swiss Federal Institute of Technology (ETH) Zurich, PO Box 474/Seestrasse 72, CH-8803 Rueschlikon, Switzerland

²Nutritional Foods Division, Taiyo Kagaku, 9-5 Akahori-Shinmachi, Yokkaichi, Mie 510-0825, Japan

(Received 14 May 2003 – Revised 4 September 2003 – Accepted 10 September 2003)

Ferric pyrophosphate is a water-insoluble Fe compound used to fortify infant cereals and chocolate-drink powders as it causes no organoleptic changes to the food vehicle. However, it is only of low absorption in man. Recently, an innovative ferric pyrophosphate has been developed (Sunactive Fe™) based on small-particle-size ferric pyrophosphate (average size 0.3 µm) mixed with emulsifiers, so that it remains in suspension in liquid products. The aim of the present studies was to compare Fe absorption of micronised, dispersible ferric pyrophosphate (Sunactive Fe™) with that of ferrous sulfate in an infant cereal and a yoghurt drink. Two separate Fe absorption studies were made in adult women (ten women/study). Fe absorption was based on the erythrocyte incorporation of stable isotopes (⁵⁷Fe and ⁵⁸Fe) 14 d after the intake of labelled test meals of infant cereal (study 1) or yoghurt drink (study 2). Each test meal was fortified with 5 mg Fe as ferrous sulfate or micronised, dispersible ferric pyrophosphate. Results are presented as geometric means. There was no statistically significant difference between Fe absorption from micronised, dispersible ferric pyrophosphate- and ferrous sulfate-fortified infant cereal (3.4 and 4.1 % respectively; *P*=0.24) and yoghurt drink (3.9 and 4.2 % respectively; *P*=0.72). The results of the present studies show that micronised, dispersible ferric pyrophosphate is as well absorbed as ferrous sulfate in adults. The high relative Fe bioavailability of micronised, dispersible ferric pyrophosphate indicates the potential usefulness of this compound for food fortification.

Iron absorption: Iron fortification: Ferric pyrophosphate: Sunactive Fe™

Food fortification programmes are usually considered the most cost-effective and sustainable approach to combat Fe deficiency. However, the success of an Fe fortification programme depends largely on the careful choice of the Fe compound (Hurrell, 1997, 1998). A cheap and highly bioavailable Fe compound that causes no organoleptic changes would be the ideal fortification compound. Unfortunately, the water-soluble compounds, which are the most bioavailable, for example, ferrous sulfate, often cause unacceptable colour or flavour changes in the food vehicle (Hurrell & Cook, 1990). Ferric pyrophosphate is a water-insoluble Fe compound often used by European food companies to fortify infant cereals and chocolate-drink powders. Its main advantage is that it causes no adverse colour and flavour changes to food vehicles. However, it is only poorly soluble in dilute acid, such as the gastric juice, and is thus only of mediocre absorption in man. Human studies have reported absorption values between 15 and 75 % relative to ferrous sulfate, depending on batch and processing (Hurrell *et al.* 1989, 1991, 2000). A further disadvantage of ferric pyrophosphate

is that it cannot be used to fortify liquid products due to its water insolubility.

Recently, a micronised, dispersible ferric pyrophosphate has been developed for food fortification. This innovative compound (Sunactive Fe™; Taiyo Kagaku (Yokkaichi, Japan) is produced from ferric chloride and sodium pyrophosphate using a dispersion technique resulting in ferric pyrophosphate particles of very small average size (approximately 0.3 µm). Further, the formation of agglomerates is avoided by adding emulsifiers. This has the additional advantage that the micronised ferric pyrophosphate is dispersible in aqueous solutions and can be used to fortify liquid foods or drinks such as milk. Micronised, dispersible ferric pyrophosphate has been reported to have a similar bioavailability as ferrous sulfate in rat Hb repletion studies (Juneja *et al.* 2003).

The aim of the present study was to compare Fe absorption from micronised, dispersible ferric pyrophosphate (Sunactive Fe™) with ferrous sulfate. Fe absorption was measured in healthy women from a wheat-based infant cereal and a yoghurt drink by using a stable-isotope

Abbreviation: RBV, relative bioavailability.

* Corresponding author: Dr Lena Davidsson, fax +41 1 704 57 10, email lena.davidsson@ilw.agrl.ethz.ch

technique based on the incorporation of Fe stable isotopes into erythrocytes 14 d after administration.

Subjects and methods

Subjects

Twenty apparently healthy adult women (20–30 years; maximum body weight 60 kg) were recruited from the student and staff population at the Swiss Federal Institute of Technology Zurich and the University of Zurich. The subjects were randomly allocated into two separate studies (ten subjects/study). Exclusion criteria included pregnancy or lactation and known gastrointestinal or metabolic disorders. No medication (except oral contraceptives) or vitamin and mineral supplements were allowed during the study. Women regularly taking vitamin and mineral supplements discontinued the supplementation 2 weeks before the start of the study.

The study protocol was reviewed and approved by the ethical committee at the Swiss Federal Institute of Technology Zurich, Switzerland. Subjects were informed orally and in writing about the aims and procedures of the study. Written informed consent was obtained from all study subjects.

Study design

Fe absorption was based on erythrocyte incorporation of Fe stable-isotope labels 14 d after the intake of labelled test meals. The Fe compounds were labelled with ^{57}Fe or ^{58}Fe and added to the different test meals as described later. All test meals were fed, after an overnight fast, on two consecutive days under strictly standardised conditions and close supervision. A crossover study design was used with each woman acting as her own control. On the day before the intake of the first test meal (day 0), a venous blood sample was drawn after an overnight fast for the determination of Fe status parameters (Hb, and plasma ferritin) and body weight and height were measured. The two test meals were fed on the following days (days 1 and 2) between 07.00 and 09.00 hours. No intake of food or fluids was allowed for 3 h after the test-meal intake. A second venous blood sample was drawn 14 d after the intake of the second test meal (day 16).

Test meals

The test meals in study 1 consisted of 50 g roller-dried wheat-based infant cereal (Nestlé PTC, Orbe, Switzerland) fed with reconstituted milk (8 g Sano Lait milk powder; Coop Schweiz, Basel, Switzerland, and 75 ml deionised water). The infant cereal was made from 79.7% partially hydrolysed wheat flour, 10% sucrose, 4% honey, 3% palm oil, 0.3% calcium carbonate and 3% water. Except for Ca, no minerals or vitamins were added. The test meals in study 2 consisted of a yoghurt drink made from 170 g unskimmed yoghurt (Joghurt Nature 3.5% fat; Migros Bio, Zurich, Switzerland) and 100 g unskimmed milk (Valflora 3.8% fat; Migros, Zurich, Switzerland). Each test meal contained 5 mg added Fe, 4 mg Fe as

$^{58}\text{FeSO}_4$ plus 1 mg Fe as FeSO_4 of natural isotopic composition or 5 mg Fe as micronised, dispersible [^{57}Fe]ferric pyrophosphate. Deionised water (200 g) was served as a drink in study 1.

Stable isotope labels

[^{57}Fe]ferrous sulfate was prepared from isotopically enriched elemental Fe (Chemgas, Boulogne, France) by dissolution in sulfuric acid and dilution to the appropriate concentration. Micronised, dispersible [^{57}Fe]ferric pyrophosphate was prepared from isotopically enriched elemental Fe (Chemgas, Boulogne, France) by firstly dissolving the elemental Fe in concentrated HCl. Formed $^{57}\text{FeCl}_2$ was oxidised to $^{57}\text{FeCl}_3$ by the addition of H_2O_2 (30%, v/v). To remove impurities (iron oxides), the soluble $^{57}\text{FeCl}_3$ was extracted into diethyl ether, followed by re-extraction into deionised water. This FeCl_3 solution was evaporated under vacuum at 80°C using a rotorvap (Rotavapor; Buechi, Flawil, Switzerland). Thereafter, the resulting dark red paste was crystallised to bright yellow $^{57}\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$. From this base compound, micronised, dispersible [^{57}Fe]ferric pyrophosphate was produced by Taiyo Kagaku (Yokkaichi, Japan) by mixing $^{57}\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, emulsifiers (enzymically hydrolysed soya lecithin and polyglycerol fatty acid ester) and sodium pyrophosphate (Nambu *et al.* 1998). Particle size was measured using a sub-micron particle sizer (NiComp 370; Particle Sizing Systems, Santa Barbara, CA, USA) and the labelled compound was found to be equivalent to commercial Sunactive FeTM with respect to particle-size distribution (average particle size 0.24 µm; Fig. 1) and visual appearance. As a comparison, the particle-size distribution of a commercial food-grade ferric pyrophosphate (Dr Paul Lohmann Ltd, Emmerthal, Germany) was measured by laser light diffraction (Mastersizer X; Malvern Instruments Ltd, Malvern, UK; Fig. 1).




Fig. 1. Particle-size distribution shown as relative volume percentage frequency curve of ^{57}Fe -labelled micronised, dispersible ferric pyrophosphate (^{57}Fe Sunactive FeTM; Taiyo Kagaku, Yokkaichi, Japan) (-○-). For comparison, the particle-size distribution of commercial micronised, dispersible ferric pyrophosphate (Sunactive FeTM; Taiyo Kagaku, Yokkaichi, Japan) (-■-) and commercial ferric pyrophosphate (Dr Paul Lohmann Ltd, Emmerthal, Germany) (-▲-) are shown. Particle-size distribution was measured by laser diffraction (NiComp 370; Particle Sizing Systems, Santa Barbara, CA, USA and Mastersizer X; Malvern Instruments Ltd, Malvern, UK).

Quantification of iron isotopes in labelled iron fortificants

Isotope-dilution MS was used to determine the concentration of ^{57}Fe and ^{58}Fe stable isotopes in the micronised, dispersible ferric pyrophosphate and ferrous sulfate solutions. An accurately measured amount of Fe of natural isotopic composition was added to samples taken from the prepared solutions of labelled Fe fortificants. The Fe standard was prepared gravimetrically from an isotopic reference material (IRMM-014; EU Institute of Reference Materials, Geel, Belgium). Isotopic analysis was performed using negative thermal ionisation MS (Walczyk, 1997). Fe concentrations in each labelled Fe fortificant solution were calculated based on the shift in Fe isotopic abundances, the determined isotopic abundances of the pure isotopic labels and the natural Fe isotopic abundances (Walczyk *et al.* 1997).

Iron status measurements

Venous blood samples (7 ml) were drawn in EDTA-treated tubes at each sampling. Samples were analysed for Fe status indices (Hb, plasma ferritin) and for the incorporation of ^{57}Fe and ^{58}Fe into erythrocytes (day 16). Whole blood samples were portioned for the analysis of Hb and isotopic composition and plasma was separated, sampled and frozen for the later analysis of plasma ferritin. Hb was measured by the cyanmethaemoglobin method (Sigma kit; Sigma, St Louis, MO, USA) and plasma ferritin by ELISA (Ramco Laboratories, Houston, TX, USA). Commercial quality-control materials (DiaMed, Cressier sur Morat, Switzerland and Ramco Laboratories, Houston, TX, USA) were analysed together with the samples analysed for Hb and plasma ferritin respectively.

Quantification of iron isotope in blood

Each isotopically enriched blood sample was analysed in duplicate for its Fe isotopic composition as previously described by Walczyk *et al.* (1997). The blood samples were mineralised by microwave digestion using a mixture of HNO_3 and H_2O_2 . Fe was separated from the matrix by anion-exchange chromatography and a solvent-solvent extraction step into diethyl ether. Isotopic analyses were performed by negative thermal ionisation MS (Walczyk, 1997).

Calculation of iron absorption

The amounts of ^{57}Fe and ^{58}Fe isotopic labels in blood 14 d after the test-meal administrations were calculated based on the shift in Fe isotope ratios and on the amount of Fe circulating in the body. The calculations were based on the principles of isotope dilution and took into account that the Fe isotopic labels were not monoisotopic (Walczyk *et al.* 1997). Circulating Fe was calculated based on blood volume and Hb concentration (Kastenmayer *et al.* 1994). Blood volume calculations were based on height and weight according to Brown *et al.* (1962). For calculations of fractional Fe absorption, 80% incorporation of the absorbed Fe into erythrocytes was assumed (Hosein *et al.* 1967).

Food analysis

All test-meal components (infant cereal and milk powder, milk and yoghurt) were analysed for Fe and Ca by electrothermal-flame atomic absorption spectroscopy (SpectrAA 400; Varian, Mulgrave, Australia) after mineralisation by microwave digestion (MLS-Ethos plus; Mikrowellen-Labor-Systeme, Leutkirch, Switzerland) in a HNO_3 - H_2O_2 mixture, using a standard addition technique to minimise matrix effects. Phytic acid in the infant cereal was determined by a modification of the Makower method (Makower, 1970) in which Ce replaced Fe in the precipitation step.

Statistics

Fractional Fe absorption values are presented as geometric means and standard deviations (-1 SD, $+1$ SD). Student's paired *t* test was used to evaluate absorption data within each study. Absorption values were logarithmically transformed before statistical analysis (Excel 2002; Microsoft Corporation, Redmond, WA, USA).

Results

None of the subjects were found to be anaemic (Hb < 120 g/l). However, nine women had no Fe stores indicated by low plasma ferritin values (< 12 $\mu\text{g/l}$).

The test meals in study 1 (infant cereal) contained 0.6 mg Fe (1.1 mg Fe/100 infant cereal, 0.15 mg Fe/100 g milk powder), 167 mg Ca (148 mg Ca/100 g infant cereal, 1159 mg Ca/100 g milk powder) and 84 mg phytic acid (168 mg phytic acid/100 g infant cereal). The yoghurt drink served in study 2 contained 0.06 mg Fe (23 μg Fe/100 g unskimmed milk, 22 μg Fe/100 g unskimmed yoghurt), and 340 mg Ca (109 mg Ca/100 g unskimmed milk, 137 mg Ca/100 g unskimmed yoghurt). The ascorbic acid content was not measured as it was assumed to be negligible in both test meals.

There was no statistically significant difference between Fe absorption from the micronised, dispersible ferric pyrophosphate- and the ferrous sulfate-fortified infant cereal (geometric mean 3.4 and 4.1% respectively; $P=0.24$) (Table 1). There was also no statistically significant difference between Fe absorption from the micronised, dispersible ferric pyrophosphate- and the ferrous sulfate-fortified yoghurt drink (geometric mean 3.9 and 4.2% respectively; $P=0.72$) (Table 2).

Discussion

When measuring Fe absorption from Fe fortification compounds using stable or radioisotope techniques it is extremely important that the physical and chemical properties of the labelled compounds are comparable with those of their commercial counterpart. In the case of ferrous sulfate, it is relatively easy to prepare a labelled compound with physical and chemical properties similar to commercially available ferrous sulfate. The production of labelled micronised, dispersible ferric pyrophosphate was however more complex. This was mainly due to the necessity to synthesise

Table 1. Iron absorption by ten healthy adult women from infant cereal (study 1) fortified with ferrous sulfate or micronised, dispersible ferric pyrophosphate (Sunactive Fe^{TM*}) (5 mg iron/meal)

Subject no.	Plasma ferritin ($\mu\text{g/l}$)	Hb (g/l)	Fe absorption (%)		Relative bioavailability (%)†
			Micronised, dispersible ferric pyrophosphate	Ferrous sulfate	
1	6.9	129	4.5	8.7	52
2	11.0	136	1.8	2.4	75
3	19.8	130	3.1	4.4	70
4	56.5	140	1.5	1.9	78
5	15.8	149	3.3	5.4	61
6	9.1	127	3.4	1.8	187
7	11.8	142	4.4	8.0	55
8	11.9	138	6.4	3.1	205
9	20.8	150	2.4	3.3	72
10	7.4	132	6.9	9.3	75
Geometric mean			3.4	4.1	83
–SD			2.2	2.2	51
+SD			5.6	7.6	133

* Taiyo Kagaku, Yokkaichi, Japan.

† Fe absorption from ferrous sulfate = 100%.

Table 2. Iron absorption by ten healthy adult women from yoghurt drink (study 2) fortified with ferrous sulfate or micronised, dispersible ferric pyrophosphate (Sunactive Fe^{TM*}) (5 mg iron/meal)

Subject no.	Plasma ferritin ($\mu\text{g/l}$)	Hb (g/l)	Fe absorption (%)		Relative bioavailability (%)†
			Micronised, dispersible ferric pyrophosphate	Ferrous sulfate	
11	20.9	141	1.1	2.7	41
12	41.1	143	2.4	2.7	88
13	7.9	129	5.3	6.0	89
14	28.6	134	4.7	2.0	237
15	9.7	129	14.0	20.0	70
16	16.2	152	5.9	5.2	114
17	7.9	129	8.1	11.6	70
18	79.3	131	0.5	0.8	66
19	23.9	145	7.1	4.0	180
20	28.4	142	5.3	4.8	109
Geometric mean			3.9	4.2	94
–SD			1.5	1.7	56
+SD			10.5	10.3	157

* Taiyo Kagaku, Yokkaichi, Japan.

† Fe absorption from ferrous sulfate = 100%.

ferric trichloride in the hexahydrate form from isotopically enriched metal, free of acid residues and iron oxides. The labelled micronised, dispersible ferric pyrophosphate was made using a down-scaled manufacturing procedure similar to the commercial production procedure and the resulting compound was found to have a similar particle-size distribution as the commercial compound (Fig. 1).

The results of the present studies showed that micronised, dispersible ferric pyrophosphate is as well absorbed as ferrous sulfate from a wheat-based infant cereal as well as from a yoghurt drink. In previous studies with adult subjects, ferric pyrophosphate has been reported to have a relative bioavailability (RBV) compared with ferrous sulfate (RBV 100%) varying from 15 to 75%. In infant cereals, the values reported were between 15 and 39% (Hurrell *et al.* 1989, 1991, 2000). The high RBV of Fe

from micronised, dispersible ferric pyrophosphate, as demonstrated in the present study, is probably related to the extremely small particle size of the Fe compound which is approximately twenty times smaller than regular ferric pyrophosphate (average particle size 7.5 μm ; Fig. 1). In rat studies, decreasing the particle size of water-insoluble Fe compounds has previously been shown to have a positive influence on Fe absorption. Shah & Belonje (1973), for example, showed that the RBV of electrolytic Fe powder increased from 12 to 32% when the proportion of particles below 10 μm was increased from 62 to 99%. Further, Motzok *et al.* (1975) demonstrated that decreasing particle size of CO-reduced Fe powders from 24–40 μm to 7–10 μm increased RBV from 11 to 31%. Fe absorption from ferric orthophosphate has also been shown to be dependent on particle size as

RBV increased nearly 8-fold (from 6 to 46%) when particle size was decreased from approximately 15 µm to below 1 µm (Harrison *et al.* 1976). In human subjects, Björn-Rasmussen *et al.* (1977) reported that Fe absorption from hydrogen-reduced elemental Fe powders was dependent on their solubility in dilute acid, which in turn was partly dependent on particle size and active surface area. In the present study, it was not technically feasible to produce labelled ferric pyrophosphate with the same particle size distribution as Sunactive Fe™ without the addition of emulsifiers. Therefore, we were not able to evaluate if the high RBV of micronised, dispersible ferric pyrophosphate was only due to the small particle size or whether the emulsifiers influenced Fe absorption significantly.

Based on the results from the present studies, micronised, dispersible ferric pyrophosphate could be a very useful Fe fortificant, especially since it can be expected to cause fewer organoleptic problems than water-soluble Fe compounds. Extensive organoleptic studies, however, still remain to be carried out. Presently, Sunactive Fe™ is being used in Japan to fortify milk and milk products. Milk products have previously been shown to be difficult to fortify with readily absorbable Fe due to organoleptic problems (Demott, 1971; Edmondson *et al.* 1971; Kurtz *et al.* 1973; Wang & King, 1973). Fe fortificants that have been shown to be suitable for fluid milk fortification include ferric ammonium citrate, ferrous bisglycinate and encapsulated ferrous sulfate (Edmondson *et al.* 1971; Wang & King, 1973; Boccio *et al.* 1997; Olivares *et al.* 1997). While ferrous bisglycinate would be expected to be at least as well absorbed as ferrous sulfate (Fox *et al.* 1998), if not better (Bovell-Benjamin *et al.* 2000; Layrisse *et al.* 2000), ferric ammonium citrate has been reported to be less well absorbed than ferrous sulfate (Grebe *et al.* 1975; Layrisse *et al.* 1976; Gonzalez *et al.* 2001).

In addition to milk products, micronised, dispersible ferric pyrophosphate is potentially a suitable Fe fortificant for food vehicles that are difficult to fortify with readily available Fe such as chocolate-drink powders, cereal products, iodised salt, and bouillon cubes. Further, the overall acceptability of simulated rice grains (Kapanidis & Lee, 1996) may be improved by using micronised, dispersible ferric pyrophosphate instead of ferrous sulfate as less discoloration of fortified rice grains can be expected.

Although not statistically different, absorption from micronised, dispersible ferric pyrophosphate relative to ferrous sulfate was somewhat lower from the infant cereal than from the yoghurt drink in the present study (Tables 1 and 2). The differences in relative Fe absorption from different meals could be related to the differences in the dissolution of micronised, dispersible ferric pyrophosphate in the gastric juice as well as gastric emptying rate which both depend on meal composition (Hallberg *et al.*, 1986). Further studies are needed to evaluate the RBV of micronised, dispersible ferric pyrophosphate added to different food vehicles.

In conclusion, the results of the present studies show that Fe absorption from micronised, dispersible ferric pyrophosphate (Sunactive Fe™) is similar to that of ferrous sulfate from a fortified infant cereal as well as from a fortified yoghurt drink. The high RBV is presumably due to the

very small particle size. Micronised, dispersible ferric pyrophosphate can be expected to provoke fewer unacceptable sensory changes than water-soluble Fe compounds in different food vehicles; however, comprehensive sensory studies are now needed to fully evaluate the usefulness of this compound.

Acknowledgement

The study was supported financially by Taiyo Kagaku, Yokkaichi, Japan.

References

- Björn-Rasmussen E, Hallberg L & Rossander L (1977) Absorption of 'fortification' iron. Bioavailability in man of different samples of reduced Fe, and prediction of the effects of Fe fortification. *Br J Nutr* **37**, 375–388.
- Boccio JR, Zubillaga MB, Caro RA, Gotelli CA, Gotelli MA & Weil R (1997) A new procedure to fortify fluid milk and dairy products with high bioavailable ferrous ferrous sulphate. *Nutr Rev* **55**, 240–246.
- Bovell-Benjamin AC, Viteri FE & Allen LH (2000) Iron absorption from ferrous bisglycinate and ferric trisglycinate in whole maize is regulated by iron status. *Am J Clin Nutr* **71**, 1563–1569.
- Brown E, Hopper J Jr, Hodges J Jr, Bradley B, Wennesland R & Yamauchi H (1962) Red cell, plasma and blood volume in healthy women measured by radiochromium cell-labeling and hematocrit. *J Clin Invest* **41**, 2182–2190.
- Demott BJ (1971) Effects on flavor of fortifying milk with iron and absorption of the iron from intestinal tract of rats. *J Dairy Sci* **54**, 1609–1614.
- Edmondson LF, Douglas FW Jr & Avants JK (1971) Enrichment of pasteurized whole milk with iron. *J Dairy Sci* **54**, 1422–1426.
- Fox TE, Eagles J & Fairweather-Tait SJ (1998) Bioavailability of iron glycine as a fortificant in infant foods. *Am J Clin Nutr* **67**, 664–668.
- Gonzalez H, Mendoza C & Viteri FE (2001) Absorption of unlabeled reduced iron of small particle size from a commercial source. A method to predict absorption of unlabeled iron compounds in humans. *Arch Latinoam Nutr* **51**, 217–224.
- Grebe G, Martinez-Torres C & Layrisse M (1975) Effect of meals and ascorbic acid on the absorption of a therapeutic dose of iron as ferrous and ferric salts. *Curr Ther Res Clin Exp* **17**, 382–397.
- Hallberg L, Brune M & Rossander L (1986) Low bioavailability of carbonyl iron in man: studies on iron fortification of wheat flour. *Am J Clin Nutr* **43**, 59–67.
- Harrison BN, Pla GW, Clark GA & Fritz JC (1976) Selection of iron sources for cereal enrichment. *Cereal Chem* **53**, 78–84.
- Hosein F, Marsaglia G & Finch CA (1967) Blood ferrokinetics in normal man. *J Clin Invest* **49**, 1–9.
- Hurrell RF (1997) Preventing iron deficiency through food fortification. *Nutr Rev* **55**, 210–222.
- Hurrell RF (1998) Improvement of trace element status through food fortification: technological, biological and health aspects. *Bibl Nutr Dieta* 40–57.
- Hurrell RF & Cook JD (1990) Strategies for iron fortification of foods. *Trends Food Sci Technol* **1**, 56–61.
- Hurrell RF, Furniss DE, Burri J, Whittaker P, Lynch SR & Cook JD (1989) Iron fortification of infant cereals: a proposal

- for the use of ferrous fumarate or ferrous succinate. *Am J Clin Nutr* **49**, 1274–1282.
- Hurrell RF, Reddy MB, Burri J & Cook JD (2000) An evaluation of EDTA compounds for iron fortification of cereal-based foods. *Br J Nutr* **84**, 903–910.
- Hurrell RF, Reddy MB, Dassenko SA & Cook JD (1991) Ferrous fumarate fortification of a chocolate drink powder. *Br J Nutr* **65**, 271–283.
- Juneja LR, Nakata H, Sakaguchi N & Nanbu H (2003) A new concept of ferric pyrophosphate fortification in foods (Sunactive Fe™). 2003 INACG Symposium, 41. <http://inacg.ilsis.org/file/inacg.pdf>
- Kapanidis AN & Lee TC (1996) Novel method for the production of color-compatible ferrous sulfate-fortified simulated rice through extrusion. *J Agric Food Chem* **44**, 522–525.
- Kastenmayer P, Davidsson L, Galan P, Cherouvrier F, Hercberg S & Hurrell RF (1994) A double stable isotope technique for measuring iron absorption in infants. *Br J Nutr* **71**, 411–424.
- Kurtz FE, Tamsma A & Pallansch MJ (1973) Effect of fortification with iron on susceptibility of skim milk and nonfat dry milk to oxidation. *J Dairy Sci* **56**, 1139–1143.
- Layrisse M, Garcia-Casal MN, Solano L, *et al.* (2000) Iron bioavailability in humans from breakfasts enriched with iron bis-glycine chelate, phytates and polyphenols. *J Nutr* **130**, 2195–2199.
- Layrisse M, Martinez-Torres C & Renzi M (1976) Sugar as a vehicle for iron fortification: further studies. *Am J Clin Nutr* **29**, 274–279.
- Makower RU (1970) Extraction and determination of phytic acid in beans (*Phaseolus vulgaris*). *Cereal Chem* **47**, 288–295.
- Motzok I, Pennell MD, Davies MI & Ross HU (1975) Effect of particle size on the biological availability of reduced iron. *J Assoc Official Anal Chem* **58**, 99–103.
- Nanbu H, Nakata K, Sakaguchi N & Yamazaki Y (1998) Mineral Composition. European Patent EP 0870435A1.
- Olivares M, Pizarro F, Pineda O, Name JJ, Hertrampf E & Walter T (1997) Milk inhibits and ascorbic acid favors ferrous bis-glycine chelate bioavailability in humans. *J Nutr* **127**, 1407–1411.
- Shah BG & Belonje B (1973) Bio-availability of reduced iron. (food additives). *Nutr Rep Int* **7**, 151–156.
- Walczyk T (1997) Iron isotope ratio measurements by negative thermal ionization mass spectrometry. *Int J Mass Spectrom Ion Proc* **161**, 217–227.
- Walczyk T, Davidsson L, Zavaleta N & Hurrell RF (1997) Stable isotope labels as a tool to determine iron absorption by Peruvian school children from a breakfast meal. *Fresenius J Anal Chem* **359**, 445–449.
- Wang CF & King RL (1973) Chemical and sensory evaluation of iron-fortified milk. *J Food Sci* **38**, 938–940.