

## Use of molecular biomarkers to estimate manganese requirements for broiler chickens from 22 to 42 d of age

Lin Lu<sup>1</sup>, Bin Chang<sup>1,2</sup>, Xiudong Liao<sup>1</sup>, Runlian Wang<sup>2</sup>, Liyang Zhang<sup>1</sup> and Xugang Luo<sup>1\*</sup>

<sup>1</sup>Mineral Nutrition Research Division, Institute of Animal Science, Chinese Academy of Agricultural Sciences (CAAS), Beijing 100193, People's Republic of China

<sup>2</sup>Department of Animal Science, Guangdong Ocean University, Zhanjiang 524088, People's Republic of China

(Submitted 22 June 2016 – Final revision received 4 September 2016 – Accepted 13 September 2016)

### Abstract

The present study was carried out to evaluate dietary Mn requirements of broilers from 22 to 42 d of age using molecular biomarkers. Chickens were fed a conventional basal maize–soyabean meal diet supplemented with Mn as Mn sulphate in graded concentrations of 20 mg Mn/kg from 0 to 140 mg Mn/kg of diet for 21 d (from 22 to 42 d of age). The Mn response curves were fitted for ten parameters including heart Mn-containing superoxide dismutase (MnSOD) mRNA and its protein expression levels and the DNA-binding activities of specificity protein 1 (Sp1) and activating protein-2 (AP-2). Heart MnSOD mRNA and protein expression levels showed significant quadratic responses ( $P < 0.01$ ), and heart MnSOD activity showed a broken-line response ( $P < 0.01$ ), whereas Mn content and DNA-binding activities of Sp1 and AP-2 in the heart displayed linear responses ( $P < 0.01$ ) to dietary Mn concentrations, respectively. The estimates of dietary Mn requirements were 101, 104 and 94 mg/kg for full expressions of MnSOD mRNA level, MnSOD protein level and MnSOD activity in the heart, respectively. Our findings indicate that heart MnSOD mRNA expression level is a more reliable indicator than heart MnSOD protein expression level and its activity for the evaluation of Mn requirement of broilers, and about 100 mg Mn/kg of diet is required for the full expression of heart MnSOD in broilers fed the conventional basal maize–soyabean meal diet from 22 to 42 d of age.

**Key words:** Gene expression: Manganese: Manganese-containing superoxide dismutase: Dietary requirements: Broilers

The importance of dietary Mn for poultry has been well established<sup>(1,2)</sup>. The dietary Mn requirements are higher for broilers than for mammals, and Mn concentrations in common feeds are low<sup>(3)</sup>. Therefore, Mn additives are usually used in broiler diets for optimal growth<sup>(3)</sup>. The parameters selected to estimate nutrient requirements are important, as the maximum responses for various criteria are different<sup>(4)</sup>. The National Research Council<sup>(5)</sup> has recommended the Mn requirement for broilers from 22 to 42 d of age to be 60 mg/kg diet, which was mainly based on studies carried out more than 70 years ago. In addition, in most of these studies, semi-purified diets were used<sup>(1,6,7)</sup>, and the parameters selected were growth performance and incidence of leg disease<sup>(7,8)</sup>. Using semi-purified diets could remove confounding effects of natural and variable Mn in practical feedstuffs; however, the requirements determined under semi-purified diets might not be applicable to conventional diets because of the absence of phytate and fibre<sup>(9–12)</sup>. Phytate and fibre could inhibit Mn absorption by binding Mn in the intestinal tract, and thus decreasing Mn bioavailability. In fact, Mn is an essential component of Mn-containing superoxide dismutase (MnSOD), which plays an important role in the detoxification of superoxide

free radicals. In our laboratory, Luo *et al.*<sup>(13)</sup> reported that MnSOD activity in the heart and Mn concentrations in tissues (heart, liver, kidney and pancreas), especially the former, were sensitive criteria to estimate the Mn requirements of broilers. On the basis of these parameters, the dietary Mn requirements for broilers from 1 to 28 d of age should be 120 mg/kg<sup>(13)</sup>. Recently, in our laboratory, heart MnSOD mRNA level has been used as a quicker and more consistent biomarker than the above-mentioned parameters to estimate dietary Mn requirement in broilers from 1 to 21 d of age<sup>(14)</sup>, and 130 mg Mn/kg of diet was required for the maximal expression of this biochemical marker<sup>(14)</sup>. Later, Li *et al.*<sup>(15)</sup> reported that dietary Mn could modulate the expression of the MnSOD gene in the heart of broilers by altering DNA-binding activities of specificity protein 1 (Sp1) and activating protein-2 (AP-2) at the transcriptional level. However, dietary Mn requirements for expressions of MnSOD mRNA or protein and the DNA-binding activities of the two transcription factors (Sp1 and AP-2) in the heart of broilers from 22 to 42 d of age have not been investigated.

Therefore, the objective of this study was to estimate dietary Mn requirements of broiler chickens fed a conventional

**Abbreviations:** AP-2, activating protein-2; MnSOD, manganese-containing superoxide dismutase; Sp1, specificity protein 1.

\* **Corresponding author:** Professor X. Luo, fax +86 10 62810184, email wlysz@263.net

maize–soyabean meal diet from 22 to 42 d of age by determining MnSOD activity and its mRNA and protein expression levels, as well as the DNA-binding activities of two transcription factors (Sp1 and AP-2) in the heart of broilers at 42 d of age.

**Methods**

*Animals, diets and experimental design*

All experimental procedures were approved by the Office of the Beijing Veterinarians. The management of the Arbor Acres male broiler chickens was in agreement with the guidelines approved by the Arbor Acres Breeding Company in Beijing. The broilers were housed in electrically heated, thermostatically controlled, stainless steel cages (length 100 cm × width 50 cm × height 45 cm) equipped with fibreglass feeders and waterers. They were maintained on a 24-h constant light schedule, and allowed *ad libitum* access to experimental diets and tap water containing no detectable Mn. Broilers were vaccinated with Newcastle disease vaccine at 7 and 24 d of age, and with infectious bursal disease virus vaccine and bronchitis virus vaccine at 14 and 21 d of age, respectively.

During 1–21 d, the chickens were fed a maize–soyabean meal-based mash diet (Table 1), containing 127 mg Mn/kg of diet by analysis, which was formulated to meet or exceed the requirements of the broilers for all nutrients. About 130 mg Mn/kg of diet was decided on the basis of the result of Li *et al.*<sup>(14)</sup> in our laboratory.

**Table 1.** Ingredients and chemical composition of the basal diet for broilers, as-fed basis

Items	Days 1–21	Days 22–42
<b>Ingredients (g/kg)</b>		
Ground maize	552.4	608.2
Soyabean meal	344.0	300.4
Fishmeal	35.0	27.0
Soyabean oil	34.0	35.0
Dicalcium phosphate	12.4	11.0
Calcium carbonate*	13.5	12.3
Sodium chloride*	3.0	3.0
DL-Met	1.8	0.6
Premix†‡	3.9	2.5
<b>Chemical composition (g/kg)</b>		
Metabolisable energy (MJ/kg)	12.72	12.99
Crude protein§	215.8	203.2
Lys	12.3	10.9
Met	5.4	3.9
Met + Cys	8.9	7.3
Ca§	10.0	8.8
Non-phytate P	4.5	3.8
Mn (mg/kg)§	127	13.7

\* Feed grade before day 21 of age and reagent grade after day 21 of age.

† Provided per kilogram of diet during 1–21 d (mg/kg): retinyl acetate, 4.12; cholecalciferol, 0.11; DL- $\alpha$ -tocopheryl acetate, 16; menadione, 2; thiamin, 1; riboflavin, 8.5; pyridoxine, 2; vitamin B<sub>12</sub>, 0.01; pantothenic acid, 10; niacin, 32.5; folic acid, 1; biotin, 0.16; choline, 700; Cu, 8; Zn, 60; Fe, 80; I, 0.35; Se, 0.15.

‡ Provided per kg of diet during 22–42 d (mg/kg): retinyl acetate, 2.48; cholecalciferol, 0.06; DL- $\alpha$ -tocopheryl acetate, 9.6; menadione, 1.2; thiamin, 0.6; riboflavin, 5.1; pyridoxine, 1.2; vitamin B<sub>12</sub>, 0.006; pantothenic acid, 6; niacin, 19.5; folic acid, 0.6; biotin, 0.09; choline, 500; Cu, 8; Zn, 60; Fe, 80; I, 0.35; Se, 0.15.

§ Values determined by analysis. Each value is based on triplicate determinations.

At 22 d of age, a total of 384 chickens were randomly divided into one of eight treatment groups with eight replicate cages (6 chickens/cage) for each treatment in a completely randomised design.

Broilers were fed a maize–soyabean meal-based mash diet supplemented with 0, 20, 40, 60, 80, 100, 120 or 140 mg Mn/kg as Mn sulphate (reagent grade MnSO<sub>4</sub>.H<sub>2</sub>O) for 21 d. The basal diet met the requirements for all nutrients<sup>(5)</sup> except for Mn (Table 1). The dietary Mn contents for the eight treatments were 13.7, 32.0, 49.8, 71.0, 90.3, 111, 128 and 147 mg/kg by analysis on an as-fed basis, respectively.

At 28, 35 and 42 d of age, chicken weight and feed intake (FI) were recorded to calculate daily weight gain (G), FI and gain: feed ratio (G:F) during the 21-d study period. Incidence of leg abnormality was calculated as a percentage of chickens with visual swelling at the tibiotarsus joint within each cage weekly<sup>(14)</sup>.

*Sample collections*

At 42 d of age, sixteen chickens (two chickens from each replicate cage) from each treatment group were killed by cervical dislocation. The heart was excised immediately. A set of heart sub-samples was snap-frozen in liquid N<sub>2</sub> and then stored at –80°C for analyses of MnSOD mRNA and protein expression levels and Sp1 and AP-2 DNA-binding activities, whereas another set of sub-samples was kept on ice and stored at –20°C for subsequent determination of Mn contents and MnSOD activities. To avoid regional variation within the heart tissue, the same region of the heart was collected for the analysis of each parameter. To reduce individual biological variation, samples from two chickens in each replicate cage were pooled into one sample of equal weight before analyses, and thus there were a total of eight replicate samples for each treatment.

*Analyses of manganese contents*

Contents of Mn in tap water, diets and heart tissue were determined by inductively coupled plasma emission spectroscopy (Thermo Fisher Scientific) as described by Li *et al.*<sup>(16)</sup>. Approximately 2 ml of tap water, or 0.5 g of feed or 0.6 g of tissue was weighed in triplicate and digested with 8 ml (tap water) or 10 ml (feed and tissue) of HNO<sub>3</sub> and 0.4 ml of HClO<sub>4</sub> at 200°C in a 50-ml calibrated flask until the solution cleared and evaporated to almost dryness; the solutions were diluted 20-fold for feed samples and 25-fold for tissue samples with 2% HNO<sub>3</sub> before analysis. Validation of the mineral analysis was conducted using bovine liver powder (GBW (E) 080193; National Institute of Standards and Technology) as a standard reference material (SRM). Approximately 0.2 g of bovine liver powder was weighed and digested as described above, and was diluted 50-fold with 2% HNO<sub>3</sub> before analysis. The actual Mn recovery rate for the bovine liver SRM was determined to be about 99% in the present study.

*Heart manganese-containing superoxide dismutase activity assay*

Heart MnSOD activity was measured by the nitrite method as previously described<sup>(16)</sup>. Duplicate samples (approximately 0.6 g)

of heart tissue were homogenised in 6 ml of 0.9% cold saline, and then sonicated at 4°C for 1 min (1 s with 2-s interval; JY92-11; Ningbo Xinzhi Bio-technology Co., Ltd). The heart homogenate was centrifuged at 1500 × *g* for 15 min, and the supernatant was diluted 50-fold with deionised H<sub>2</sub>O for MnSOD activity analysis. The MnSOD activity in the heart was expressed as nitrite units (NU) per gram of fresh weight, and 1 NU was defined as the amount of enzyme needed to obtain 50% inhibition of nitrite formation.

#### RNA extraction and quantitative RT-PCR

Total RNA in heart tissue was isolated using Trizol reagent (Invitrogen Life Technologies) according to the manufacturer's protocol. Concentrations of total RNA were analysed by measuring UV light absorbance at 260 nm with a spectrophotometer (ND-100; NanoDrop Technologies). Complementary DNA was synthesised using the SuperScript<sub>III</sub> First-Strand Synthesis for RT-PCR kit (cat no. 18080-051; Invitrogen) with DNA Engine PCR instrument (Bio-RAD). Quantitative PCR was performed in triplicate on an Abi Prism 7500 apparatus (Applied Biosystems) according to optimised PCR protocols<sup>(15)</sup>. The primers for MnSOD (forward, 5'-CACTCTTCCTGACCTGCCTTAC-3'; reverse, 5'-TAGACGTCCCTGCTCCTTATTA-3') and reference gene  $\beta$ -actin (forward, 5'-CAGCTACGTTGGTGATGAAGCC-3'; reverse, 5'-CAAGAAAGATGGCTGGAAGAGG-3') were used for the amplification reactions, respectively. We used the relative standard curve method to quantify gene expression, as previously described<sup>(15)</sup>. The results are expressed as the ratio of MnSOD mRNA abundance: $\beta$ -actin mRNA abundance.

#### Measurements of specificity protein 1 and activating protein-2 DNA-binding activities

The DNA-binding activities of transcriptional factors Sp1 and AP-2 were analysed using the electrophoretic mobility shift assay (EMSA)<sup>(15)</sup>. Minced tissue samples were homogenised in HEPES buffer and incubated on ice for 30 min. After the addition of Nonidet P-40 to a final concentration of 5 g/l, the homogenate was vortexed vigorously for 15 s and centrifuged at 14 000 × *g* for 1 min at 4°C. The pellet was incubated with HEPES buffer containing 420 mmol/l NaCl for 30 min on ice. The nuclear extract supernatant was obtained by centrifugation at 12 000 × *g* for 5 min at 4°C. EMSA was performed using consensus sequences for Sp1 (59-CGAGGAGGGGCGGGCGTGTGAG-39) and AP-2 (59-GCAGTGCGGGCGCAGGCGGG-39). The complementary single-stranded oligonucleotide was end-labelled separately using biotin-11-dUTP (deoxyuridine triphosphate) and terminal deoxynucleotidyl transferase according to instructions of the biotin 3' end DNA labelling kit (Pierce) and then annealed before being used in the binding reactions. The binding specificity between Sp1 or AP-2 protein and its respective consensus sequences were verified by competition and supershift assay. Samples were then subjected to a 6% non-denaturing PAGE in Tris borate glycine buffer and transferred to a nitrocellulose membrane. The biotin-labelled DNA-protein complex was detected using streptavidin-horseradish peroxidase conjugate and chemiluminescent substrate according to instructions of the Lightshift Chemiluminescent EMSA kit (Pierce). The data are expressed as

densitometry units of the DNA-protein complex in each experimental group.

#### Western blotting assay

The heart samples were homogenised and cleaved using radio immunoprecipitation assay lysis buffer (Beyotime Institute of Biotechnology), and then sonicated at 4°C for 2 min (2 s with 10-s intervals). After centrifugation at 12 000 × *g* for 5 min at 4°C, the supernatants were collected for SDS-PAGE and Western blot analysis. Total protein contents in the supernatants were determined by Bradford Protein Assay Kit (Beyotime Institute of Biotechnology). Other procedures for Western blot analysis were performed as described by Li *et al.*<sup>(15)</sup>.

#### Statistical analyses

The effect of dietary Mn treatment was analysed by one-way ANOVA using the general liner model procedure of SAS (version 8.0; SAS Institute Inc.). Differences between means were assessed by Duncan's multiple tests. Cage was the experimental unit. Percentage data for incidence of leg abnormality were transformed to arcsine for analysis. Orthogonal comparisons were applied for linear and quadratic responses of dependent variables to independent variables. Regression analyses were used to estimate Mn requirements (the inflection point from a broken-line model, or the maximum response from a quadratic model) whenever the response was significant ( $P < 0.05$ )<sup>(14,17)</sup>.

## Results

#### Growth performance and incidence of leg abnormality

The G, FI and G:F were not affected ( $P > 0.05$ ) by dietary Mn concentration (data not shown). Incidence of leg abnormality tended to decrease with increasing dietary Mn concentrations, although treatment differences were not significant ( $P > 0.05$ ) (data not shown).

#### Heart manganese content

Dietary Mn had an effect ( $P < 0.01$ ) on Mn content in the heart (Table 2). Heart Mn content increased linearly ( $P < 0.001$ ) with increasing dietary Mn concentrations.

#### Heart manganese-containing superoxide dismutase activity

Dietary Mn concentration significantly affected ( $P < 0.0001$ ) heart MnSOD activity (Table 2). Heart MnSOD activity increased linearly ( $P < 0.0001$ ) and quadratically ( $P < 0.001$ ) as dietary Mn concentrations increased, and reached a plateau visually at about a supplemental Mn concentration of 80 mg/kg.

#### Heart manganese-containing superoxide dismutase mRNA

Heart MnSOD mRNA was significantly affected ( $P < 0.05$ ) by dietary Mn treatment, and increased linearly ( $P < 0.05$ ) and quadratically ( $P < 0.01$ ) with increasing dietary Mn concentrations (Table 2). The highest MnSOD mRNA expression level was observed visually at about a supplemental Mn concentration of 80 mg/kg.

**Table 2.** Effect of dietary manganese concentration on manganese content, manganese-containing superoxide dismutase (MnSOD) mRNA and protein expression levels and MnSOD activity in the heart of broilers at 42 d of age (Mean values with their pooled standard errors, *n* 8)

Supplemental Mn (mg/kg)	Mn content (µg/g fresh tissue)	MnSOD mRNA (arbitrary units)*	MnSOD protein (arbitrary units)*	MnSOD activity (NU/g fresh tissue)†
0	0.335 <sup>d</sup>	1.339 <sup>b</sup>	0.709 <sup>b</sup>	2598 <sup>e</sup>
20	0.371 <sup>d,c</sup>	1.405 <sup>a</sup>	0.803 <sup>a</sup>	2869 <sup>d</sup>
40	0.391 <sup>b,c</sup>	1.421 <sup>a</sup>	0.807 <sup>a</sup>	2990 <sup>d,c,b</sup>
60	0.407 <sup>a,b,c</sup>	1.423 <sup>a</sup>	0.813 <sup>a</sup>	3064 <sup>a,c</sup>
80	0.442 <sup>a</sup>	1.455 <sup>a</sup>	0.815 <sup>a</sup>	3154 <sup>a</sup>
100	0.448 <sup>a</sup>	1.439 <sup>a</sup>	0.820 <sup>a</sup>	3167 <sup>a</sup>
120	0.445 <sup>a</sup>	1.433 <sup>a</sup>	0.810 <sup>a</sup>	3196 <sup>a</sup>
140	0.446 <sup>a</sup>	1.415 <sup>a</sup>	0.810 <sup>a</sup>	3134 <sup>a,b</sup>
Pooled SEM	0.020	0.024	0.021	0.020
<i>P</i>				
Dietary Mn	0.0009	0.049	0.006	<0.0001
Mn linear	0.0007	0.017	0.004	<0.0001
Mn quadratic	0.071	0.007	0.005	0.0002

a,b,c,d,e Mean values within a column with unlike superscript letters were significantly different (*P* < 0.05).

\* Expressed as the ratio of MnSOD mRNA abundance or MnSOD protein band intensity:β-actin mRNA or β-actin protein.

† One nitrite unit (NU) was defined as the amount of enzyme needed to obtain 50% inhibition of nitrite formation.

### Heart manganese-containing superoxide dismutase protein expression level

Dietary Mn had a significant effect (*P* < 0.01) on heart MnSOD protein expression levels (Table 2). As dietary Mn concentrations increased, linear and quadratic (*P* < 0.01) responses were observed in MnSOD protein expression levels with the highest MnSOD protein expression level visually at about a supplemental Mn concentration of 100 mg/kg.

### DNA-binding activity of specificity protein 1

Dietary Mn significantly affected (*P* < 0.01) the DNA-binding activity of Sp1 in the heart (Table 3). With increasing dietary Mn concentrations, DNA-binding activities of Sp1 in the heart showed a linear response (*P* < 0.0001) to dietary Mn concentrations.

### DNA-binding activity of activating protein-2

Dietary Mn significantly affected (*P* < 0.05) the DNA-binding activity of AP-2 in the heart (Table 3). As dietary Mn concentrations increased, the DNA-binding activities of AP-2 in the heart decreased linearly (*P* < 0.001).

### Estimations of the dietary manganese requirements of broilers

Results of dietary Mn requirements of broilers as estimated by the non-linear regression analyses are shown in Table 4. On the basis of a broken-line model (*P* < 0.01) of heart MnSOD activity and quadratic models of heart MnSOD mRNA (*P* < 0.001) and protein (*P* < 0.05) expression levels with dietary Mn concentrations, dietary Mn requirements for the full expression of heart MnSOD activity and mRNA and protein expression levels were 94, 101 and 104 mg/kg, respectively for broiler chickens fed a conventional maize–soyabean meal diet from 22 to 42 d of age. These requirement values are higher than those (80, 80 and

**Table 3.** Effect of dietary manganese concentration on the DNA-binding activities of specificity protein 1 (Sp1) and activating protein-2 (AP-2) in the heart of broilers at 42 d of age (Mean values with their pooled standard errors, *n* 8)

Supplemental Mn (mg/kg)	Sp1 DNA-binding activity*	AP-2 DNA-binding activity*
0	14 041 <sup>d</sup>	19 902 <sup>a</sup>
20	18 446 <sup>c</sup>	19 366 <sup>a,b</sup>
40	18 707 <sup>b,c</sup>	19 161 <sup>a,c,d</sup>
60	18 827 <sup>b,c</sup>	17 216 <sup>b,d,e</sup>
80	21 303 <sup>b,c</sup>	16 040 <sup>e</sup>
100	20 831 <sup>b,c</sup>	16 915 <sup>d,e</sup>
120	21 694 <sup>a,b</sup>	16 610 <sup>d,e</sup>
140	21 038 <sup>b,c</sup>	16 138 <sup>e</sup>
Pooled SEM	1277	960
<i>P</i>		
Dietary Mn	0.002	0.036
Mn linear	<0.0001	0.0004
Mn quadratic	0.052	0.393

a,b,c,d,e Mean values within a column with unlike superscript letters were significantly different (*P* < 0.05).

\* Data are expressed as densitometry units of the RNA–protein complex in each experimental treatment.

100 mg/kg) derived by visual approximation from the tabulated data, but more accurate and valid than the latter as they were derived by statistical evaluation of the regression models.

### Discussion

In earlier studies, growth performance and incidence of leg abnormality were often used to assess Mn requirements in chickens fed semi-purified diets<sup>(7,8)</sup>. However, growth performance was usually not sensitive when chickens were fed a conventional maize–soyabean meal diet<sup>(3,13,15)</sup>. Incidence of leg weakness could not be completely prevented by Mn supplementation<sup>(18)</sup>, and was also not a sensitive criterion for estimation of Mn requirement<sup>(14)</sup>. The results from the current study are consistent with those from previous studies, illustrating

**Table 4.** Estimations of dietary manganese requirements based on non-linear regressions of heart manganese-containing superoxide dismutase (MnSOD) mRNA and protein expression levels and MnSOD activity on dietary supplemental manganese concentrations

Dependent variables	Regression equation*	R <sup>2</sup>	P	Dietary Mn requirement† (mg/kg)
Heart MnSOD mRNA expression level	$Y = 1.3490 + 0.00223X - 0.000012708X^2$	0.92	0.0002	101
Heart MnSOD protein expression level	$Y = 0.733 + 0.0021X - 0.00011631X^2$	0.79	0.025	104
Heart MnSOD activity	$Y = 2598 + 6.78X; Y = 2674 + 5.84X$	0.93	0.008	94

\* Y is the dependent variable and X the supplemental Mn concentration (mg/kg).

† Dietary Mn requirement = the optimal supplemental Mn concentration + the analysed Mn in the basal diet (13.7 mg/kg).

that growth performance and incidence of leg abnormality were not suitable parameters to estimate Mn requirements of broilers when a maize–soyabean meal diet is used.

Numerous studies have indicated that heart Mn concentration is a sensitive indicator of Mn deficiency in animals<sup>(13,19,20)</sup>. A study with lambs demonstrated that heart Mn could reflect the bioavailability of Mn sources because it showed a good linear response to dietary supplemental Mn (500–4000 mg/kg)<sup>(19)</sup>. Similar results in broilers were obtained by Li *et al.*<sup>(16,21)</sup> in our laboratory. The linear response of heart Mn to dietary Mn in the present study indicates that heart Mn is not a useful marker for the assessment of Mn requirement of broilers.

As a Mn-dependent enzyme, MnSOD activity is typically reduced by Mn deficiency in animals<sup>(16,20,22)</sup>. Luo *et al.*<sup>(13)</sup> found that heart MnSOD activity was a sensitive criterion for Mn requirement estimation of broilers from 1 to 28 d. Li *et al.*<sup>(14)</sup> observed a quadratic curve for heart MnSOD activity relative to dietary Mn concentration, and that the full activity expression of MnSOD in the heart of broilers from 1 to 21 d occurred at 135–156 mg Mn/kg of diet. Our results showed that dietary Mn had a broken-line effect on heart MnSOD activity, and 94 mg Mn/kg was required by broilers from 22 to 42 d of age based on this parameter. Therefore, dietary Mn needs for the full expressions of heart MnSOD activity in broilers showed a substantial decrease with increasing age of chickens.

The MnSOD gene in the heart of broilers is regulated transcriptionally in a Mn-dependent manner<sup>(15,21,23,24)</sup>. Mn deficiency in mice resulted in low Mn concentration, low MnSOD activity and low MnSOD mRNA in the liver<sup>(25)</sup>. Li *et al.*<sup>(15)</sup> also reported that dietary Mn could modulate MnSOD gene expression in heart tissue of broilers at the transcriptional level. In addition, they found that the heart MnSOD mRNA level could be used for a quicker and consistent evaluation of dietary Mn requirement of broilers from 1 to 21 d of age<sup>(14)</sup>. The results from the current study indicated that MnSOD mRNA levels in the heart increased quadratically as dietary Mn concentrations increased, and heart MnSOD mRNA level was a sensitive marker to assess Mn status and Mn requirements for broilers from 22 to 42 d of age. Therefore, our findings are in line with the observations of previous studies. On the basis of the heart MnSOD mRNA level, dietary Mn requirement for broilers from 22 to 42 d of age was 101 mg Mn/kg of diet, which was lower than the Mn requirement value (130 mg Mn/kg of diet) for broilers from 1 to 21 d of age in the study of Li *et al.*<sup>(14)</sup>.

Early researchers demonstrated that the promoter sequences of the chicken MnSOD gene contained multiple Sp1 and AP-2 binding sites, and the transcription of the MnSOD gene might

be controlled by these regulatory elements<sup>(26,27)</sup>. The results from the present study indicated that Sp1 DNA-binding activities increased and AP-2 DNA-binding activities decreased linearly as dietary Mn concentrations increased. The enhanced Sp1 DNA-binding activities and decreased AP-2 DNA-binding activities, as well as increased MnSOD mRNA levels, due to dietary Mn additions revealed that Sp1 up-regulated and AP-2 down-regulated MnSOD gene expression in broilers, respectively, as previously reported by Li *et al.*<sup>(15)</sup>. The above results also suggest that the DNA-binding activities of Sp1 and AP-2 might be sensitive indices to assess Mn status and bioavailability, but may not be suitable indices to access Mn requirements for broilers.

Clerch *et al.*<sup>(28)</sup> demonstrated that during exposure to hyperoxia (>95% O<sub>2</sub>), the lungs of rats were substantially damaged, and lung MnSOD activity dropped by approximately 50%. They also found that the fall in lung MnSOD activity was not due to a failure to increase MnSOD mRNA, but rather due to impaired translational efficiency<sup>(28,29)</sup>. These previous studies imply that the synthesis of MnSOD protein is essential for tolerance to hyperoxia in animals. The results from the present study illustrated that the addition of Mn to broiler diets increased the synthesis of MnSOD protein, and thus might enhance the ability of broilers to resist the damage caused by oxygen free radicals and other stress factors. Until now, information is still limited regarding the translational regulations of MnSOD gene expression in chickens by dietary Mn. Li *et al.*<sup>(15)</sup> found that chickens fed Mn-supplemented diets had higher MnSOD protein concentrations in heart than those fed Mn-deficient diets, suggesting that dietary Mn could modulate MnSOD gene expression in heart tissue of broilers at the translational level. Furthermore, they demonstrated that addition of Mn elevated MnSOD protein expression level and its activities in the primary broiler myocardial cells<sup>(30)</sup>. Our findings highlighted that MnSOD protein concentration was a new and valuable parameter to assess Mn requirement for broilers from 22 to 42 d of age. In the current study, the Mn requirement for the full expression of heart MnSOD protein in broilers from 22 to 42 d of age was 104 mg Mn/kg of diet, which was a little higher than the Mn requirements (101 and 94 mg/kg) for the full expression of heart MnSOD mRNA and activity.

As for the evaluations of nutrient requirements of animals, the indicators used in fitted non-linear regression models with lower P values and higher R<sup>2</sup> values should be more reliable, because lower P values could indicate the significance and validation of fitted non-linear regression models, and higher R<sup>2</sup> values could reflect better correlations between the response

criteria and dietary nutrient intakes. The above rationale and approach have been successfully used in previous studies on dietary mineral requirements of broilers<sup>(4,14,17,31,32)</sup>. In the current study, the fitted non-linear regression model for the heart MnSOD mRNA expression level had the lowest *P* value (0.0002) and higher *R*<sup>2</sup> value (0.92); therefore, the heart MnSOD mRNA expression level is a more reliable indicator for the evaluation of Mn requirements of broilers.

In conclusion, the results from the present study indicate that 94 mg Mn/kg is adequate for the full expression of MnSOD activity in the heart of broilers. Full expressions of MnSOD mRNA and protein in the heart of broilers require 101 and 104 mg Mn/kg diet, respectively. In addition, heart MnSOD mRNA expression level is a more suitable and reliable indicator than heart MnSOD activity and MnSOD protein expression level for the assessment of Mn requirement of broilers. It appears that about 100 mg Mn/kg of diet is required for broilers fed the conventional maize–soyabean meal diet from 22 to 42 d of age.

### Acknowledgements

The present study was supported by the Key International Cooperation Program of the National Natural Science Foundation of China (grant no. 31110103916; Beijing), the Agricultural Science and Technology Innovation Program (grant no. ASTIP-IAS08; Beijing) and the China Agriculture Research System (grant no. CARS-42; Beijing).

The authors' contributions are as follows: X. L. and L. L. designed the experiment; L. L. drafted the manuscript; X. L. and R. W. participated in writing and editing of the manuscript; B. C. conducted most of the experiments and analysed the data; L. Z. performed the Mn analysis; X. L. had primary responsibility for the final content. All authors read and approved the final version of the manuscript.

The authors have no personal conflicts of interest to declare.

### References

1. Wilgus HS, Norris LC & Heuser GF (1937) The role of manganese and certain other trace elements in the prevention of perosis. *J Nutr* **14**, 155–167.
2. Wang J, Wang ZY, Wang ZJ, *et al.* (2015) Effects of manganese deficiency on chondrocyte development in tibia growth plate of Arbor Acres chicks. *J Bone Miner Metab* **33**, 23–29.
3. Attia YA, Qota EM, Bovera F, *et al.* (2010) Effect of amount and source of manganese and/or phytase supplementation on productive and reproductive performance and some physiological traits of dual purpose cross-bred hens in the tropics. *Br Poult Sci* **51**, 235–245.
4. Huang YL, Lu L, Luo XG, *et al.* (2007) An optimal dietary zinc level of broiler chicks fed a corn-soybean meal diet. *Poult Sci* **86**, 2582–2589.
5. National Research Council (1994) *Nutrient Requirements of Poultry*, 9th ed. Washington, DC: National Academies Press.
6. Insko WM, Lyons M & Martin JH (1938) The quantitative requirement of the growing chick for manganese. *J Nutr* **15**, 621–627.
7. Gallup WD & Norris LC (1939) The amount of manganese required to prevent perosis in the chick. *Poult Sci* **18**, 76–82.

8. Southern LL & Baker DH (1983) *Eimeria acervulina* infection in chicks fed deficient or excess levels of manganese. *J Nutr* **113**, 172–177.
9. Halpin KM & Baker DH (1986) Long-term effects of corn, soybean meal, wheat bran, and fish meal on manganese utilization in the chick. *Poult Sci* **65**, 1371–1374.
10. Halpin KM & Baker DH (1986) Manganese utilization in the chick: effects of corn, soybean meal, fish meal, wheat bran, and rice bran on tissue uptake of manganese. *Poult Sci* **65**, 995–1003.
11. Halpin KM & Baker DH (1987) Mechanism of the tissue manganese-lowering effect of corn, soybean meal, fish meal, wheat bran, and rice bran. *Poult Sci* **66**, 332–340.
12. Baker DH & Halpin KM (1987) Efficacy of a manganese-protein chelate compared with that of manganese sulfate for chicks. *Poult Sci* **66**, 1561–1563.
13. Luo XG, Su Q, Huang JC, *et al.* (1991) A study on the optimal manganese (Mn) level in a practical diet of broiler chicks. *Chin J Anim Vet Sci* **22**, 313–317.
14. Li S, Lin Y, Lu L, *et al.* (2011) An estimation of the manganese requirement for broilers from 1 to 21 days of age. *Biol Trace Elem Res* **143**, 939–948.
15. Li S, Lu L, Hao S, *et al.* (2011) Dietary manganese modulates expression of the manganese-containing superoxide dismutase gene in chickens. *J Nutr* **141**, 189–194.
16. Li S, Luo X, Liu B, *et al.* (2004) Use of chemical characteristics to predict the relative bioavailability of supplemental organic manganese sources for broilers. *J Anim Sci* **82**, 2352–2363.
17. Jiang Y, Lu L, Li SF, *et al.* (2016) An optimal dietary non-phytate phosphorus level of broilers fed a conventional corn-soybean meal diet from 4 to 6 weeks of age. *Animal* **10**, 1626–1634.
18. Lu L, Ji C, Luo XG, *et al.* (2006) The effect of supplemental manganese in broiler diets on abdominal fat deposition and meat quality. *Anim Feed Sci Technol* **129**, 49–59.
19. Black JR, Ammerman CB & Henry PR (1985) Effects of high dietary manganese as manganese oxide or manganese carbonate in sheep. *J Anim Sci* **60**, 861–866.
20. Pallauf J, Kauer C, Most E, *et al.* (2012) Impact of dietary manganese concentration on status criteria to determine manganese requirement in piglets. *J Anim Physiol Anim Nutr* **96**, 993–1002.
21. Li SF, Luo XG, Lu L, *et al.* (2005) Bioavailability of organic manganese sources in broilers fed high dietary calcium. *Anim Feed Sci Technol* **124**, 703–715.
22. Zhu YW, Lu L, Li WX, *et al.* (2015) Effect of dietary manganese on antioxidant status and expression levels of heat-shock proteins and factors in tissues of laying broiler breeders under normal and high environmental temperatures. *Br J Nutr* **114**, 1965–1974.
23. Li SF, Luo XG, Lu L, *et al.* (2008) Effect of intravenously injected manganese on the gene expression of manganese-containing superoxide dismutase in broilers. *Poult Sci* **87**, 2259–2265.
24. Luo XG, Li SF, Lu L, *et al.* (2007) Gene expression of manganese-containing superoxide dismutase as a biomarker of manganese bioavailability for manganese sources in broilers. *Poult Sci* **86**, 888–894.
25. Borrello S, De Leo ME & Galeotti T (1992) Transcriptional regulation of MnSOD by manganese in the liver of manganese-deficient mice and during rat development. *Biochem Int* **28**, 595–601.
26. DiSilvestre D, Kleeberger SR, Johns J, *et al.* (1995) Structure and DNA sequence of the mouse MnSOD gene. *Mamm Genome* **6**, 281–284.

27. Xu Y, Porntadavity S & St Clair DK (2002) Transcriptional regulation of the human manganese superoxide dismutase gene: the role of specificity protein 1 (Sp1) and activating protein-2 (AP-2). *Biochem J* **362**, 401–412.
28. Clerch LB & Massaro D (1993) Tolerance of rats to hyperoxia. Lung antioxidant enzyme gene expression. *J Clin Invest* **91**, 499–508.
29. Clerch LB, Massaro D & Berkovich A (1998) Molecular mechanisms of antioxidant enzyme expression in lung during exposure to and recovery from hyperoxia. *Am J Physiol* **274**, L313–L319.
30. Li SF, Lu L, Liao XD, *et al.* (2016) Manganese elevates manganese superoxide dismutase protein level through protein kinase C and protein tyrosine kinase. *Biometals* **29**, 265–274.
31. Liao XD, Li A, Lu L, *et al.* (2013) Optimal dietary zinc levels of broiler chicks fed a corn-soybean meal diet from 22 to 42 days of age. *Anim Prod Sci* **53**, 388–394.
32. Liu SB, Liao XD, Lu L, *et al.* (2016) Dietary non-phytate phosphorus requirement of broilers fed a conventional corn-soybean meal diet from 1 to 21 d of age. *Poult Sci* (publication ahead of print version 2 August 2016).