

NOTES FOR AUTHORS

Proceedings of the Royal Society of Edinburgh: Section A is a general journal, and papers in all areas of mathematics will be considered. Papers to be considered for publication should be sent to the Publications Manager, The Royal Society of Edinburgh, 22 George Street, Edinburgh EH2 2PQ, Scotland.

A paper by more than one author must be submitted with a statement, signed by each author, to the effect that the paper in its entirety is approved by the joint authors and naming the author who will be responsible for correspondence with the Society.

Authors will receive fifty (50) offprints free of charge, this number to be shared between joint authors. Additional offprints may be obtained, in units of fifty, at a fixed scale of prices given on a form which will be attached to the proof.

Authors must prepare their papers as concisely as possible. Manuscripts should be submitted in triplicate, on single-sided A4 paper, double spaced with adequate margins. Authors are advised to retain a copy of their papers as the Society cannot accept responsibility for any loss.

Every paper must be accompanied by a Synopsis, in general not exceeding two hundred words, which will be printed in small type at the beginning of the paper.

References within the text should be indicated by bold numbers in square brackets, e.g. [2] or [3, p. 167]. References at end of text should be in alphabetical order, numbered sequentially.

Authors should ensure that punctuation carries through the mathematics in the proper manner. The use of hyphens should be consistent. In the text avoid such abbreviations as: iff, w.r.t. and thm.

Footnotes should be avoided. Headings should not be underlined. Every effort should be made to avoid complicated subscripts, superscripts, ranges of summation and integration. Replace $e^{(\dots)}$ by $\exp[\dots]$ if the expression in parenthesis is complicated. Use the prime ' or d/dx , but preferably not a dot, to denote ordinary differentiation. If possible use subscripts to denote partial differentiation of $\partial/\partial x$ etc. Bars reaching over several letters should be avoided: use $\sqrt{()}$ or the exponent $1/2$ for the square root.

Note that confusion very often arises between 1 (one) and l (ell); 0 (zero) and O (Capital oh); \circ (composition) and o (lower case oh); x and \times ; U and \cup ; c and \subset ; \in (belongs to) and ε (epsilon); \emptyset (empty set) and ϕ (phi); $_1$ and comma ,; prime ' and 1 ; K and κ ; p and ρ ; w and ω ; \sum (summation) and Σ (capital sigma); \prod (product) and Π (capital pi); v (lower case vee) and ν (Greek nu); a (lower case a) and α (Greek alpha); y (lower case y) and γ (Greek gamma). Please provide pencilled indicators in the margin where necessary. Where capitals and lower case of the same shape have to be printed, please indicate accordingly. Show italics by single underlining (except in the formulae which are set up normally in italics), bold face/Clarendon by wavy underlining and Greek by red underlining.

The statement of theorems, lemmas, et cetera, will be printed in italics and should be underlined. In definitions key words only should be in italics.

Equations should be indicated by numbers in parentheses in the right-hand margin.

Proofs of papers will be sent to the author. The cost of *authors' corrections in excess of five per cent* of the printers' charge for the setting of a particular paper *will be charged to the author*.

Copyright

© 1996 The Royal Society of Edinburgh

Except as otherwise permitted under the Copyright, Designs and Patents Act, 1988, this publication may only be reproduced, stored or transmitted, in any form or by any means, with the prior permission in writing of the publisher, or, in the case of reprographic reproduction, in accordance with the terms of a licence issued by the Copyright Licensing Agency. In particular, the Society permits the making of a single photocopy of an article from this issue (under Sections 29 and 38 of the Act) for an individual for the purposes of research or private study.

CONTENTS

TASSILO KÜPPER and ACHILLES TERTIKAS	
A global branch of positive solutions above the continuous spectrum for problems with indefinite nonlinearities	465
A. NOVICK-COHEN and SONGMU-ZHENG	
The Penrose–Fife-type equations: counting the one-dimensional stationary solutions	483
P. B. BAILEY, W. N. EVERITT and A. ZETTL	
Regular and singular Sturm–Liouville problems with coupled boundary conditions	505
I. ALONSO-MALLO and C. PALENCIA	
On the convolution operators arising in the study of abstract initial boundary value problems	515
ZVI ARTSTEIN and ALEXANDER VIGODNER	
Singularly perturbed ordinary differential equations with dynamic limits	541
JERRY L. BONA and BING-YU ZHANG	
The initial-value problem for the forced Korteweg–de Vries equation	571
P. KORMAN, YI LI and TIANCHENG OUYANG	
Exact multiplicity results for boundary value problems with nonlinearities generalising cubic	599
KENNETH F. ANDERSEN	
Cesàro averaging operators on Hardy spaces	617
CARME CASCANTE and JOAQUÍN ORTEGA	
A characterisation of tangential exceptional sets for H_α^p , $\alpha p = n$	625
W. ALLEGRETTO and A. BARABANOVA	
Positivity of solutions of elliptic equations with nonlocal terms	643
FABIÁN FLORES-BAZÁN	
Some remarks about relaxation problems in the Calculus of Variations	665
MARIA E. SCHONBEK and MICHAEL WIEGNER	
On the decay of higher-order norms of the solutions of Navier–Stokes equations	677