

Galaxy And Mass Assembly (GAMA)

Simon P. Driver¹ and the GAMA team[†]

¹(SUPA) School of Physics and Astronomy, University of St Andrews
St Andrews, Fife, KY16 8RS, Scotland
email: spd3@st-and.ac.uk

Abstract. The GAMA survey aims to deliver 250,000 optical spectra (3–7Å resolution) over 250 sq. degrees to spectroscopic limits of $r_{AB} < 19.8$ and $K_{AB} < 17.0$ mag. Complementary imaging will be provided by GALEX, VST, UKIRT, VISTA, HERSCHEL and ASKAP to comparable flux levels leading to a definitive multi-wavelength galaxy database. The data will be used to study all aspects of cosmic structures on 1kpc to 1Mpc scales spanning all environments and out to a redshift limit of $z \approx 0.4$. Key science drivers include the measurement of: the halo mass function via group velocity dispersions; the stellar, HI, and baryonic mass functions; galaxy component mass-size relations; the recent merger and star-formation rates by mass, types and environment. Detailed modeling of the spectra, broad SEDs, and spatial distributions should provide individual star formation histories, ages, bulge-disc decompositions and stellar bulge, stellar disc, dust disc, neutral HI gas and total dynamical masses for a significant subset of the sample ($\sim 100k$) spanning both the giant and dwarf galaxy populations. The survey commenced March 2008 with 50k spectra obtained in 21 clear nights using the Anglo Australian Observatory's new multi-fibre-fed bench-mounted dual-beam spectroscopic system (AAΩ).

Keywords. galaxies: general, galaxies: structure, galaxies: formation, galaxies: evolution

1. GAMA Motivation

Galaxy And Mass Assembly (GAMA) is a major expansion of the Millennium Galaxy Catalogue (MGC) survey (Liske *et al.* 2003; Allen *et al.* 2006, Driver *et al.* (2005) and a natural extension of the extremely productive nearby “Legacy” surveys (e.g., SDSS, 2MASS, HIPASS etc). In comparison to the superb SDSS survey GAMA will only sample 250 sq degrees of sky but will extend to significantly fainter spectroscopic limits ($12\times$ the redshift density of SDSS main, $5\times$ stripe 82), to higher spatial (0.6" FWHM) and spectral (3–7Å) resolutions, as well as moving to a far broader wavelength coverage (UV to Radio). GAMA has come about by parallel technological developments leading to a suite of new facilities whose survey sensitivities, resolutions, and capabilities are reasonably well matched. Until now the study of galaxies has generally been restricted to either large samples of limited wavelength data or multi-wavelength studies of small (and often biased) samples. However galaxy systems are extremely complex and diverse, exhibiting strong environmental and mass dependencies and containing distinct but interlinked

[†] I.K. Baldry (LJMU), S. Bamford (Nott), J. Bland-Hawthorn (USyd), T. Bridges (AAO), E. Cameron (StA), C. Conselice (Nott), W.J. Couch (Swinburne), S. Croom (USyd), N.J.G. Cross (Edin), S.P. Driver (StA), L. Dunne (Nott.), S. Eales (Cardiff), E. Edmondson (Ports), S.C. Ellis (USyd), C.S. Frenk (Durham), A.W. Graham (Swinburne), H. Jones (AAO), D. Hill (StA), A. Hopkins (USyd), E. van Kampen (Inns), K. Kuijken (Leiden), O. Lahav (UCL), J. Liske (ESO), J. Loveday (Sussex), B. Nichol (Ports.), P. Norberg (Edin), S. Oliver (Sussex), H. Parkinson (Edin), J.A. Peacock (Edin), S. Phillipps (Bristol), C.C. Popescu (UCLan), M. Prescott (LJMU), R. Proctor (Swinburne), R. Sharp (AAO), L. Staveley-Smith (UWA), W. Sutherland (QMW), R.J. Tuffs (MPIK), S. Warren (Imperial).




Figure 1. Facilities contributing to the final GAMA database.

components (AGN, nucleus, bulge, pseudo-bulge, bar, disc etc) and constituents (SMBH, plasma, stars, gas, dust etc). It then follows that a clear understanding of galaxy formation and evolution may only come about via the construction of a comprehensive survey which simultaneously samples all of these facets. The GAMA team aims to provide this data. In addition to the provision of a generic galaxy database, the GAMA project also includes a number of more focussed science goals, in particular:

1. Measurement of the Halo Mass Function via virialised group velocity dispersions to directly test the *numerical* prediction from CDM (and WDM) simulations.
2. Measurement of the dynamic, baryonic, HI and stellar mass functions to LMC masses versus redshift, environment, type, and component (as well as higher order relations, e.g., mass-spin [$M - \lambda$]).
3. Measurement of the recent merger rates and star formation rates versus type, mass and environment over a 3–4 Gyr baseline.

2. Facilities Contributing to GAMA

Fig. 1 shows the facilities currently contributing to the GAMA project along with the approximate time allocations within the GAMA sky regions (see Fig. 2 and Tables 1 & 2). The expected source resolution and detection sensitivities (5σ point source) are shown in Fig. 3 (upper and lower) in arcseconds and milliJanskys. Overlaid on the lower panel is the modelled NGC891 spectra (Popescu *et al.* 2000) with a weak AGN added and transposed to $z \approx 0.1$. The UKIRT data is provided courtesy of the UKIDSS LAS Public




Figure 2. (black rectangles) Survey regions of first year data obtained in 21 clear nights as compared to the 2dFGRS (green outline) and MGC (blue band) surveys. Also shown are galaxies within 10Mpc (red dots), abell clusters (yellow), bright stars (cyan) and the NGC catalogue (magenta). The dashed regions show the final GAMA survey extent.

Survey while the VST and VISTA data are provided via the KIDS and VIKING ESO Public Surveys (whose teams include GAMA members). The Herschel data is provided as part of the broader Herschel-ATLAS survey and a proposal is currently pending to complete GALEX medium depth observations of the GAMA regions (~50% already covered with MIS). A major advancement over previous surveys will be the inclusion of radio data via ASKAP (Australian Square Kilometer Array Pathfinder, see Johnston *et al.* 2007) which should allow HI mass, dynamical mass and continuum measurements for all GAMA galaxies with high or normal neutral gas content. The deep ASKAP pointing (a single $6^\circ \times 6^\circ$ field of 1 year integration) is predicted to have an $n(z)$ distribution comparable to that derived for GAMA (see Fig. 4). While the exact location of the ASKAP deep pointing has not been finalised it is highly likely, given the similarity in the $n(z)$ distributions that one of the GAMA fields will be adopted (nominally the 12hr field).

Table 1. The current extent of the GAMA survey showing the year 1 regions and two possible expansion options currently under consideration.

GAMA Field ID	Year 1 Regions		Extension 1		Extension 2	
	RA(deg)	δ (deg)	RA(deg)	δ (deg)	RA(deg)	δ (deg)
G09	129.0 – 141.0	-1 – +2	129.0 – 141.0	-1 – +3	129.0 – 141.0	-3 – +3
G12	174.0 – 186.0	-2 – +1	174.0 – 186.0	-2 – +2	129.0 – 186.0	-3 – +3
G14	211.5 – 223.5	-1 – +2	211.5 – 223.5	-2 – +2	211.5 – 223.5	-3 – +3
G03	–	–	45.0 – 57.0	-28 – -31	–	–
G22	–	–	348.0 – 360.0	-28 – -31	–	–




Figure 3. (lower) Spectroscopic, photometric and (upper) resolution sensitivities in mJy and arcseconds respectively. Overlaid are the typical AGN (green), stellar (blue) and dust emissions (red; based on NGC891 at $z \approx 0.1$ with weak AGN added).

3. First Light

The survey commenced March 2008 with 50k spectra obtained in 21 clear nights using the Anglo Australian Observatory's new multi-fibre-fed bench-mounted dual-beam spectroscopic system (AA Ω). Fig. 5 shows the areas of sky surveyed (upper) and the resulting cone plot (lower). This includes the existing 25k redshifts within these regions from the MGC, SDSS and 2dFGRS. AA Ω represents an upgrade of the pre-existing 2dF system using the same fibre positioner/tumbler but replacing the two telescope mounted spectrographs with a single bench-mounted, double-beam spectrograph (see Sharp *et al.* 2006 or the AAO website). The facility can be used for both multi-fibre and integral field spectroscopy and in multi-fibre mode is capable of obtaining 350–400 spectra in a single 2° diameter field. During an 8hr observation period the system is capable of obtaining ~ 3000 spectra. Data are reduced in real-time and redshifts also obtained in real-time via cross-correlation with a template library. All data are later re-reduced and processed with GANDALF (see Schawinski *et al.* 2007) to obtain line indices and velocity dispersion measurements. The GAMA survey at the AAT uses the 580V and 385R gratings yielding a resolution of 1300 or $3\text{--}7\Delta\text{FWHM}$.




Figure 4. The GAMA $n(z)$ distribution compared to that expected by the ASKAP 1 year deep stare, arbitrarily normalised to give a comparable height peak.

Table 2. Time allocations, resolutions and sensitivities of the facilities contributing to the GAMA survey. Limits are for 5σ detections in AB mag or mJsky ($1Jsky = 3631 \times 10^{-0.4m_{AB}}$).

Window	Facility	Collab.(Time)	Detection limits					Resol.	GAMA Fields
UV	GALEX	MIS + 25hrs pend.	FUV		NUV		FUV/NUV	G09,G12,G15	
			23.0		23.0		4-5''		
Opt	SDSS	DR6	u	g	r	i	z	u-z	G09,G12,G15
			22.0	22.0	22.2	21.3	20.5	1.0''-2.5''	
	VST	KIDS (70n)	24.8	25.4	25.2	24.2	-	0.6''-1.0''	All
Near-IR	UKIRT	LAS (10n)	Z	Y	J	H	K	Z-K	G09,G12,G15
			-	20.9	20.6	20.3	20.1	0.6''-1.0''	
	VISTA	VIKING (40n)	23.1	22.3	22.1	21.5	21.1	0.6''-0.8''	All
Far-IR	Herschel	ATLAS (200hrs)	110 μ m	170 μ m	250 μ m	350 μ m	500 μ m	110-500 μ m	All
			67mJ	94mJ	45mJ	62mJ	53mJ	8-35''	
Radio	ASKAP	DEEP (1 year)	0.7-1.8 GHz				0.7-1.8GHz	G12?	
			10 μ J				\sim 10''		

4. Summary

The GAMA project has commenced with data flows imminent from a number of international facilities. The survey will allow for a comprehensive study of structure on 1kpc to 1Mpc scales as well as the subdivision of the galaxy population into its distinct




Figure 5. (upper panel) The GAMA sky coverage and (lower panel) cone plot from the first year of GAMA observations at the AAT.

components and constituents. Progress and data releases (1st data release forecast for Dec 2009) can be monitored via the GAMA website: <http://www.eso.org/~jliske/gama/> and anyone interested in further details should contact Simon Driver at spd3@st-and.ac.uk.

The GAMA team acknowledges all staff of the Anglo Australian Observatory for provision of the superb AAΩ facility and the continued smooth running of the Anglo Australian Telescope. SPD thanks the IAU254 organisers for a most enjoyable meeting.

References

- Allen, P. D., Driver, S. P., Graham, A. W., *et al.*, 2006, MNRAS, 371, 2
 Driver, S. P., Liske, J., Cross, N. J. G., *et al.*, 2005, MNRAS, 360, 81
 Driver, S. P., Popescu, C. C., Tuffs, R. J., *et al.*, 2008, ApJL, 678, 101
 Johnston, S., *et al.*, 2007, PASA, 24, 174
 Liske, J., Lemon, D. J., Driver, S. P., *et al.*, 2003, MNRAS, 344, 307
 Popescu, C. C., Misiriotis, A., Kylafis, N. D., *et al.*, 2000, A&A, 362, 138
 Sharp, R., *et al.*, 2006, SPIE, 6269, 14
 Schawinski, K., Thomas, D., Sarzi, M., *et al.*, 2007, MNRAS, 382, 1415