

## Effect of *Bifidobacterium lactis* HN019 on inflammatory markers and oxidative stress in subjects with and without the metabolic syndrome

Luciana J. Bernini<sup>1</sup>, Andréa N. Colado Simão<sup>2,3</sup>, Cíntia H. B. de Souza<sup>1</sup>, Daniela F. Alfieri<sup>3</sup>, Liliane G. Segura<sup>1</sup>, Giselle N. Costa<sup>1\*</sup> and Isaias Dichi<sup>4</sup>

<sup>1</sup>Programa de Mestrado em Ciência e Tecnologia de Leite e Derivados, Pitágoras-UNOPAR Londrina, Parana, 86041-140, Brazil

<sup>2</sup>Departamento de Patologia, Análises Clínicas e Toxicologia, Universidade Estadual de Londrina, Londrina, Parana, 86038-440, Brazil

<sup>3</sup>Pesquisador do Laboratório de Imunologia Aplicada, Universidade Estadual de Londrina, Londrina, Parana, 86038-440, Brazil

<sup>4</sup>Departamento de Medicina Interna, Universidade Estadual de Londrina, Londrina, Parana, 86038-440, Brazil

(Submitted 8 February 2018 – Final revision received 7 June 2018 – Accepted 9 June 2018 – First published online 30 July 2018)

### Abstract

Beneficial effects of probiotics have been reported on body weight, lipid and carbohydrate metabolism, inflammatory state and oxidative stress in healthy subjects and in many metabolic and inflammatory diseases. The aim of this study was to evaluate the effects of *Bifidobacterium lactis* HN019 on inflammatory state and nitro-oxidative stress in patients with and without the metabolic syndrome (MetS). The usual diets of the thirty-three subjects were supplemented with probiotic milk for 90 d. Inflammatory markers and oxidative measurements were performed. In relation to the baseline values, subjects in both groups showed a decrease in homocysteine ( $P=0.02$  and  $P=0.03$ , respectively), hydroperoxides ( $P=0.02$  and  $P=0.01$ , respectively) and IL-6 levels ( $P=0.02$ ). Increases in adiponectin ( $P=0.04$ ) and nitric oxide metabolites (NOx,  $P=0.001$ ) levels were only seen in the group with the MetS in relation to the baseline values, whereas only the individuals without the MetS had increases in total radical-trapping antioxidant parameter levels ( $P=0.002$ ). In conclusion, *B. lactis* HN019 have several beneficial effects on inflammatory and oxidative biomarkers in healthy subjects and the MetS patients. Patients with the MetS showed a specific improvement in adiponectin and NOx levels, whereas a specific favourable effect was shown in the antioxidant defenses in healthy subjects. If the results obtained in the present study are confirmed, supplementation of fermented milk with probiotics in healthy subjects and patients with the MetS must be further discussed.

**Key words:** Probiotics: Metabolic syndrome: Cytokines: Oxidative stress: Nitrosative stress

The metabolic syndrome (MetS) can be defined as a set of metabolic risk factors involved in the development of CVD and/or type 2 diabetes<sup>(1)</sup>. Abdominal obesity and insulin resistance play a key role in the genesis of this syndrome.

The MetS has increased in epidemic proportions worldwide<sup>(2–5)</sup>. Currently, the search for other factors that cause the MetS, or alternatives that may prevent its occurrence, is often associated with the intestinal microbiota. Indeed, recent findings indicate that altered gut microbial composition and diversity have a vital role in the pathophysiology of metabolic disorders, as the commensal gut microbes modulate host metabolism by affecting energy extraction from food and by biochemically converting molecules derived from the host or from the gut microbes themselves<sup>(5–7)</sup>.

The imbalance of the intestinal microbiota has been strongly associated with the occurrence of cardiovascular and neurological

diseases, decrease of immune function, increase in oxidative stress and many other dysbiosis<sup>(8–11)</sup>.

In contrast, consumption of probiotics has been associated with the gut microbial balance and, consequently, various positive actions on host health are reported, including beneficial effects on lipid and glucose metabolism, inflammatory state and oxidative stress in healthy subjects and in those with many metabolic and inflammatory diseases<sup>(12,13)</sup>. Although the effects of probiotics on oxidative stress are still conflicting<sup>(14)</sup>, several reports showed that probiotics in patients with metabolic diseases resulted in favourable metabolic consequences especially reducing oxidative stress in diabetic patients<sup>(14,15)</sup>. However, studies of probiotics in relation to oxidative stress in patients with the MetS have not been performed yet.

In a previous study, we reported the effects of *Bifidobacterium lactis* HN019 in patients with the MetS<sup>(16)</sup>. The group that

**Abbreviations:** hsCRP, high-sensitivity C-reactive protein; MetS, metabolic syndrome; NOx, nitric oxide metabolites; TRAP, total radical-trapping antioxidant parameter.

\* **Corresponding author:** G. N. Costa, fax +55 43 3371 7834, email gcnobre@gmail.com

ingested probiotic milk for 45 d presented favourable effects on lipid profile and inflammatory biomarkers (TNF- $\alpha$  and IL-6) when compared with the control group, which did not receive the probiotic milk.

However, some questions have emerged after the aforementioned study: which differences would be shown in a study comparing healthy subjects with the MetS patients? How would oxidative stress biomarkers behave in such a comparison? Would the beneficial effects of the same probiotic last for even longer periods of time? Thus, the aim of this study was to evaluate the effects on inflammatory state and oxidative stress of *B. lactis* HN019 in patients with and without the MetS, whose diet was supplemented with the probiotic fermented milk for 90 d.

## Methods

### Study design

The study was conducted according to the guidelines in the Declaration of Helsinki and approved by the Ethical Committee involving humans of the University of Londrina, Parana, Brazil, protocol number 469.555/201. This was a randomised and double-blind study. A total of thirty-three patients, men and women, aged between 18 and 60 years were recruited from a health facility in the city of Jataizinho, Parana State, Brazil, and received detailed information about the purpose and procedures of the study and agreed to participate and signed a consent form. Thus, the volunteers were divided in two groups: patients without the MetS (control group, *n* 14) and patients with the MetS (*n* 19). Patients with the MetS were selected among volunteers who presented three or more of the following five criteria: (1) waist circumference (WC)  $\geq 102$  cm in men and 88 cm in women, (2) fasting TAG levels  $\geq 150$  mg/dl ( $\geq 1.7$  mmol/l), (3) HDL  $\leq 40$  mg/dl ( $\leq 1.0$  mmol/l) in men or  $\leq 50$  mg/dl ( $\leq 1.3$  mmol/l) in women, (4) blood pressure  $\geq 130/85$  mmHg (or anti-hypertensive medication use) and (5) fasting glucose levels  $\geq 100$  mg/dl ( $\geq 5.6$  mmol/l) or the use of antidiabetic medication<sup>(1)</sup>.

All individuals daily ingested 80 ml of the probiotic milk containing an average of  $3.4 \times 10^8$  colony-forming units (CFU)/ml of *Bifidobacterium animalis* subsp. *lactis* subsp. nov. HN019 for 90 d. Anthropometric measurements and biochemical parameters were assessed in blood or plasma samples before (T0) and after 90 d (T90) of probiotic milk ingestion.

### Inclusion and exclusion criteria

Inclusion criteria were the ages between 18 and 60 years, and the MetS diagnosis criteria according to the National Cholesterol Education Program, Adult Treatment Panel III<sup>(1)</sup>. Exclusion criteria were thyroid, adrenal, renal, hepatic, gastrointestinal, infectious or oncological diseases; the utilisation of hormone replacement therapy; vitamins; antioxidant supplements; antibiotics and anti-inflammatories; and ingestion of foods or supplements containing probiotic micro-organisms, besides lactose-intolerant individuals. The individuals of both groups did not drink alcohol regularly. In addition, the volunteers were instructed not to modify their eating habits and/or physical

activities and other lifestyle factors, as well as to not use another probiotic or antibiotic throughout the intervention period.

### Milk fermented with probiotics

The commercial strain Bifido HOWARU<sup>®</sup> *Bifidobacterium lactis* HN019 (Danisco) was used to produce the probiotic milk with  $3.4 \times 10^8$  CFU/ml. The probiotic milk was produced and packaged aseptically at the Laboratory of Microbiology of the University of North Parana, according to the current Brazilian legislation<sup>(17)</sup>. The product ready for consumption presented 0.38% acidity (lactic acid), a pH of 5.15, 2.80% protein, 7.54% carbohydrates, 0.0% fat, 12.08% total solids and 0.48% ash, which resulted in 33.09 energy content/bottle. In addition, the final product contained 0.01% (v/v) orange and/or vanilla flavour<sup>(16)</sup>. Pathogens were not detected, as required by legislation<sup>(18)</sup>; thus, the product was characterised as suitable for consumption. Weekly, seven bottles of 80 ml were distributed to each volunteer during 13 weeks (90 d), and the patients were instructed to drink one bottle daily for breakfast.

### Waist circumference and blood pressure measurements

WC was measured with a soft tape on subjects in a standing position midway between the lowest rib and the iliac crest. Three blood pressure measurements, systolic (SBP) and diastolic blood pressures (DBP), taken with a 1-min interval between them after the subject had been seated were recorded. The mean of these measurements was used in the analysis<sup>(19)</sup>.

### Measurements of serum high-sensitivity C-reactive protein, cytokines and immune mediators

Blood samples were obtained after 12 h of fasting at the Central Health facility in the city of Jataizinho, Parana State, Brazil. The samples were collected and taken to the Clinical Immunology Laboratory at the Londrina University Hospital for analysis. The plasma and serum were divided into aliquots and stored in a freezer at  $-70^\circ\text{C}$  (Indrel) until analysis. For further analysis, the samples followed routine examinations in that laboratory. Serum high-sensitivity C-reactive protein (hsCRP) was measured using a nephelometric assay (DadeBehring). Serum homocysteine level was determined by chemiluminescence microparticle immunoassay (Architect; Abbott Laboratory). In addition, IL-6, leptin and adiponectin were measured by a sandwich ELISA using a commercial immunoassay (R&D System).

### Oxidative and nitrosative stress markers

**Tert-butyl hydroperoxide-initiated chemiluminescence.** Plasma levels of lipid hydroperoxide were evaluated by tert-butyl hydroperoxide-initiated chemiluminescence (CL-LOOH) as described previously<sup>(20)</sup>, and the results were expressed in relative light units.

**Determination of nitric oxide metabolites.** Serum levels of nitric oxide metabolites (NOx) were assessed by nitrite (NO<sub>2</sub><sup>-</sup>)



concentration according to the Griess reaction, supplemented by the reduction of nitrate to nitrite with cadmium<sup>(21,22)</sup>.

**Total radical-trapping antioxidant parameter.** Total radical-trapping antioxidant parameter (TRAP) was determined as reported by Repetto *et al.*<sup>(23)</sup>. This method detects hydrosoluble and/or liposoluble plasma antioxidants by measuring the chemiluminescence inhibition time induced by 2,2-azobis (2-amidinopropane). The system was calibrated with the vitamin E analogue Trolox. Serum uric acid levels were determined using a biochemical auto-analyser (Dimension Dade AR; Dade Behring) and were used to correct the TRAP values. TRAP measurements in conditions associated with hyperuricaemia, such as the MetS, may be inaccurate because uric acid concentration accounts for 60% of total plasma antioxidant capacity. Thus, a correction of TRAP based on uric acid concentration was performed<sup>(24)</sup>.

**Sulphydryl groups.** Sulphydryl (SH) groups of proteins were evaluated in plasma samples by a spectrophotometric assay based on 2,2-dithiobisnitrobenzoic acid, as reported previously, and the results are expressed in  $\mu\text{M}$ <sup>(25)</sup>.

**Statistical analysis**

The sample size was calculated to obtain a statistically significant result when changes of the parameters evaluated were at >10% level. This study was performed to obtain 80% statistic power. The Shapiro–Wilk test was used to assess normality of distribution. Logarithmic transformation of continuous data was used in the analyses (SBP, DBP, hsCRP and leptin) when the variables were not normally distributed or when there was heterogeneity of variance (as assessed with the Levene test). Analyses of contingency tables ( $\chi^2$  or Fisher’s test) were used to check the associations between categorical variables (sex, ethnicity, smoking, physical exercise, hypertension, diabetes mellitus, the MetS and insulin resistance), and data were

expressed as absolute number (*n*) and percentage (%). Student’s *t* test (WC, blood pressure measurements, homocysteine, hsCRP and oxidative and nitrosative stress (O&NS) markers) or Mann–Whitney (IL-6) test was performed to compare differences between parameters of groups at baseline and differences across treatment groups (intergroup changes). Paired Student’s *t* test (WC, blood pressure measurements, homocysteine, hsCRP and O&NS markers) or Wilcoxon matched pairs test (IL-6) was performed to verify changes from baseline (intra-group changes), and data were expressed as means with their standard errors. All statistical analyses were performed using IBM SPSS windows version 22<sup>(26)</sup>. Tests were two-tailed, and an  $\alpha$  level of 0.05 indicated statistically significant results.

**Results**

All participants completed the study after 90 d. There were no differences between the groups without the MetS and with the MetS in relation to age, sex, ethnicity, smoking habit, physical exercise and the number of patients with diabetes mellitus and insulin resistance. The majority of the patients (58%) from the MetS group presented with hypertension ( $P < 0.001$ ), whereas this characteristic was not seen in any subject of the control group (Table 1). A total of three patients in each group used statins, whereas none of the patients in the control group and six patients in the MetS group used anti-hypertensive medications (data not shown).

There were no significant inter- or intra-group differences observed in SBP, DBP, hsCRP, SH and leptin levels (Table 2). On the other hand, in relation to the baseline values (intra-group changes), subjects in both groups showed a decrease in homocysteine ( $P = 0.02$  and  $P = 0.03$ , respectively), hydroperoxides ( $P = 0.02$  and  $P = 0.01$ , respectively) and IL-6 levels ( $P = 0.02$ ). In addition, increased adiponectin and NOx levels were observed only in the group with the MetS (Fig. 1(a) and (b)). In contrast, only the individuals without the MetS had increases in TRAP levels ( $P = 0.002$ ) (Fig. 1(c)).

**Table 1.** Characteristics of the study population\* (Numbers and percentages; mean values with their standard errors)

Parameters	MetS <sup>-</sup> (n 14)		MetS <sup>+</sup> (n 19)		P
	n	%	n	%	
Age (years)					0.059
Mean	41.77		48.05		
SEM	2.66		1.43		
Sex					0.485
Female	10	71.4	11	57.9	
Male	4	28.6	8	42.1	
Ethnicity					0.620
Caucasian	13	92.9	16	84.2	
Non-Caucasian	1	7.1	3	15.8	
Smoking (yes)	1	7.1	3	15.8	0.620
Physical exercise (yes)	2	14.2	0	0.00	0.172
Hypertension (yes)	0	0.0	11	57.9	<0.001
DM (yes)	0	0.0	5	26.3	0.057
IR (yes)	6	46.2	14	77.8	0.128

MetS<sup>-</sup>, subjects without the metabolic syndrome; MetS<sup>+</sup>, subjects with the metabolic syndrome; DM, diabetes mellitus; IR, insulin resistance (homeostatic model assessment for insulin resistance  $\geq 2.5$ ).

\* Student’s *t* test was used.

**Table 2.** Effects of consuming *Bifidobacterium lactis* HN019 in subjects with the metabolic syndrome (MetS, MetS<sup>+</sup>) and without the MetS (MetS<sup>-</sup>) evaluated at the beginning of the experiment (T0) and after 90 d (T90) (Mean values with their standard errors)


	MetS <sup>-</sup> (n 14)				P*	MetS <sup>+</sup> (n 19)				P*
	T0		T90			T0		T90		
	Mean	SEM	Mean	SEM		Mean	SEM	Mean	SEM	
WC (cm)	98.42	4.47	95.71	4.56	0.126	106.29	3.02	105.88	3.00	0.775
SBP (mmHg)	117.14	3.69	120.71	3.70	0.355	142.35	5.03	140.58	5.52	0.654†
DBP (mmHg)	72.85	2.44	78.57	2.74	0.055	91.17	2.82	94.70	4.03	0.320†
Homocysteine (μmol/l)	9.46	0.48	8.15	0.67	0.015	11.97	0.82	10.94	0.86	0.031
hsCRP (mg/l)	3.50	0.70	3.08	0.68	0.128†	7.94	1.85	5.88	1.16	0.752†
SH (μM)	0.289	0.016	0.318	0.013	0.113	0.297	0.006	0.299	0.014	0.878
Hydroperoxide (cpm)	163.94	8.36	138.42	5.08	0.021	171.62	7.31	155.44	5.92	0.010
Leptin (ng/ml)	17.56	4.14	20.87	5.45	0.302†	17.92	2.79	15.26	3.59	0.184†
IL-6 (pg/ml)	5.43	1.6	1.54	0.38	0.018‡	4.70	1.06	2.12	0.50	0.016‡

WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; hsCRP, high-sensitivity C-reactive protein; SH, sulfhydryl groups of proteins; cpm, counts per million.

\* Paired Student's *t* test was used.

† Variables processed in logarithmic transformation.

‡ Non-parametric paired test: Wilcoxon.



**Fig. 1.** Concentrations of adiponectin (a), nitric oxide metabolites (NOx) (b) and total radical-trapping antioxidant parameter (TRAP) (c) in patients with the metabolic syndrome (MetS<sup>+</sup>) and without (MetS<sup>-</sup>), evaluated at the beginning of the study (T0, □) and after 90 d (T90, ■) of probiotic milk ingestion. The *t* test was performed to compare differences between parameters of groups at baseline and differences across treatment groups (intergroup changes). Values are means, with their standard errors represented by vertical bars.

## Discussion

### Principal findings

The main finding of this study was the beneficial effects of *B. lactis* HN019 on adiponectin and NOx levels in the group of patients with the MetS and on TRAP in the group of patients without the MetS. In addition, *B. lactis* HN019 showed beneficial effects on homocysteine, hydroperoxides and IL-6 independently of the presence of the MetS.

### Beneficial effects of *Bifidobacterium lactis* HN019 in patients with and without the metabolic syndrome

Although inflammation has been recently considered as a mechanism of immune defence and repair<sup>(27)</sup>, the intriguing pro- and/or anti-inflammatory actions of IL-6 is still a matter of discussion<sup>(28,29)</sup>. In the present study, it is feasible to suggest that the reduction in IL-6, which occurred together with the decrease in homocysteine and hydroperoxide levels, can be considered a beneficial effect. Our group has performed some studies that also reported that the consumption of *Lactobacillus*

*plantarum*<sup>(30)</sup> and *B. lactis*<sup>(16)</sup> reduced IL-6 levels in patients with the MetS. In addition, the reduction of cytokine IL-6 corroborates the study of Cani *et al.*<sup>(31)</sup> that associated the decrease of pro-inflammatory cytokines with the increased levels of *Bifidobacterium* species in high-fat-diet-induced diabetes in mice.

We are not aware of any study to date that has evaluated oxidative stress with probiotics in patients with the MetS. Probiotics have been shown to reduce pro-oxidant biomarkers in patients with diabetes mellitus type 2<sup>(32,33)</sup>. Ejtahed *et al.*<sup>(32)</sup> reported a reduction in malondialdehyde (MDA) levels using a yogurt enriched with *B. lactis* and *Lactobacillus acidophilus*, whereas Hariri *et al.*<sup>(33)</sup> found a decrease in 8-hydroxy-2'-deoxyguanosine levels using probiotic soya milk containing *L. plantarum*. In addition, in healthy volunteers with high levels of  $\gamma$ -glutamyltransferase, Kang *et al.*<sup>(34)</sup> showed a decrease in MDA levels with *Lactobacillus brevis*. Although we used a different methodology (CL-LOOH) to evaluate oxidative stress, our results with *B. lactis* in healthy subjects and patients with the MetS are in agreement with the aforementioned studies. In contrast, other reports have not shown changes in oxidative

stress in healthy aged individuals using the probiotic VSL#3, a multistrain cocktail composed of *Streptococcus thermophilus* and several species of *Lactobacillus* and bifidobacteria<sup>(35)</sup>, and in patients with diabetes mellitus type 2 using *L. acidophilus* and *B. lactis*<sup>(36)</sup>. The decrease in inflammation and oxidative stress by probiotics has been attributed to their effects on scavenging superoxide and hydroxyl radicals, decreasing expression of IL-6 in adipocytes and decreasing adiposity<sup>(37)</sup>.

Regarding homocysteine, studies have proposed that increased concentration of total plasma homocysteine could represent an additional independent risk factor for CVD<sup>(38)</sup>. A meta-analysis of prospective cohort studies demonstrated that, after accounting for known CVD risk, a 25% lower homocysteine level was associated with about an 11% lower risk of ischaemic heart disease and about a 19% lower risk of stroke<sup>(39)</sup>. Elevated homocysteine levels can occur owing to a lack of vitamins, in particular folate and/or cobalamin, which are required cofactors for homocysteine metabolism through the re-methylation pathway<sup>(40)</sup>. Milk is a moderate dietary source of folic acid<sup>(41,42)</sup>. Nevertheless, some micro-organisms that are important for fermented food and/or gut microbiota, such as bifidobacteria, can synthesise folate *de novo*<sup>(43–45)</sup>. The recommended daily intake of folate is 300 µg for men, 400 µg for women and 500 µg for lactating or pregnant women. However, these levels are not easy to reach<sup>(45)</sup>. Milk provides 20–50 µg folate/l, whereas in fermented milk this amount increases to 110 µg/l<sup>(41)</sup>. In this study, the beneficial effect found on homocysteine levels in both groups permits one to suggest that even a moderate quantity of folic acid present in probiotic milk can decrease this CVD risk factor. This is the first study, to our knowledge, to show decreased homocysteine levels with a probiotic *Bifidobacterium* species. However, other studies performed with *L. plantarum*<sup>(30)</sup> and *L. acidophilus*<sup>(46)</sup> have already reported this finding.

#### Link between the effects of homocysteine and oxidative stress

The association between homocysteine and oxidative stress has been shown in patients with type 2 diabetes mellitus, hypertension, atherosclerosis and the MetS<sup>(47,48)</sup>. Tyagi<sup>(47)</sup> showed that homocysteine induces oxidative stress by increasing nitric oxide synthase while decreasing endothelial nitric oxide synthase (eNOS) in cardiac microvascular endothelial cells. In the present study, the effects of probiotics decreasing concomitantly homocysteine and hydroperoxide levels seem to reinforce this association.

#### Beneficial effects of Bifidobacterium lactis HN019 only demonstrated in patients with the metabolic syndrome

The adipose tissue also secretes adiponectin, a protein showing anti-inflammatory activity, which inhibits TNF- $\alpha$  production, adhesion molecule expression and nuclear transcriptional factor  $\kappa$ B signalling, a pivotal pathway in inflammatory reactions in endothelial cells. In addition, adiponectin is anti-atherogenic and is an insulin-sensitising agent<sup>(49,50)</sup>. Obesity, especially

visceral obesity, may lead to a decreased secretion of adiponectin through feedback inhibition, thereby suppressing the beneficial effects of adiponectin on insulin sensitivity, and decreased adiponectin levels have been found in patients with obesity, type 2 diabetes mellitus, arterial hypertension and the MetS<sup>(51,52)</sup>. Similarly to the results obtained in the current work, adiponectin levels also increased in obese subjects using *Lactobacillus gasseri*<sup>(53)</sup>. In addition, an experimental study showed an increase in adiponectin levels with *Lactobacillus rhamnosus* supplementation in high-fat-diet-fed mice<sup>(54)</sup>.

The mechanisms by which probiotics increase adiponectin levels are largely unknown. Although it is conceivable to suggest an effect related to the decrease in pro-inflammatory cytokines, such as IL-6, demonstrated in the present study, other mechanistic insights cannot be discharged. For example, conjugated linoleic acid is produced by some species of *Lactobacillus* (e.g. *acidophilus*, *plantarum*, *paracasei* and *casei*) and has the potential to act as a PPAR $\gamma$  agonist, which up-regulates adiponectin and down-regulates inflammation and adiposity<sup>(55)</sup>.

Although endothelial dysfunction has been considered an important issue in patients with the MetS, results of studies on serum NOx levels in these patients have been contradictory<sup>(56–58)</sup>. The majority of the studies support the hypothesis that even in low-grade inflammatory processes, such as in the MetS, NO is consumed in a reaction with superoxide anion yielding a strong oxidant species, ONOO<sup>-</sup>, which in turn accelerates the lipid peroxidation reaction<sup>(59)</sup>. The increase in NOx levels with *B. lactis* is coherent with both improvement in endothelium dysfunction and decreased oxidative stress found in this study. Other reports also showed an increase in NOx after utilisation of probiotics in rats; VSL#3 prevented endothelial dysfunction in rats with portal hypertension<sup>(60)</sup>, whereas kefir, a probiotic containing different strains, restored the production and release of NOx in streptozocin-induced diabetic rats<sup>(61)</sup>. Esposito *et al.*<sup>(62)</sup> reported that VSL#3 reduced several pro-inflammatory and pro-oxidant biomarkers, including NF- $\kappa$ B, in rats with non-alcoholic fatty liver disease, a condition strongly associated with the MetS. In addition, the same authors reported that probiotics increased PPAR $\alpha$  expression, which explained the important anti-inflammatory and antioxidant effects on O&NS demonstrated with VSL#3.

#### Link between the effects of adiponectin and nitric oxide

Epidemiological studies based on different ethnic groups have identified adiponectin deficiency as an independent risk factor for endothelial dysfunction, hypertension, CHD, myocardial infarction and other cardiovascular complications. On the other hand, decreased bioavailability and/or impaired production of NO is considered a major cause of endothelial dysfunction, which is now recognised as one of the earliest changes in atherosclerosis<sup>(63,64)</sup>.

In experimental studies, adiponectin has shown to enhance NO production in cultured aortic endothelial cells<sup>(65)</sup>, to significantly increase eNOS (83%) and reduce inducible nitric oxide synthase (70%) in hyperlipidaemic rats<sup>(66)</sup>, and to improve obesity-related hypertension in mice<sup>(67)</sup>. Adiponectin



has anti-inflammatory effects and augments blood flow by enhancing nitric oxide production and activating eNOS, and it may act as a modulator of vascular remodelling by suppressing smooth muscle cell migration, which possibly plays a role in the regulation of atherosclerosis<sup>(68)</sup>. In a previous study, our group has already demonstrated that the association of hypoadiponectinaemia and increased NOx is one of the main reasons of decreasing blood pressure in patients with the MetS using fish oil and soya<sup>(69)</sup>.

### *Beneficial effects of Bifidobacterium lactis HN019 only demonstrated in subjects without the metabolic syndrome*

Several studies have shown the increase in antioxidant defences with probiotics in healthy people and diabetic patients, but, to our knowledge, this is the first study that reported the antioxidant status in patients with the MetS using probiotics. In healthy subjects, significant increases in antioxidants have been seen with different strains of *Lactobacillus*<sup>(34,37,70,71)</sup> and with different methodologies, such as total antioxidant capacity (TAC)<sup>(70,71)</sup>, superoxide dismutase (SOD)<sup>(34)</sup> and total GSH<sup>(72)</sup>. Similarly, in diabetic patients, these increases have been reported not only with *L. plantarum*<sup>(33)</sup> but also with *L. acidophilus* associated with *B. lactis*<sup>(32)</sup> and cocktails that used some strains of *Lactobacillus*, *Bifidobacterium* and *S. thermophilus* (VSL#3)<sup>(37)</sup>; these beneficial effects have been demonstrated by increased TAC<sup>(32)</sup>, SOD<sup>(32,33)</sup> and GSH<sup>(37)</sup>. On the other hand, a study using VSL#3 in diabetic patients did not find significant changes in TAC<sup>(36)</sup>. The antioxidant action of *Lactobacillus* strains has been attributed to the antioxidative effect of bioactive peptides released during fermentation by proteolytic lactic acid bacteria, the expression of MnSOD activity, elimination of hydroxyl and peroxyl radicals and as having the complete glutathione system necessary for glutathione recycling, transporting and synthesis<sup>(32,33)</sup>. However, the mechanism by which *Bifidobacterium* strains decreased TAC has still to be elucidated. Unexpectedly, in the current study, antioxidant defences only improved in healthy subjects, but not in the MetS patients. Nevertheless, it is feasible to suggest that lack of TRAP increase in the group with the MetS could be due to the presence of higher body fat, which becomes difficult because of adequate responses of the antioxidant defence mechanisms<sup>(73)</sup>.

### *Limitations and strengths*

There are some limitations that must be considered in this study. The first limitation is the small number of subjects in the groups. Second, this study does not allow extrapolation of the results with different strains, especially the data on oxidative stress, which can be highly specific<sup>(36)</sup>. However, some strengths of the study can be pointed out. The study is original under some aspects, as follows: first, this is a unique report with probiotics, in which oxidative stress was studied in patients with the MetS; second, even in patients with diabetes mellitus type 2, studies with *B. lactis* are still scarce; third, the original design of the study compares healthy subjects and patients with the MetS using the same fermented milk with probiotics.

In conclusion, *B. lactis* HN019 has several beneficial effects on inflammatory and oxidative biomarkers in healthy subjects and the MetS patients supplemented with the same probiotics as our previous study, but for a longer period. Patients with the MetS showed a specific improvement in adiponectin and NOx levels, whereas a specific favourable effect was shown in the antioxidant defences in healthy subjects. If the results obtained in the present study are confirmed, discussion on supplementation of fermented milk with probiotics in healthy subjects and diseases in which insulin resistance is the *pièce de résistance* may be warranted.

### **Acknowledgements**

The authors thank the CONFEPAR – Agro-Industrial Cooperativa Central for milk donation, Duas Rodas Industrial for flavors donation and the PROSUP – Programa de Suporte à Pós-Graduação de Instituições de Ensino Particulares – for financial support to L. G. S.

This work was supported by Fundação Nacional de Desenvolvimento de Ensino Superior Particular.

L. J. B. was responsible for recruiting the patients, interpretation of the results and the writing of the manuscript. C. H. B. d. S. and L. G. S. were responsible for writing of the manuscript; D. F. A. was responsible for recruiting the patients and the laboratory analysis. A. N. C. S. and I. D. were responsible for interpretation of the results and the writing of the manuscript. G. N. C. was responsible for the original concept of the study, the study design, interpretation of the results and the writing of the manuscript. All authors read and approved the final manuscript.

The authors declare that there are no conflicts of interest.

### **References**

1. Grundy SM, Cleeman JI, Daniels SR, *et al.* (2005) Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* **112**, 2735–2752.
2. Lakka H, De L, Ta L, *et al.* (2002) The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* **288**, 2709–2716.
3. Kahn R, Buse J, Ferrannini E, *et al.* (2005) The metabolic syndrome: time for a critical appraisal: joint statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* **28**, 2289–2304.
4. Cornier MA, Dabelea D, Hernandez TL, *et al.* (2008) The metabolic syndrome. *Endocr Rev* **29**, 777–822.
5. Erejuwa OO, Sulaiman SA & Ab Wahab MS (2014) Modulation of gut microbiota in the management of metabolic disorders: the prospects and challenges. *Int J Mol Sci* **15**, 4158–4188.
6. Caricilli A & Saad M (2013) The role of gut microbiota on insulin resistance. *Nutrients* **5**, 829–851.
7. Shen J, Obin MS & Zhao L (2013) The gut microbiota, obesity and insulin resistance. *Mol Aspects Med* **34**, 39–58.
8. Bäckhed F, Ley RE, Sonnenburg JL, *et al.* (2005) Host-bacterial mutualism in the human intestine. *Science* **307**, 1915–1920.
9. Forsythe P, Sudo N, Dinan T, *et al.* (2010) Mood and gut feelings. *Brain Behav Immun* **24**, 9–16.
10. Jones RM, Mercante JW & Neish AS (2012) Reactive oxygen production induced by the gut microbiota: pharmacotherapeutic implications. *Curr Med Chem* **19**, 1519–1529.



11. Carding S, Verbeke K, Vipond DT, *et al.* (2015) Dysbiosis of the gut microbiota in disease. *Microb Ecol Health Dis* **26**, 26191.
12. Nagpal R, Kumar A, Kumar M, *et al.* (2012) Probiotics, their health benefits and applications for developing healthier foods: a review. *FEMS Microbiol Lett* **334**, 1–15.
13. Miglioranza Scavuzzi B, Miglioranza LH, Henrique FC, *et al.* (2015) The role of probiotics on each component of the metabolic syndrome and other cardiovascular risks. *Expert Opin Ther Targets* **19**, 1127–1138.
14. Salehi-Abargouei A, Ghasvand R & Hariri M (2017) Prebiotics, prosynbiotics and synbiotics: can they reduce plasma oxidative stress parameters? A systematic review. *Probiotics Anti-microb Proteins* **9**, 1–11.
15. Mirmiran P (2014) Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: a review. *World J Diabetes* **5**, 267.
16. Bernini LJ, Simão ANC, Alfieri DF, *et al.* (2016) Beneficial effects of *Bifidobacterium lactis* on lipid profile and cytokines in patients with metabolic syndrome: a randomized trial. Effects of probiotics on metabolic syndrome. *Nutrition* **32**, 716–719.
17. Brasil (2007) Ministério da Agricultura Pecuária e Abastecimento. Instrução Normativa nº 46, de 23 de Outubro de 2007. Regulamento Técnico de Identidade e Qualidade de Leites Fermentados. Diário Of da República Fed do Bras Brasília DF, 24 out. 2007. Seção 1, 5 (Technical Regulation on the Identity and Quality of Fermented Milks). <http://www.cidasc.sc.gov.br/inspecao/files/2012/08/instru%C3%87%C3%83o-normativa-n%C2%BA-46-de-23-de-outubro-de-2007.pdf>
18. Brasil (2001) Ministério da Saúde. Agência Nacional de Vigilância Sanitária - ANVISA Resolução RDC nº 12, de 02/01/2001. M. Regulamento Técnico sobre padrões microbiológicos para alimentos. Diário Of da República Fed do Bras Brasília DF, 10 January 2001. Seção 1, 45–53 (Technical Regulation on microbiological standards for food). [http://portal.anvisa.gov.br/documents/33880/2568070/RDC\\_12\\_2001.pdf/15ffddf6-3767-4527-bfac-740a0400829b](http://portal.anvisa.gov.br/documents/33880/2568070/RDC_12_2001.pdf/15ffddf6-3767-4527-bfac-740a0400829b)
19. Pickering TG, Hall JE, Appel LJ, *et al.* (2005) Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation* **111**, 697–716.
20. Gonzalez Flecha B, Llesuy S & Boveris A (1991) Hydroperoxide-initiated chemiluminescence: an assay for oxidative stress in biopsies of heart, liver, and muscle. *Free Radic Biol Med* **10**, 93–100.
21. Guevara I, Iwanejko J, Dembińska-Kieć A, *et al.* (1998) Determination of nitrite/nitrate in human biological material by the simple Griess reaction. *Clin Chim Acta* **274**, 177–188.
22. Navarro-González JA, García-Benayas C & Arenas J (1998) Semiautomated measurement of nitrate in biological fluids. *Clin Chem* **44**, 679–681.
23. Repetto M, Reides C & Gomez Carretero ML (1996) Oxidative stress in blood of HIV infected patients. *Clin Chim Acta* **255**, 107–117.
24. Venturini D, Simão ANC, Sripes NA, *et al.* (2012) Evaluation of oxidative stress in overweight subjects with or without metabolic syndrome. *Obesity* **20**, 2361–2366.
25. Hu ML (1994) Measurement of protein thiol groups and glutathione in plasma. *Methods Enzymol* **233**, 380–385.
26. IBM Corp. (2013) *IBM SPSS Statistics for Windows, Version 22.0*. Armonk, NY: IBM Corp.
27. Lu L, Yu Y, Guo Y, *et al.* (2015) Transcriptional modulation of intestinal innate defense/inflammation genes by preterm infant microbiota in a humanized gnotobiotic mouse model. *PLOS ONE* **10**, e0124504.
28. Eder K, Baffy N, Falus A, *et al.* (2009) The major inflammatory mediator interleukin-6 and obesity. *Inflamm Res* **58**, 727–736.
29. Scheller J, Chalaris A, Schmidt-Arras D, *et al.* (2011) The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochim Biophys Acta* **1813**, 878–888.
30. Barreto FM, Colado Simão AN, Morimoto HK, *et al.* (2014) Beneficial effects of *Lactobacillus plantarum* on glycemia and homocysteine levels in postmenopausal women with metabolic syndrome. *Nutrition* **30**, 939–942.
31. Cani PD, Neyrinck AM, Fava F, *et al.* (2007) Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia* **50**, 2374–2383.
32. Ejtahed HS, Mohtadi-Nia J, Homayouni-Rad A, *et al.* (2012) Probiotic yogurt improves antioxidant status in type 2 diabetic patients. *Nutrition* **28**, 539–543.
33. Hariri M, Salehi R, Feizi A, *et al.* (2015) The effect of probiotic soy milk and soy milk on anthropometric measures and blood pressure in patients with type II diabetes mellitus: a randomized double-blind clinical trial. *ARYA Atheroscler* **11**, 1–8.
34. Kang YM, Lee BJ, Kim JI, *et al.* (2012) Antioxidant effects of fermented sea tangle (*Laminaria japonica*) by *Lactobacillus brevis* BJ20 in individuals with high level of  $\gamma$ -GT: a randomized, double-blind, and placebo-controlled clinical study. *Food Chem Toxicol* **50**, 1166–1169.
35. Valentini L, Pinto A, Bourdel-Marchasson I, *et al.* (2015) Impact of personalized diet and probiotic supplementation on inflammation, nutritional parameters and intestinal microbiota – the ‘RISTOMED project’: randomized controlled trial in healthy older people. *Clin Nutr* **34**, 593–602.
36. Tonucci LB, Olbrich dos Santos KM, Licursi de Oliveira L, *et al.* (2017) Clinical application of probiotics in type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled study. *Clin Nutr* **36**, 85–92.
37. Asemi Z, Zare Z, Shakeri H, *et al.* (2013) Effect of multispecies probiotic supplements on metabolic profiles, hs-CRP, and oxidative stress in patients with type 2 diabetes. *Ann Nutr Metab* **63**, 1–9.
38. Fruchart J-C (2004) New risk factors for atherosclerosis and patient risk assessment. *Circulation* **109**, Suppl. 23, III-15–III-19.
39. Homocysteine Studies Collaboration (2002) Homocysteine and risk of ischemic heart disease and stroke: a meta-analysis. *JAMA* **288**, 2015–2022.
40. Selhub J (2006) The many facets of hyperhomocysteinemia: studies from the Framingham cohorts. *J Nutr* **136**, 1726–1730.
41. Sybesma W, Starrenburg M, Tijsseling L, *et al.* (2003) Effects of cultivation conditions on folate production by lactic acid bacteria. *Appl Environ Microbiol* **69**, 4542–4548.
42. Konstantinova SV, Vollset SE, Berstad P, *et al.* (2007) Dietary predictors of plasma total homocysteine in the hordaland homocysteine study. *Br J Nutr* **98**, 201–210.
43. Deguchi Y, Morishita T & Mutai M (1985) Comparative studies on synthesis of water-soluble vitamins among human species of bifidobacteria. *Agric Biol Chem* **49**, 13–19.
44. Camilo E, Zimmerman J, Mason JB, *et al.* (1996) Folate synthesized by bacteria in the human upper small intestine is assimilated by the host. *Gastroenterology* **110**, 991–998.
45. D’Aimmo MR, Mattarelli P, Biavati B, *et al.* (2012) The potential of bifidobacteria as a source of natural folate. *J Appl Microbiol* **112**, 975–984.
46. Mohammad MA, Molloy A, Scott J, Hussein L, *et al.* (2006) Plasma cobalamin and folate and their metabolic markers

- methylnalonic acid and total homocysteine among Egyptian children before and after nutritional supplementation with the probiotic bacteria *Lactobacillus acidophilus* in yoghurt matrix. *Int J Food Sci Nutr* **57**, 470–480.
47. Tyagi N (2005) Mechanisms of homocysteine-induced oxidative stress. *Am J Physiol Heart Circ Physiol* **289**, H2649–H2656.
  48. Dominguez LJ, Galioto A, Pineo A, *et al.* (2010) Age, homocysteine, and oxidative stress: relation to hypertension and type 2 diabetes mellitus. *J Am Coll Nutr* **29**, 1–6.
  49. Ouchi N, Kihara S, Arita Y, *et al.* (2000) Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF-kappaB signaling through a cAMP-dependent pathway. *Circulation* **102**, 1296–1301.
  50. Hotta K, Funahashi T, Bodkin NL, *et al.* (2001) Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in rhesus monkeys. *Diabetes* **50**, 1126–1133.
  51. Santaniemi M, Kesäniemi YA & Ukkola O (2006) Low plasma adiponectin concentration is an indicator of the metabolic syndrome. *Eur J Endocrinol* **155**, 745–750.
  52. Simão ANC, Lozovoy MAB, Simão TNC, *et al.* (2012) Adiponectinemia is associated with uricemia but not with proinflammatory status in women with metabolic syndrome. *J Nutr Metab* **2012**, 418094.
  53. Kim SW, Park KY, Kim B, *et al.* (2013) *Lactobacillus rhamnosus* GG improves insulin sensitivity and reduces adiposity in high-fat diet-fed mice through enhancement of adiponectin production. *Biochem Biophys Res Commun* **431**, 258–263.
  54. Kadooka Y, Sato M, Imaizumi K, *et al.* (2010) Regulation of abdominal adiposity by probiotics (*Lactobacillus gasseri* SBT2055) in adults with obese tendencies in a randomized controlled trial. *Eur J Clin Nutr* **64**, 636–643.
  55. Nakamura YK & Omaye ST (2012) Metabolic diseases and pro- and prebiotics: mechanistic insights. *Nutr Metab* **9**, 60.
  56. Sun Y, Hu S, XH Z, *et al.* (2006) Plasma levels of vWF and NO in patients with metabolic syndrome and their relationship with metabolic disorders. *Zhejiang Da Xue Xue Bao Yi Xue Ban* **35**, 315–318.
  57. Zahedi Asl S, Ghasemi A & Azizi F (2008) Serum nitric oxide metabolites in subjects with metabolic syndrome. *Clin Biochem* **41**, 1342–1347.
  58. Simão ANC, Lozovoy MAB, Simão TNC, *et al.* (2011) Immunological and biochemical parameters of patients with metabolic syndrome and the participation of oxidative and nitroactive stress. *Braz J Med Biol Res* **44**, 707–712.
  59. Tao L, Gao E, Jiao X, *et al.* (2007) Adiponectin cardioprotection after myocardial ischemia/reperfusion involves the reduction of oxidative/nitrative stress. *Circulation* **115**, 1408–1416.
  60. Rashid SK, Khodja NI, Auger C, *et al.* (2014) Probiotics (VSL#3) prevent endothelial dysfunction in rats with portal hypertension: role of the angiotensin system. *PLOS ONE* **9**, e97458.
  61. Maciel FR, Punaro GR, Rodrigues AM, *et al.* (2016) Immunomodulation and nitric oxide restoration by a probiotic and its activity in gut and peritoneal macrophages in diabetic rats. *Clin Nutr* **35**, 1066–1072.
  62. Esposito E, Iacono A, Bianco G, *et al.* (2009) Probiotics reduce the inflammatory response induced by a high-fat diet in the liver of young rats. *J Nutr* **139**, 905–911.
  63. Goldstein BJ & Scalia R (2004) Adiponectin: a novel adipokine linking adipocytes and vascular function. *J Clin Endocrinol Metab* **89**, 2563–2568.
  64. Zhu W, Cheng KKY, Vanhoutte PM, *et al.* (2008) Vascular effects of adiponectin: molecular mechanisms and potential therapeutic intervention. *Clin Sci* **114**, 361–374.
  65. Tan KCB, Xu A, Chow WS, *et al.* (2004) Hypoadiponectinemia is associated with impaired endothelium-dependent vasodilation. *J Clin Endocrinol Metab* **89**, 765–769.
  66. Li R, Wang W-Q, Zhang H, *et al.* (2007) Adiponectin improves endothelial function in hyperlipidemic rats by reducing oxidative/nitrative stress and differential regulation of eNOS/iNOS activity. *Am J Physiol Endocrinol Metab* **293**, E1703–E1708.
  67. Ohashi K, Kihara S, Ouchi N, *et al.* (2006) Adiponectin replenishment ameliorates obesity-related hypertension. *Hypertension* **47**, 1108–1116.
  68. Ziemke F & Mantzoros CS (2010) Adiponectin in insulin resistance: lessons from translational research. *Am J Clin Nutr* **91**, 258–261.
  69. Simão ANC, Lozovoy MAB, Bahls LD, *et al.* (2012) Blood pressure decrease with ingestion of a soya product (kinako) or fish oil in women with the metabolic syndrome: role of adiponectin and nitric oxide. *Br J Nutr* **108**, 1435–1442.
  70. Kullisaar T, Songisepp E, Mikelsaar M, *et al.* (2003) Antioxidative probiotic fermented goats' milk decreases oxidative stress-mediated atherogenicity in human subjects. *Br J Nutr* **90**, 449.
  71. Songisepp E, Kals J, Kullisaar T, *et al.* (2005) Evaluation of the functional efficacy of an antioxidative probiotic in healthy volunteers. *Nutr J* **4**, 22.
  72. Asemi Z, Jazayeri S, Najafi M, *et al.* (2012) Effect of daily consumption of probiotic yogurt on oxidative stress in pregnant women: a randomized controlled clinical trial. *Ann Nutr Metab* **60**, 62–68.
  73. Chrysohoou C, Panagiotakos DB, Pitsavos C, *et al.* (2007) The implication of obesity on total antioxidant capacity in apparently healthy men and women: the ATTICA study. *Nutr Metab Cardiovasc Dis* **17**, 590–597.