

DEBATE

Comment on ‘Changing relationships between HIV prevalence and circumcision in Lesotho’, and ‘Age-incidence and prevalence of HIV among intact and circumcised men: an analysis of PHIA surveys in Southern Africa’

Brian J. Morris¹  and Joya Banerjee²

¹School of Medical Sciences, University of Sydney, Sydney, NSW, Australia and ²Be the Change Group, Vancouver, BC, Canada

Corresponding author: Brian J. Morris; Email: brian.morris@sydney.edu.au

(Received 3 July 2023; revised 18 March 2024; accepted 9 April 2024)

Abstract

Two articles by Garenne (2023a,b) argue that voluntary medical male circumcision does not reduce human immunodeficiency virus transmission in Africa. Here we point out key evidence and analytical flaws that call into question this conclusion.

Keywords: Human immunodeficiency virus/acquired immunodeficiency syndrome; circumcision; Southern Africa; reproductive health; voluntary medical male circumcision

Contrary findings ignored

In Garenne (2023b), six countries with circumcision ‘prevalence’ of 12.7–71.2% were studied. However, these prevalence figures incorrectly conflate voluntary medical male circumcision (VMMC) with traditional male circumcision (TMC), which can increase risk of human immunodeficiency virus (HIV) infection through use of unsterilised contaminated instruments on multiple youths, some already infected (Brewer *et al.* 2007, 2009; Ndiwane 2008). TMC of 16–20-year-olds is common in Lesotho where VMMC, but not TMC, is associated with HIV risk reduction (Coburn *et al.* 2013; Carrasco *et al.* 2020; Makatjane *et al.* 2016). These observations render the study’s finding of no association questionable.

Confounding

These studies ignore confounding from antiretroviral therapy (ART). ART was rolled out alongside VMMC in sub-Saharan Africa concurrently with VMMC. While ART can reduce HIV infection, it increases HIV prevalence because people who previously would have died from acquired immunodeficiency syndrome-related illnesses now live with HIV (Shafer *et al.* 2013; Zaidi *et al.* 2013). The articles should have documented incidence. A modelling study found VMMC was the third most effective intervention after ART and condoms (Johnson *et al.* 2022). Furthermore, uptake of VMMC has been high in young adolescent boys who are not yet sexually active (UNAIDS and WHO 2021), leading to a delay between VMMC and reduction in new HIV cases.

Data limitations

The primary data sources use self-reported circumcision status. Self-reporting is unreliable. This could affect any study using such data, but it seems to be a particular issue with Lesotho, which is a country Garenne focused on in the first paper we criticise. A study in Lesotho found that only half of men claiming to be circumcised actually were, and a further 26.6% were only partially circumcised (Thomas *et al.* 2011). Partial TMC is also common in Malawi (Renne *et al.* 2016).

Selective and misleading literature citations

The studies ‘cherry-pick’ often dated opinion pieces that fail to consider some of the issues described above. The articles ignore reviews of >30 studies linking VMMC to HIV risk reduction (Siegfried *et al.* 2003; Addanki *et al.* 2008). The studies also fail to point out that most subjects in Connolly *et al.* (2008) had TMC, not VMMC. And findings in Rosenberg *et al.* (2018) are attributed to self-selection, not ineffectiveness of VMMC.

Van Howe’s meta-regression analyses in 2015 are cited (Van Howe 2015), but not the detailed critique undermining his statistics (Morris *et al.* 2018). Van Howe’s reply in 2018 (Van Howe 2018) was rebutted (Morris *et al.* 2017). A meta-analysis (Van Howe 1999) was discredited (Moses *et al.* 1999; O’Farrell & Egger 2000) and became a textbook example of how not to do a meta-analysis (Borenstein *et al.* 2009). Yet the articles cite it (Garenne & Matthews 2019; Garenne 2023a,b). All other meta-analyses confirm VMMC is effective against female-to-male HIV transmission (O’Farrell & Egger 2000; Weiss *et al.* 2000; Byakika-Tusiime, 2008; Lei *et al.* 2015; Sharma *et al.* 2018; Farley *et al.* 2020) but these are ignored.

Conclusion

These recent articles on VMMC and HIV are problematic. The issues described above are well-known to researchers in the field but are ignored. Unfortunately, the articles are now being cited (Garenne 2023c) to support an opposition to VMMC. Most authorities find that VMMC is biologically- and cost-effective against HIV infection in Africa (Farley *et al.* 2020; Bershteyn *et al.* 2022; Bansi-Matharu *et al.* 2023).

Funding statement. None.

Competing interests. Brian J. Morris is a member of the Circumcision Academy of Australia, a not-for-profit, government registered, medical society that provides accurate, evidence-based information on male circumcision to parents, practitioners, and others, as well as contact details of doctors who perform the procedure in Australia and New Zealand. This author’s interest in male circumcision for disease prevention began after he co-invented and competitively patented the first use of PCR for viral detection, applying this technology to human papillomavirus detection in screening for cervical cancer, the prevalence of which is lower in women with circumcised male partners.

Ethical standard. Not applicable.

References

- Addanki KC, Pace DG and Bagasra O (2008) A practice for all seasons: male circumcision and the prevention of HIV transmission. *Journal of Infection in Developing Countries* 2(5), 328–334.
- Bansi-Matharu L, Mudimu E, Martin-Hughes R, Hamilton M, Johnson L, ten Brink D, Stover J, Meyer-Rath G, Kelly SL, Jamieson L, Cambiano V, Jahn A, Cowan FM, Mangenah C, Mavhu W, Chidarikire T, Toledo C, Revill P, Sundaram M, Hatzold K, Yansaneh A, Apollo T, Kalua T, Mugurungi O, Kiggundu V, Zhang S, Nyirenda R, Phillips A, Kripke K and Bershteyn A (2023) Cost-effectiveness of voluntary medical male circumcision for HIV prevention across sub-Saharan Africa: results from five independent models. *Lancet Global Health* 11(2), e244–e255.

- Bershteyn A, Mudimu E, Platais I, Mwalili S, Zulu JE, Mwanza WN and Kripke K** (2022) Understanding the evolving role of voluntary medical male circumcision as a public health strategy in Eastern and Southern Africa: opportunities and challenges. *Current HIV/AIDS Reports* **19**(6), 526–536.
- Borenstein M, Hedges L, Higgins JPT and Rothstein HR** (2009) *Introduction to Meta-Analysis*. John Wiley and Sons: West Sussex.
- Brewer DD, Potterat JJ, Roberts JM Jr and Brody S** (2007). Male and female circumcision associated with prevalent HIV infection in virgins and adolescents in Kenya, Lesotho, and Tanzania. *Annals of Epidemiology* **17**, 217–226.
- Brewer DD, Potterat JJ, Roberts JM and Brody S** (2009) Unhygienic male circumcision procedures and HIV transmission. *South African Medical Journal* **99**(1), 11.
- Byakika-Tusiime J** (2008) Circumcision and HIV infection: assessment of causality. *AIDS Behavior* **12**(6), 835–841.
- Carrasco MA, Rosen JG, Maile L, Manda R, Amzel A and Kiggundu V** (2020) Medically, traditionally, and dually circumcised men in Lesotho: population-based measurements of HIV/STI infections, sexual risk behaviors, and service use patterns. *AIDS and Behavior* **24**(7), 2112–2118.
- Coburn BJ, Okano JT and Blower S** (2013) Current drivers and geographic patterns of HIV in Lesotho: implications for treatment and prevention in Sub-Saharan Africa. *BMC Medicine* **11**, 224.
- Connolly C, Simbayi LC, Shanmugam R and Nqeketo A** (2008) Male circumcision and its relationship to HIV infection in South Africa: Results of a national survey in 2002. *South African Medical Journal (SAMJ)* **98**, 789–794.
- Farley TMM, Samuelson J, Grabowski MK, Ameyan W, Gray RH and Baggaley R** (2020) Impact of male circumcision on risk of HIV infection in men in a changing epidemic context – systematic review and meta-analysis. *Journal of the International AIDS Society* **23**, e25490.
- Garenne M** (2023a) Changing relationships between HIV prevalence and circumcision in Lesotho. *Journal of Biosocial Science* **55**(3), 463–478.
- Garenne M** (2023b) Age-incidence and prevalence of HIV among intact and circumcised men: an analysis of PHIA surveys in Southern Africa. *Journal of Biosocial Science* **55**(6), 1156–1168.
- Garenne M** (2023c) HIV prevention in Africa: is VMMC useful and acceptable? *International Journal of Impotence Research* **35**, 279–281.
- Garenne M and Matthews A** (2019) Voluntary medical male circumcision in Zambia: expectations and observations. *Journal of Biosocial Science* **52**(4), 560–572.
- Johnson LF, Meyer-Rath G, Dorrington RE, Puren A, Seathodi T, Zuma K and Feizzadeh A** (2022) The effect of HIV programs in South Africa on national HIV incidence trends, 2000–2009. *Journal of Acquired Immune Deficiency Syndrome* **90**(2), 115–123.
- Lei JH, Liu LR, Wei Q, Yan SB, Yang L, Song TR, Yuan HC, Lv X and Han P** (2015) Circumcision status and risk of HIV acquisition during heterosexual intercourse for both males and females: a meta-analysis. *PLoS One* **10**(5), e0125436.
- Makatjane T, Hlabana T and Letete E** (2016) *Male circumcision and HIV in Lesotho: Is the Relationship Real or Spurious? Analysis of the 2009 Demographic and Health Survey*. Rockville, Maryland, USA: ICF International.
- Morris BJ, Barboza G, Wamai RG and Krieger JN** (2017) Expertise and ideology in statistical evaluation of circumcision for protection against HIV infection. *World Journal of AIDS* **7**(3), 179–203.
- Morris BJ, Barboza G, Wamai RG and Krieger JN** (2018). Circumcision is a primary preventive against HIV infection: critique of a contrary meta-regression analysis by Van Howe. *Global Public Health* **13**(12), 1889–1899.
- Moses S, Nagelkerke NJ and Blanchard J** (1999) Analysis of the scientific literature on male circumcision and risk for HIV infection. *International Journal of STD & AIDS* **10**(9), 626–628.
- Ndiwane A** (2008). Laying down the knife may decrease risk of HIV transmission: cultural practices in Cameroon with implications for public health and policy. *Journal of Cultural Diversity* **15**, 76–80.
- O'Farrell N and Egger M** (2000) Circumcision in men and the prevention of HIV infection: a 'meta-analysis' revisited. *International Journal of STD & AIDS* **11**, 137–142.
- Renne S, Perry B, Corneli A, Chilungo A and Umar E** (2016) Perceptions of voluntary medical male circumcision among circumcising and non-circumcising communities in Malawi. *Global Public Health* **10**(5–6), 679–691.
- Rosenberg MS, Gómez-Olivé FX, Rohr JK, Kahn K and Bärnighausen TW** (2018) Are circumcised men safer sex partners? Findings from the HAALSI cohort in rural South Africa. *PLoS One* **13**(8), e0201445.
- Shafer LA, Nsubuga RN, Chapman R, O'Brien K, Mayanja BN and White RG** (2013) The dual impact of antiretroviral therapy and sexual behaviour changes on HIV epidemiologic trends in Uganda: a modelling study. *Sexually Transmitted Infections* **90**, 423–429.
- Sharma SC, Raison N, Khan S, Shabbir M, Dasgupta P and Ahmed K** (2018) Male circumcision for the prevention of human immunodeficiency virus (HIV) acquisition: a meta-analysis. *BJU International* **121**, 515–526.
- Siegfried N, Muller M, Volmink J, Deeks JJ, Egger M, Low NN, Weiss HH, Walker SA and Williamson PR** (2003). Male circumcision for prevention of heterosexual acquisition of HIV in men. *Cochrane Database of Systematic Reviews* **3**, CD003362.

- Thomas AG, Tran BR, Cranston M, Brown MC, Kumar R and Tlelai M** (2011) Voluntary medical male circumcision: a cross-sectional study comparing circumcision self-report and physical examination findings in Lesotho. *PLoS One* **6**(11), e27561.
- UNAIDS and WHO** (2021) *Voluntary Medical Male Circumcision*. Geneva, Switzerland: UNAIDS & WHO.
- Van Howe RS** (1999) Circumcision and HIV infection: review of the literature and meta-analysis. *International Journal of STD & AIDS* **10**(1), 8–16.
- Van Howe RS** (2015). Circumcision as a primary HIV preventive: extrapolating from the available data. *Global Public Health* **10**(5–6), 607–625.
- Van Howe RS** (2018). Expertise or ideology? A response to Morris *et al.* 2016, ‘circumcision is a primary preventive against HIV infection: critique of a contrary meta-regression analysis by Van Howe’. *Global Public Health* **13**(12), 1900–1918.
- Weiss HA, Quigley MA and Hayes RJ** (2000) Male circumcision and risk of HIV infection in sub-Saharan Africa: a systematic review and meta-analysis. *AIDS* **14**(15), 2361–2370.
- Zaidi J, Grapsa E, Tanser F, Newell M-L and Bärnighausen T** (2013) Dramatic increases in HIV prevalence after scale-up of antiretroviral treatment: a longitudinal population-based HIV surveillance study in rural Kwazulu-Natal. *AIDS* **27**(14), 2301–2305.

Cite this article: Morris BJ and Banerjee J (2024). Comment on ‘Changing relationships between HIV prevalence and circumcision in Lesotho’, and ‘Age-incidence and prevalence of HIV among intact and circumcised men: an analysis of PHIA surveys in Southern Africa’. *Journal of Biosocial Science*. <https://doi.org/10.1017/S0021932024000208>