

## EPV0740

### Comparative study of the effectiveness of EMDR G-TPE and CBT group protocols for the treatment of trauma in children exposed to a conflict context

E. Dozio<sup>1\*</sup>, C. Bizouerne<sup>2</sup> and V. Wamba<sup>3</sup>

<sup>1</sup>Action contre la Faim; <sup>2</sup>Indipendent, Paris, France and <sup>3</sup>Action contre la Faim, Bangui, Central African Republic

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1378

**Introduction:** 357 million children live in conflict zones. Children's mental health is a major but complex issue as needs and interventions depend on the age of the child, caregivers, daily safety and protection, etc. EMDR and CBT are the recommended therapies to treat PTSD according to the WHO, but there is not enough standardized evidence-based protocol for children. Testing and evaluating trauma management systems for children is essential for trauma treatment interventions to be implemented in emergency contexts, such as war and conflict situations.

**Objectives:** This research compares two intervention protocols for children aged 6 to 17 years suffering from Post Traumatic Stress Disorder after exposure to conflict traumatic events in the Central African Republic: the protocol "Kôno" developed by Action contre la Faim, based on a CBT and narrative approaches and the EMDR/G-TPE (Group Traumatic Events Protocol). After a psychoeducation session, the children were assigned to the ACF-KONO or EMDR/G-TPE groups for 5 sessions. The Child Psychosocial Distress Screener (CPDS) to measure general well-being and functioning and the Child Revised Impact of Events Scale (CRIES) to assess trauma, were administered before and after treatment.

**Methods:** 793 children participated in the research, 391 were included in the ACF-KONO protocol and 402 in the EMDR/G-TPE protocol. Both protocols have been shown to be equally effective in improving well-being and reducing traumatic symptoms. 185 children (90 ACF-KONO and 95 EMDR/G-TPE) were also re-evaluated after 5 months. The CPDS and CRIES-8 measurements reveal that the results are stable over time, with the use of both protocols. Detailed results will be presented.

**Results:** This research contributes to the discussion on a framework for group protocols for children in mental health and psychosocial support interventions in humanitarian programs. The two protocols tested showed very good results in reducing symptoms of PTSD in children. How to choose between EMDR/G-TPE and CBT? What contextual and cultural adaptations are to be expected? Are there differences in children's appreciation? And in that of mental health practitioners? Ideas for reflection will be shared.

**Conclusions:** It is possible to widen access to therapeutic treatment of PTSD for children in emergency situations such as war and conflict. Further research in other contexts is needed. These studies should explore qualitative elements, such as children's appreciations, but also the impact of these different protocols on the vicarious trauma of professionals involved in the treatment of children's trauma.

**Disclosure of Interest:** None Declared

## EPV0742

### Post-traumatic stress disorder after childbirth: A Tunisian study

N. Sghaier<sup>1\*</sup>, R. Ben Soussia<sup>1</sup>, H. Bouchahda<sup>2</sup>, I. Belhadj<sup>2</sup>, H. Ben Garouia<sup>3</sup> and H. Khiari<sup>3</sup>

<sup>1</sup>Psychiatry Departement; <sup>2</sup>Gynecology and obstetrics Department, Taher Sfar Hospital, Mahdia and <sup>3</sup>Psychiatry Departement, Ibn Jazzar Hospital, Kairouan, Tunisia

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1379

**Introduction:** Childbirth is a special time for every woman, bringing pregnancy to an end and marking the birth of a new baby. This transitional event presents countless physical and psychological changes. Post-traumatic stress disorder (PTSD), the result of particularly intense stress, is often linked to the perception of childbirth as a traumatic event, requiring optimized follow-up and screening.

**Objectives:** The aim of this current study is to estimate the prevalence of post-partum post-traumatic stress disorder in a sample group of Tunisian women and to determine factors associated with childbirth-related post-traumatic stress disorder.

**Methods:** This is a longitudinal prospective descriptive study conducted among women hospitalized for childbirth in the obstetrics and gynecology department and those who consulted the prenatal outpatient clinic at Taher Sfar Mahdia Hospital. The duration of the study is 7 months, from March 15, 2020 to September 15, 2020. Data collection was based on a pre-established questionnaire determining the various socio-demographic and clinical characteristics. Psychometric assessment was carried out using the Posttraumatic Stress Disorder Checklist Scale (PCL-S).

**Results:** We enrolled 120 women with a mean age of 28.2±5.3 years. Few women had a psychiatric history of depression (1.2%) or anxiety (3%), and 29% had a pathological obstetric history. Nevertheless, 12.5% of patients were hospitalized during pregnancy. Eighty-seven patients expressed anticipatory fear of childbirth, and 102 women had good marital and social support. Almost half of deliveries (48.3%) were vaginal, and almost a third (27.5%) were emergency caesarean sections. Level 3 pain was reported in 73.3% of cases. Psychometric assessment revealed a prevalence of PTSD of 5.8%, with PTSD symptomatology in 18.4% of women. PTSD was statistically associated with low level of education (p=0.02), postpartum complications (p=0.05) and gender of newborn (p=0.01).

**Conclusions:** Postpartum PTSD is a major public health problem affecting the healthy development of the newborn, the overall mental and physical recovery and well-being of the mother. Our findings suggest several intervention points for healthcare practitioners, including careful prenatal screening of past trauma history, social support, pain management and expectations about the birth, within a multidisciplinary approach.

**Disclosure of Interest:** None Declared