

Effect of supplementation of ewes with barley or maize during the last week of pregnancy on colostrum production

G. E. Banchemo¹, G. Quintans², A. Vazquez², F. Gigena², A. La Manna¹, D. R. Lindsay³
and J. T. B. Milton^{3†}

¹National Institute of Agricultural Research, INIA La Estanzuela, 70003, Ruta 50 Km 12, Uruguay; ²National Institute of Agricultural Research, INIA Treinta y Tres, 33000, Ruta 8 Km 281, Uruguay; ³School of Animal Biology, The University of Western Australia, Crawley, WA 6009, Australia

(Received 11 September 2006; Accepted 13 December 2006)

The experiment tested the hypothesis that short-term feeding of barley just before lambing would be as effective as maize in stimulating early production of colostrum. Both grains are high in starch, and should provide a substrate for lactose which, in turn, promotes lactogenesis. Thirty-five Corriedale ewes bearing single foetuses and 25 bearing twin-foetuses from a synchronised mating were fed on pasture during most of gestation. Fourteen days before the expected time of lambing the single- and twin-bearing ewes were allocated to three treatments and fed (1) a basal diet of lucerne hay to meet their nutrient requirements, (2) the basal diet plus a supplement of whole barley; or (3) the basal diet plus a supplement of cracked maize. The twin-bearing control ewes accumulated more colostrum than the single-bearing control ewes at birth (292 v. 190 g). However, supplementation with barley or maize increased the colostrum at birth to 360 and 541 g in singles and 648 and 623 g in twins. We conclude that barley is a good alternative to maize to stimulate production of colostrum especially in twin-bearing ewes whose lambs are the most likely to benefit from the supplement.

Keywords: lactose, lambs, pre-partum feeding, starch, survival

Introduction

Twin or single-bearing ewes may produce insufficient colostrum for their lambs even when grazing lush green pastures (Murphy *et al.*, 1996; McNeill *et al.*, 1998; Banchemo *et al.*, 2003). This may be due to the ewes' having a low intake of pasture in the last weeks of pregnancy (Forbes, 1968) or the low nutrient density of a diet based solely on pastures. Consequently, a supplement of grain in late pregnancy provides the ewe with a more concentrated diet and can overcome the limitations imposed by roughages alone (Hall *et al.*, 1992; Murphy *et al.*, 1996; Banchemo *et al.*, 2004a and b). Banchemo *et al.* (2004a) showed that supplementation of single- and twin-bearing ewes with cracked maize during the last 7 days of pregnancy more than doubled the amount of colostrum accumulated at lambing compared with that produced by unsupplemented, but otherwise well-fed ewes consuming a roughage diet. In those studies, the maize increased the lactose concentration of the colostrum because it increased the glucose supply to the udder. We argued that it is probably the starch in the maize that provided the substrate for

the lactose that improved production of colostrum and therefore barley, also high in starch, should be equally effective in stimulating early production of colostrum. Maize is not universally available in regions where sheep are bred, especially those with little summer rainfall so a more accessible grain like barley would be valuable if it were equally effective.

Material and methods

The experiment was conducted in accordance with the directives of the Experimental Unit 'Palo a Pique' of INIA Treinta y Tres in Uruguay with regard to the use of animals for experimentation.

Experimental treatments

The experiment was conducted at the Experimental Unit 'Palo a Pique' of INIA Treinta y Tres, Uruguay (35°S), in August 2003 using adult Corriedale ewes, 35 bearing single foetuses (44.3 ± 0.52 kg, body condition score 1.48 ± 0.03; scale 0 to 5 of Russell *et al.* (1969) measured at 128 days after mating) and 25 bearing twin foetuses (50.8 ± 0.91 kg, body condition score 1.52 ± 0.02).

†E-mail: jmilton@animals.uwa.edu.au

The experimental ewes were selected from a flock of 260 that had been synchronised using a single injection of 160 µg Delprostenate (Glandinex™ Universal Lab., Uruguay). They were mixed with Corriedale rams (1 per 12.5 ewes) during the second heat after synchronisation and scanned for pregnancy with single or twin fetuses 90 days later. The ewes grazed *ad libitum* native pastures (71 g crude protein (CP) and 6.6 MJ metabolisable energy (ME) per kg dry matter (DM)) from mating until they were housed indoors in individual pens on day 128 after mating (day -21 before lambing was expected to commence; day 0 = lambing). The ewes were fed 100% of their ME requirements according to Ministry of Agriculture, Fisheries and Food (MAFF, 1975) to maintain them in moderate body condition (about condition score 1.5). The twin-bearing ewes were each offered a basal diet of 1.4 kg DM of lucerne hay (153 g CP and 9.0 MJ ME per kg DM) per day. The single-bearing ewes were each offered a basal diet of 1.0 kg DM of lucerne hay per day. All ewes had access to a mineral mix (Cobalfosal Ovino Total® Barraca Deambrossi, Uruguay) and water *ad libitum*. Fourteen days before the expected time of lambing the ewes were allocated to three treatments and fed either the basal diet alone (control, 12 single- and 8 twin-bearers), the basal diet plus a supplement of whole barley (88.4% DM, 107 g CP and 13.5 MJ ME per kg DM; 'whole barley', 11 single- and 8 twin-bearers) or the basal diet plus a supplement of cracked maize (86.6% DM, 80 g CP and 13.3 MJ ME per kg DM; 'cracked maize', 12 single- and 9 twin-bearers). Both supplements were introduced to the ewes progressively at the rate of 0.2, 0.2, 0.3, 0.3, 0.4, 0.4, 0.5, 0.5 kg per head per day from day 134 to 141 of gestation (day -15 to day -8 before birth) and thereafter the ewes were offered 0.6 kg per head per day until they lambed (day 0). The lucerne hay was offered each morning at 0800 h and the whole barley or cracked maize was offered to the supplemented ewes 30 min later. Any lucerne hay or grain that was not eaten was weighed and recorded before the ewes were fed the next day.

Live weight and body condition

The body weight and condition score of ewes were recorded on days 65 and 130 of gestation.

Udder measures, colostrum production, viscosity and composition

The volume of the udder for each ewe was calculated just before and immediately after lambing by measuring the lateral and longitudinal semi-circumference of the udder with a flexible tape measure. Udder volume was derived considering the udder as a hemisphere and using the formula of Mellor and Murray (1985) modified by Bencini and Purvis (1990). The full volume of the udder at lambing was calculated from its linear dimensions and the empty volume was calculated by subtracting from the full volume, the amount of colostrum that could be expressed from the udder immediately after lambing.

The ewes were observed for 24 h/day and, immediately after lambing, each ewe received an intramuscular injection of 5 IU of oxytocin (Hipofamina™ Lab. Dispert, Uruguay). One teat was then completely milked-out by hand and covered with tape to prevent the lamb or lambs from sucking. The other teat was left uncovered for the lamb or lambs to suck. The colostrum expressed from the teat was weighed and classified on appearance and consistency according to the 0 to 7 scale of McCance and Alexander (1959): where 0 is no expressible secretion; 1 is a clear, serous straw-coloured liquid grading to 7 which is an opaque, white liquid similar to normal milk. A 20-ml sample of the colostrum had a preservative added (50 µl 10% potassium dichromate) and was stored at 4°C until analysed for components. The same teat from each ewe was milked following an intramuscular injection of 5 IU of oxytocin at 1, 3, 6 and 10 h after lambing. The resulting colostrum was weighed each time and a 20-ml sample was treated and stored in the same way as the initial sample until analysed. The colostrum expressed from the teat at lambing represented that which had accumulated prior to lambing and the colostrum expressed at each subsequent milking represented the quantity secreted since the previous milking. The samples of colostrum were analysed for fat, lactose, protein and total solids using a Milkoscan (133 Foss Electric Denmark).

Birth weight and identification of lambs

The lambs were weighed and identified soon after birth and each lamb was allowed to suck from the uncovered teat. When colostrum production was judged to be insufficient for the lamb's requirement (Mellor and Murray, 1986) the lamb(s) were given a supplement of colostrum that had been warmed to about 38°C.

Statistical analysis

The effects of supplementation on the weight and characteristics of the colostrum secreted, udder volume and lamb birth weight were analysed by least-squares analysis of variance using the SAS ProcGLM procedure (Statistical Analysis Systems Institute (SAS), 2001).

The model used was: $Y_{ij} = \mu + \tau_{ij} + \varepsilon_{ij}$ where: Y_{ij} is the random variable ($i =$ treatment 1...6; $j =$ repetitions); μ is the general mean for each treatment; $\tau_{ij} = A_i + B_j + (AB)_{ij}$, being A_i the effect of birth type (1,2) and B_j the effect of the supplement (1,2,3) and $(AB)_{ij}$ the interaction between birth type and supplement; ε_{ij} is the experimental error.

The viscosity of the colostrum was analysed as discrete data using the SAS GenMod procedure. Regression analysis was used to establish correlations between colostrum production and viscosity. Effects were deemed significant when the level of probability was 5% or less. The interaction between treatments and birth type was presented only if it was significant.

Results

Production of colostrum

At lambing, ewes supplemented with whole barley or cracked maize had accumulated more than double the quantity of colostrum accumulated by unsupplemented ewes and the twin-bearing ewes produced more than the single-bearing ewes ($P < 0.05$, Table 1). The colostrum produced by twin-bearing ewes supplemented with barley or maize did not differ but, single-bearing ewes supplemented with maize produced 50% more than the ewes supplemented with barley ($P < 0.05$). Subsequent production of colostrum remained greater in supplemented than in unsupplemented ewes for up to 3 h after lambing although, the production of colostrum did not differ between supplemented groups. Colostrum secretion was also higher in twin- than in single-bearing ewes up to 1 h after lambing, but there was no difference by 3 h after lambing ($P < 0.05$). The total quantity of colostrum produced (that accumulated at lambing plus subsequent production up to 10 h after lambing) was higher in supplemented than in unsupplemented ewes and was also higher in twin- compared with single-bearing ewes ($P < 0.05$). In single-bearing ewes, supplementation with maize increased colostrum production more than supplementation with barley but in twin-bearing ewes, barley produced more colostrum than maize ($P < 0.05$).

Udder development

At lambing, the udder volume of the supplemented ewes was only 16% greater ($P = 0.10$) than that of the unsupplemented ewes, but the full and empty udders of the twin-bearing ewes were much larger ($P < 0.001$) than those of the single-bearing ewes (Table 2).

Viscosity of the colostrum

Unsupplemented ewes had colostrum that was more viscous than that of ewes supplemented with either barley or maize (Table 3 and Annex, $P < 0.001$) irrespective of whether they bore single or twin lambs. The viscosity of

the colostrum decreased for all groups from parturition up to 10 h after birth. Immediately after parturition, unsupplemented ewes continued producing colostrum that was more viscous ($P > 0.01$) than that of supplemented ewes but by 10 h after birth there was no difference. The viscosity of the colostrum was negatively correlated with both the amount of colostrum accumulated at parturition ($n = 60$, $r = 0.54$, $P < 0.05$) and the yield of colostrum from lambing up to 1 h after birth ($n = 58$, $r = 0.36$, $P < 0.05$) or 3 h after birth ($n = 57$, $r = 0.37$, $P < 0.05$). The viscosity of the colostrum was not significantly correlated with the yield of colostrum 6 h after birth.

Components of colostrum

Supplementation also affected the main colostrum components at lambing ($P > 0.05$, Table 4). The total weight of solids in the colostrum secreted up to 10 h after birth were similar for the supplemented ewes (300 ± 26.5 g for those fed barley and 336 ± 25.2 g for those fed maize) and higher than unsupplemented ewes (209 ± 26 g) ($P < 0.01$) even though the percentage of solids were significantly lower in the supplemented groups. Twin-bearing ewes produced more total solids than single-bearing ewes (326 ± 22.8 g v. 237 ± 19.3 g, $P < 0.01$). At birth, the percentage of fat did not differ between treatments but the percentage of protein was lower and the percentage of lactose was higher in supplemented than in unsupplemented ewes ($P < 0.05$). There was no difference in the protein content of the colostrum produced by ewes supplemented with either barley or maize but the lactose content of the colostrum produced by single-bearing ewes supplemented with barley was lower than that produced by single-bearing ewes supplemented with maize ($P < 0.001$). From 1 h after birth the percentage of fat was similar regardless of supplementation. On the other hand, protein percentage continued to be lower in supplemented ewes up to 6 h after lambing ($P < 0.01$). In contrast, the percentage of lactose remained higher ($P < 0.01$) in supplemented than in unsupplemented ewes up to 3 h after

Table 1 Mean (\pm s.e.) colostrum (g) accumulated at birth and secreted 0 to 1, 1 to 3, 3 to 6 and 6 to 10 h after lambing and total production of colostrum in single- and twin-bearing Corriedale ewes supplemented with barley, maize or unsupplemented in the week before lambing

Treatments	Birth type						Significance [†]	
	Single-bearing ewes			Twin-bearing ewes				
	Control	Whole barley	Cracked maize	Control	Whole barley	Cracked maize	Birth type	Supplement
Colostrum								
At birth	190 (44.3)	360 (81.2)	541 (69.3)	292 (116)	648 (95.3)	623 (87.6)	*	***
Birth to 1 h	84.5 (18.1)	111 (12.1)	134 (22.8)	133 (25.8)	205 (45.3)	162 (27.1)	***	*
1–3 h after birth	77.8 (10.8)	107 (15.9)	117 (12.3)	90.5 (19.7)	124 (22.9)	140 (16.8)	NS	*
3–6 h after birth	87.3 (10.8)	99.6 (14.1)	115 (14.4)	89.5 (15.9)	130 (29.9)	112 (14.0)	NS	NS
6–10 h after birth	140 (17.5)	165 (19.0)	168 (15.4)	141 (22.9)	159 (21.7)	147 (38.9)	NS	NS
Total produced	580.3 (8.1)	838 (102.5)	1126 (13.3)	746 (120.2)	1245 (52.0)	1185 (13.3)	*	**

[†] Significant differences for values within rows: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$. NS, values within rows are not different.

Table 2 Mean (\pm s.e.) udder volume (ml) at lambing in single- and twin-bearing Corriedale ewes supplemented with barley, maize or unsupplemented in the week before lambing

Treatments	Birth type						Significance [†]	
	Single-bearing ewes			Twin-bearing ewes			Birth type	Supplement
	Control	Whole barley	Cracked maize	Control	Whole barley	Cracked maize		
Udder volume								
Full	1787 (258)	1894 (269)	2220 (258)	2359 (316)	2962 (316)	3592 (298)	***	*
Empty	1596 (221)	1534 (231)	1678 (221)	2067 (271)	2314 (271)	2969 (256)	***	NS

[†] Significant differences for values within rows: * $P \leq 0.05$, *** $P \leq 0.001$. NS, values within rows are not different.

Table 3 Mean viscosity scores (0 to 7) for colostrum accumulated at birth and secreted 0 to 1, 1 to 3, 3 to 6 and 6 to 10 h after lambing in single- and twin-bearing Corriedale ewes supplemented with barley, maize or unsupplemented in the week before lambing

Treatments	Birth type					
	Single-bearing ewes			Twin-bearing ewes		
	Control	Whole barley	Cracked maize	Control	Whole barley	Cracked maize
Viscosity scores						
At birth	4.2	6.0	6.3	4.3	6.3	6.1
Birth to 1 h	4.3	6.3	6.6	4.0	6.7	6.3
1–3 h after birth	5.7	6.8	7.0	5.5	7.0	6.8
3–6 h after birth	6.1	6.9	7.0	6.1	7.0	7.0
6–10 h after birth	6.8	7.0	7.0	6.8	7.0	7.0

birth. There was no difference between supplemented treatments for any of the components of colostrum.

At parturition the colostrum of single- and twin-bearing ewes did not differ in composition. The total weight of solids was similar for single and twin-bearing ewes for the first three milkings was higher for the twin-bearing ewes at 6 h ($P < 0.05$) becoming similar again at 10 h (Table 4). After lambing, the percentage of fat and protein remained similar in both groups but the single-bearing ewes had

a higher percentage of lactose in their milk than twin-bearing ewes at 1 and 6 h after birth ($P < 0.05$).

Birth weight of lambs

Supplementation of the ewes prior to lambing did not affect the birth weight of their lambs but twin lambs were lighter than single lambs ($P < 0.001$; Table 5).

Consumption of roughages and supplements

Single-bearing ewes consumed all of the roughage diet offered, as did control twin-bearing ewes. On the other hand, twin-bearing ewes supplemented with maize or barley left 7% each. Single-bearing ewes supplemented with maize left 5% of the grain diet whereas the ewes supplemented with barley left 9%. Supplemented twin-bearing ewes ate their entire supplement.

Table 3 Annex Comparison between treatments

Factors	Comparisons		Significance [†]
	Single	Twin	
Birth type			NS
Supplement	Barley	Control	***
	Barley	Maize	NS
	Control	Maize	***
Time of milking	At birth	Birth to 1 h	NS
	At birth	1–3 h after birth	*
	At birth	3–6 h after birth	**
	At birth	6–10 h after birth	***
	Birth to 1 h	1–3 h after birth	NS
	Birth to 1 h	3–6 h after birth	*
	Birth to 1 h	6–10 h after birth	**
	1–3 h after birth	3–6 h after birth	NS
	1–3 h after birth	6–10 h after birth	NS
	3–6 h after birth	6–10 h after birth	NS

[†] Significant differences for values within rows: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$. NS, values within rows are not different.

Discussion

Supplementing ewes before lambing either with whole barley or cracked maize increased the quantity of colostrum accumulated at birth and its subsequent secretion during the following hours. Supplemented ewes produced between 1.9 to 2.8 times more colostrum at birth than unsupplemented ewes despite the unsupplemented ewes being fed to meet their estimated ME requirements (MAFF, 1975). The increase in production of colostrum was of the same order as that found by Banchero *et al.* (2004a and b) using maize alone. This strengthens our hypothesis that

Table 4 Mean values (\pm s.e.) for the components of colostrum at different milking times in single- and twin-bearing Corriedale ewes supplemented with barley, maize or unsupplemented in the week before lambing

Treatments	Birth type						Significance [†]	
	Single-bearing ewes			Twin-bearing ewes			Birth type	Supplement
	Control	Whole barley	Cracked maize	Control	Whole barley	Cracked maize		
Fat (%)								
At birth	14.0 (1.45)	11.7 (1.45)	11.7 (1.32)	14.2 (1.73)	13.3 (1.62)	13.7 (1.53)	NS	NS
1 h after birth	17.0 (1.70)	19.1 (1.70)	18.9 (1.60)	16.3 (2.00)	23.2 (2.10)	22.7 (1.80)	NS	*
3 h after birth	18.0 (2.20)	17.2 (2.20)	19.2 (2.10)	15.1 (2.60)	17.7 (2.80)	20.3 (2.40)	NS	NS
6 h after birth	12.6 (1.90)	10.4 (2.00)	11.9 (1.90)	11.4 (2.30)	18.7 (2.30)	17.9 (2.20)	*	NS
10 h after birth	12.7 (1.50)	12.0 (1.60)	12.1 (1.50)	11.0 (1.90)	13.9 (2.00)	15.5 (1.80)	NS	NS
Protein (%)								
At birth	20.2 (1.17)	17.5 (1.17)	15.3 (1.07)	22.9 (1.40)	14.8 (1.31)	15.8 (1.24)	NS	***
1 h after birth	17.9 (1.30)	14.3 (1.20)	12.7 (1.20)	20.1 (1.40)	12.2 (1.50)	13.1 (1.40)	NS	***
3 h after birth	12.5 (1.10)	10.2 (1.10)	8.80 (1.00)	15.4 (1.30)	8.60 (1.40)	10.4 (1.20)	NS	***
6 h after birth	11.2 (1.20)	7.60 (1.20)	6.60 (1.20)	10.6 (1.40)	6.20 (1.40)	8.00 (1.30)	NS	**
10 h after birth	7.70 (1.00)	5.60(1.10)	5.00 (1.00)	8.40 (1.30)	5.10 (1.40)	8.20 (1.20)	NS	NS
Lactose (%)								
At birth	1.6 (0.20)	2.5 (0.20)	3.1 (0.20)	1.4 (0.30)	2.8 (0.30)	2.4 (0.20)	NS	***
1 h after birth	1.5 (0.20)	2.1 (0.20)	2.5 (0.20)	1.2 (0.20)	2.0 (0.20)	2.1 (0.20)	*	***
3 h after birth	2.1 (0.30)	3.0 (0.30)	3.0 (0.20)	2.2 (0.30)	3.2 (0.30)	3.0 (0.30)	NS	**
6 h after birth	3.2 (0.30)	4.3 (0.30)	4.0 (0.30)	3.4 (0.40)	3.3 (0.40)	4.3 (0.30)	*	NS
10 h after birth	3.8 (0.20)	4.4 (0.20)	4.4 (0.20)	3.8 (0.30)	4.0 (0.30)	4.4 (0.20)	NS	NS
Total solids (%)								
At birth	35.8 (1.70)	31.7(1.70)	30.1 (1.60)	38.5 (2.10)	30.9 (1.90)	31.9 (1.80)	NS	*
1 h after birth	36.4 (1.30)	35.4 (1.20)	34.2 (1.20)	37.6 (1.50)	37.4 (1.60)	37.6 (1.40)	NS	NS
3 h after birth	32.6 (2.00)	30.5 (2.00)	31.0 (1.90)	32.6 (2.40)	29.6 (2.60)	33.0 (2.20)	NS	NS
6 h after birth	27.0 (1.90)	22.3 (1.90)	22.5 (1.90)	25.4 (2.30)	28.2 (2.30)	28.8 (2.20)	*	NS
10 h after birth	24.1 (1.60)	21.9 (1.60)	21.5 (1.60)	23.2 (1.90)	22.9 (2.10)	27.5 (1.80)	NS	NS

[†] Significant differences for values within rows: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$. NS, values within rows are not different.

Table 5 Mean (\pm s.e.) birth weight of lambs born to single- and twin-bearing Corriedale ewes supplemented with barley, maize or unsupplemented in the week before lambing

Treatments	Birth type						Significance [†]	
	Single-bearing ewes			Twin-bearing ewes			Birth type	Supplement
	Control	Whole barley	Cracked maize	Control	Whole barley	Cracked maize		
Lamb birth weight	4.5 (0.08)	4.6 (0.19)	4.5 (0.18)	3.5 (0.27)	4.1 (0.15)	3.8 (0.23)	***	NS

[†] Significant differences for values within the row: *** $P \leq 0.001$. NS, values within the row are not different.

the starch supplied by both grains is the key to stimulating the production of colostrum. Large amounts of starch may pass into the small intestine and can account for a large proportion of the starch digested in the whole digestive tract (Nocek and Tamminga, 1991). This may increase the entry rate of glucose (Knowlton *et al.*, 1998; Landau *et al.*, 1999) and, subsequently, the mammary uptake of glucose and the synthesis of lactose by the mammary gland (Linzell, 1974).

Barley is fermented to a greater extent than maize in the rumen (Nocek and Tamminga, 1991) and therefore would be expected to provide less starch for digestion to glucose in the small intestine. Indeed, in the single-bearing ewes, barley was less effective than maize to increase the

amount of colostrum produced but it was just as effective in twin-bearing ewes. In twin-bearing ewes Weston (1988) showed that the rate of passage of digesta from the rumen is higher than in single-bearing ewes due to the compression by the twin foetuses. So it is likely that more unfermented barley passed from the rumen to the intestine in our twin-bearing ewes resulting in a similar glucose entry rate from both supplements.

The greater response in the yield of colostrum of twin-bearing ewes over single-bearing ewes after supplementation is consistent with previous work (Hall *et al.*, 1992; Banchemo *et al.*, 2004a). Twin-bearing ewes developed bigger udders than single-bearing ewes (Mellor, 1988) with the capacity to synthesise more colostrum.

Supplemented ewes not only produced more colostrum than unsupplemented ewes but also the colostrum they produced was more liquid and this, in turn, was associated with higher levels of lactose. Lactose is osmotically active (Leong *et al.*, 1990) so draws water from the blood and, in doing so, decreases the viscosity of the colostrum. The ewes with colostrum of highest viscosity and at the same time the lowest concentration of lactose were the unsupplemented ewes regardless of whether they bore twin or single lambs.

From a practical standpoint, short-term, strategic supplementation did not significantly increase the weight of the lambs and therefore it did not increase the chances of difficult births. On the contrary, it ensured that the volume of colostrum available at birth met, or exceeded the estimated requirements of 50 ml/kg of lamb birth weight (Robinson *et al.*, 2002). In addition, the fact that the colostrum was less viscous means that it should be more readily taken in by the lamb when it first attempts to suck and this, in turn, should aid in establishing a stronger maternal bond (Nowak and Poindron, 2006). This suggests that supplementation is a worthwhile strategy for increasing the chances of survival of newborn lambs. The results presented here show that the form of supplementation, maize or barley, can be chosen on economic grounds as the lambs that are usually most at risk, the twins, receive equivalent benefit when their mothers receive either supplement.

Acknowledgements

We thank Gabriel and Fernando García, Damián Gonzalez and Jonny Nogueira for their technical help. We thank Dr Gabriel Ciappesoni for assistance with the statistical analysis.

References

- Banchero GE, Quintans G, Martin GB, Lindsay DR and Milton JTB 2004a. Nutrition and colostrum production in sheep. 1. Metabolic and hormonal responses to a high-energy supplement in the final stages of pregnancy. *Reproduction Fertility and Development* 16, 633-643.
- Banchero GE, Quintans G, Martin GB, Milton JTB and Lindsay DR 2004b. Nutrition and colostrum production in sheep. 2. Metabolic and hormonal responses to different energy sources in the final stages of pregnancy. *Reproduction Fertility and Development* 16, 645-653.
- Banchero GE, Quintans G, Martin GB, Milton JTB and Lindsay DR 2003. Production of colostrum by Polwarth ewes grazing lucerne pasture: effect of birth type and body condition. *Proceedings, world congress of animal production; Porto Alegre, Brazil*, 224 (abstr.)
- Bencini R and Purvis IW 1990. The yield and composition of milk from Merino sheep. *Animal Production in Australia. Proceedings of the Australian Society of Animal Production* 18, 144-147.
- Forbes JM 1968. The physical relationships of the abdominal organs in the pregnant ewe. *Journal of Agricultural Science, Cambridge* 70, 171-177.
- Hall DG, Holst PJ and Shutt DA 1992. The effect of nutritional supplements in late pregnancy on ewe colostrum production, plasma progesterone and IGF-1 concentrations. *Australian Journal of Agricultural Research* 43, 325-337.
- Knowlton KF, Dawson TE, Glenn BP, Huntington GB and Erdman RA 1998. Glucose metabolism and milk yield of cows infused abomasally or ruminally with starch. *Journal of Dairy Science* 81, 3248-3258.
- Landau S, Nitsan Z, Zoref Z and Madar Z 1999. The influence of extruding corn grain on glucose metabolism in pregnant ewes. *Reproduction, Nutrition, Development* 39, 181-187.
- Leong WS, Navaratnam N, Stankiewicz MJ, Wallace AV, Ward S and Kuhn N 1990. Subcellular compartmentation in the synthesis of the milk sugars lactose and -2,3-sialyllactose. *Protoplasm* 159, 144-156.
- Linzell JL 1974. Mammary blood flow and methods of identifying and measuring precursors of milk. In *Lactation: a comprehensive treatise volume 1*, (ed. BL Larson and VR Smith), pp. 143-225. Academic Press, New York.
- McCance I and Alexander G 1959. The onset of lactation in the Merino ewe and its modification by nutritional factors. *Australian Journal of Agricultural Research* 10, 699-719.
- McNeill DM, Murphy PM and Lindsay DR 1998. Blood lactose v. milk lactose as a monitor of lactogenesis and colostrum production in Merino ewes. *Australian Journal of Agricultural Research* 49, 581-587.
- Ministry of Agriculture, Fisheries and Food 1975. Energy allowances and feeding systems for ruminants. Technical bulletin 33. MAFF, London.
- Mellor DJ 1988. Integration of perinatal events, pathophysiological changes and consequences for the newborn lamb. *British Veterinary Journal* 144, 552-569.
- Mellor DJ and Murray L 1985. Effects of maternal nutrition on udder development during late pregnancy and on colostrum production in Scottish Blackface ewes with twin lambs. *Research in Veterinary Science* 39, 230-234.
- Mellor DJ and Murray L 1986. Making the most of colostrum at lambing. *Veterinary Record* 118, 351-353.
- Murphy PM, McNeill DM, Fisher JS and Lindsay DR 1996. Strategic feeding of Merino ewes in late pregnancy to increase colostrum production. *Animal production in Australia. Proceedings of the Australian Society of Animal Production* 21, 227-230.
- Nocek JE and Tamminga S 1991. Site of digestion of starch in the gastrointestinal tract of dairy cows and its effect on milk yield and composition. *Journal of Dairy Science* 74, 3598-3629.
- Nowak R and Poindron P 2006. From birth to colostrum: early steps leading to lamb survival. *Reproduction Nutrition and Development* 46, 431-446.
- Robinson JJ, Rooke JA and McEvoy TG 2002. Nutrition for conception and pregnancy. In *Sheep nutrition* (ed. M Freer and H Dove), pp. 189-211, CABI Publishing in association with CSIRO Publishing, Canberra.
- Russell AJF, Doney JM and Gunn RG 1969. Subjective assessment of body fat in live sheep. *Journal of Agricultural Science, Cambridge* 72, 451-454.
- Statistical Analysis Systems Institute 2001. SAS version 8.02. SAS Institute, Cary, NC, USA.
- Weston RH 1988. Factors limiting the intake of feed by sheep. 11. The effect of pregnancy and early lactation on the digestion of a medium-quality roughage. *Australian Journal of Agricultural Research* 39, 659-669.