



ON SOME SEMI-PARAMETRIC ESTIMATES FOR EUROPEAN OPTION PRICES

CARLO MARINELLI *, *University College London*

Abstract

We show that an estimate by de la Peña, Ibragimov, and Jordan for $\mathbb{E}(X - c)^+$, with c a constant and X a random variable of which the mean, the variance, and $\mathbb{P}(X \leq c)$ are known, implies an estimate by Scarf on the infimum of $\mathbb{E}(X \wedge c)$ over the set of positive random variables X with fixed mean and variance. This also shows, as a consequence, that the former estimate implies an estimate by Lo on European option prices.

Keywords: Probabilistic inequalities; bounds for option prices

2020 Mathematics Subject Classification: Primary 60E15

Secondary 91G20; 91G80

1. Introduction

A remarkable result by Scarf [11] provides an explicit solution to the problem of minimizing $\mathbb{E}(X \wedge c)$, with c a positive constant, over the set of all positive random variables X with given mean and variance (see Theorem 4.1 below). The infimum is shown to have two different expressions depending on whether the parameter c is above or below a certain threshold and to be attained by a random variable taking only two values. About thirty years after the publication of [11], Lo [7] noticed that Scarf's result immediately implies an upper bound for $\mathbb{E}(X - c)^+$, with X and c as before. This has an obvious financial interpretation as an upper bound for the price at time zero of a European call option with strike c on an asset with value at maturity equal to X (in discounted terms, assuming that expectation is meant with respect to a pricing measure). More recently, de la Peña, Ibragimov, and Jordan [5] obtained, among other things, a sharp upper bound for $\mathbb{E}(X - c)^+$ over the set of random variables X for which mean and variance as well as the probability $\mathbb{P}(X \leq c)$ are known (see Theorem 3.1 below).

Our goal is to prove that the estimate by de la Peña, Ibragimov, and Jordan is stronger than Scarf's in the sense that the former implies the latter. This may appear somewhat counterintuitive, as the former estimate requires the extra input $\mathbb{P}(X \leq c)$, while the latter has an extra positivity constraint.

The proof by Scarf, while relatively elementary, is quite ingenious. A different proof, also covering substantial generalizations, has been obtained in [2] using duality methods in semi-definite optimization. The arguments used in [5] are instead based on classical probabilistic

Received 23 June 2023; accepted 31 October 2023.

* Postal address: Department of Mathematics, University College London, Gower Street, London WC1E 6BT, UK.
Email: c.marinelli@ucl.ac.uk

© The Author(s), 2024. Published by Cambridge University Press on behalf of Applied Probability Trust. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

inequalities and a ‘toy’ version of decoupling. The proof of Scarf’s estimate given here is entirely elementary and self-contained. Starting from an alternative proof of the relevant estimates in [5], another proof of Scarf’s result is obtained that is certainly not as deft as the original, but that would hopefully seem more natural to anyone who, like the author, would hardly ever come up with the ingenious idea used in [11]. Our proof is based, roughly speaking, on a representation of the set of random variables with given mean and variance as a union of subsets of equivalent random variables, where two random variables X_1 and X_2 are equivalent if $\mathbb{P}(X_1 \leq c) = \mathbb{P}(X_2 \leq c)$. This allows to establish a link between the two inequalities and to reduce the problem of proving a version of Scarf’s result without the positivity constraint to the minimization of a function of one real variable. Finally, the positivity constraint is taken into account, thus establishing the full version of Scarf’s result. Moreover, the fact that optimizers exist and are given by two-point distributed random variables appears in a natural way and plays an important role in the proof.

The result proved in this article may also clarify, or at least complement, several qualitative remarks made in [5] about the relation between the two abovementioned inequalities. For instance, the authors note that their inequality is simpler than Lo’s in the sense that the right-hand side does not depend on the value of c . Here we show that this is only due to the positivity constraint in [11], with very explicit calculations showing how the threshold value for c arises. Moreover, the sharpness of their inequality is proved in a much more natural way, i.e. showing that a two-point distributed random variable always attains the bound.

Apart from the application to bounds for option prices, estimates of (functions of) $X \wedge c$, sometimes called the Winsorization of X , are important in several areas of applied probability and statistics (see, e.g., [10]). For results in this direction, as well as for an informative discussion with references to the literature, we refer to [9].

2. Preliminaries

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, on which all random elements will be defined. We shall write, for simplicity, L^2 to denote $L^2(\Omega, \mathcal{F}, \mathbb{P})$, and $\|\cdot\|_2$ for its norm. For any $m \in \mathbb{R}$ and $\sigma \in \mathbb{R}_+$, the sphere of L^2 of radius σ centered in m will be denoted by $\mathcal{X}_{m,\sigma}$, and just by \mathcal{X} if $m = 0$ and $\sigma = 1$. In other words, $\mathcal{X}_{m,\sigma}$ stands for the set of random variables X such that $\mathbb{E}X = m$ and $\text{Var}(X) = \mathbb{E}(X - m)^2 = \sigma^2$. It is clear that $\mathcal{X}_{m,\sigma} = m + \sigma \mathcal{X}$. The intersection of $\mathcal{X}_{m,\sigma}$ with the cone of random variables bounded below by $\alpha \in \mathbb{R}$ will be denoted by $\mathcal{X}_{m,\sigma}^\alpha$. It is easily verified that, for any $m \in \mathbb{R}$ and $\sigma \in \mathbb{R}_+$,

$$m + \sigma \mathcal{X}^\alpha = \mathcal{X}_{m,\sigma}^{m+\sigma\alpha}. \quad (2.1)$$

Recall that a random variable is said to be two-point distributed if it takes only two (distinct) values. The set of two-point distributed random variables belonging to \mathcal{X} can be parametrized by the open interval $]0, 1[$: let X take the values $x, y \in \mathbb{R}$, $x < y$, and set $p := \mathbb{P}(X = x)$, $1 - p = \mathbb{P}(X = y)$. Then $X \in \mathcal{X}$ if and only if $px + (1 - p)y = 0$ and $px^2 + (1 - p)y^2 = 1$, which implies

$$x = -\left(\frac{1-p}{p}\right)^{1/2}, \quad y = \left(\frac{p}{1-p}\right)^{1/2}. \quad (2.2)$$

Note that $p = 0$ and $p = 1$ are not allowed, and hence $p \in]0, 1[$. This is also obvious a priori, as there is no degenerate random variable with mean zero and variance one. The following simple observations, the proofs of which are immediate consequences of (2.2) and elementary algebra, will be useful.

Lemma 2.1. Let $X \in \mathcal{X}$ be the $\{x, y\}$ -valued random variable identified by the parameter $p \in]0, 1[$, and $c \in \mathbb{R}$.

- (i) If $c \geq 0$, then $x < c \leq y$ if and only if $p \geq c^2/(1 + c^2)$.
- (ii) If $c < 0$, then $x \leq c < y$ if and only if $p \leq 1/(1 + c^2)$.

We shall also need the following elementary lattice identities.

Lemma 2.2. Let $a, b, c \in \mathbb{R}$. The following hold:

- (i) $a + (b \wedge c) = (a + b) \wedge (a + c)$;
- (ii) if $a \geq 0$, then $a(b \wedge c) = (ab) \wedge (ac)$;
- (iii) $(a - b)^+ = a - (a \wedge b)$.

Proof. The identities in (i) and (ii) are clear. The identity in (iii) can be verified ‘case by case’, but it can also be deduced from the identity $(a - b)^+ = (a - b) + (a - b)^-$, where, using (i),

$$(a - b)^- = -((a - b) \wedge 0) = b - (a \wedge b),$$

from which the claim follows immediately. □

3. de la Peña–Ibragimov–Jordan bound

The following sharp estimates are proved in [5].

Theorem 3.1. (de la Peña, Ibragimov, and Jordan) Let $X \in \mathcal{X}_{m,\sigma}$ and $c \in \mathbb{R}$. Setting $p_0 := \mathbb{P}(X > c)$, we have

$$(m - c)p_0 \leq \mathbb{E}(X - c)^+ \leq (m - c)p_0 + \sigma(p_0 - p_0^2)^{1/2}. \tag{3.1}$$

The proof of (3.1) in [5] is very elegant, the main idea being the introduction of an independent copy of the random variable X . Here we give an alternative, entirely elementary, proof.

Proof. Let us start with the lower bound. We can assume, without loss of generality, that $m \geq c$, otherwise there is nothing to prove. Since

$$\mathbb{E}(X - c)^+ = \mathbb{E}(X - c)\mathbf{1}_{\{X > c\}} = \mathbb{E}X\mathbf{1}_{\{X > c\}} - c\mathbb{P}(X > c),$$

it suffices to show that $\mathbb{E}X\mathbf{1}_{\{X > c\}} \geq m\mathbb{P}(X > c)$. To this purpose, note that

$$\mathbb{E}X\mathbf{1}_{\{X > c\}} = \mathbb{E}X - \mathbb{E}X\mathbf{1}_{\{X \leq c\}},$$

where, thanks to the assumption $m \geq c$,

$$\mathbb{E}X\mathbf{1}_{\{X \leq c\}} \leq \mathbb{E}c\mathbf{1}_{\{X \leq c\}} = c\mathbb{P}(X \leq c) \leq m\mathbb{P}(X \leq c),$$

and hence $\mathbb{E}X\mathbf{1}_{\{X > c\}} \geq m - m\mathbb{P}(X \leq c) = m\mathbb{P}(X > c)$.

To prove the upper bound, note that

$$\mathbb{E}(X - c)\mathbf{1}_{\{X > c\}} - (m - c)p_0 = \mathbb{E}(X\mathbf{1}_{\{X > c\}} - m\mathbf{1}_{\{X > c\}}),$$

hence, adding and subtracting $\mathbb{E}Xp_0 = mp_0$ on the right-hand side, the Cauchy–Schwarz inequality yields

$$\mathbb{E}(X - c)^+ - (m - c)p_0 = \mathbb{E}(X - m)(\mathbf{1}_{\{X > c\}} - p_0) \leq \sigma \|\mathbf{1}_{\{X > c\}} - p_0\|_2 = \sigma(p_0 - p_0^2)^{1/2},$$

thus completing the proof. \square

Remark 3.1. Even if $m < c$, the inequality $\mathbb{E}X\mathbf{1}_{\{X > c\}} \geq m\mathbb{P}(X > c)$ is still true. In fact, $m\mathbb{P}(X > c) \leq c\mathbb{P}(X > c) = \mathbb{E}c\mathbf{1}_{\{X > c\}} \leq \mathbb{E}X\mathbf{1}_{\{X > c\}}$.

Theorem 3.1 implies useful one-sided Chebyshev-like bounds.

Corollary 3.1. Let $X \in \mathcal{X}$ and $c \in \mathbb{R}$. The following hold:

- (i) if $c \geq 0$, then $\mathbb{P}(X \leq c) \geq \mathbb{P}(X < c) \geq c^2/(1 + c^2)$;
- (ii). if $c < 0$, then $\mathbb{P}(X \leq c) \leq 1/(1 + c^2)$.

Proof. The proof of Theorem 3.1 remains valid with $p_1 := \mathbb{P}(X \geq c)$ in place of p_0 ; hence, as the right-hand side of (3.1) must be positive, $\sqrt{p_1(1 - p_1)} \geq cp_1$. If $c \geq 0$, squaring both sides yields a linear inequality that is satisfied if and only if $p_1 \leq 1/(1 + c^2)$. This proves (i).

For (ii), if $c < 0$, $\mathbb{P}(X \leq c) = \mathbb{P}(-X \geq -c) = 1 - \mathbb{P}(-X < -c)$. Since $-X \in \mathcal{X}$ and $-c > 0$, (i) implies that $\mathbb{P}(-X < -c) \geq c^2/(1 + c^2)$, and hence

$$\mathbb{P}(X \leq c) \leq 1 - \frac{c^2}{1 + c^2} = \frac{1}{1 + c^2}. \quad \square$$

Remark 3.2. Let $X \in \mathcal{X}$ and $c \in \mathbb{R}_+$. By reasoning entirely analogous to the proof of Corollary 3.1(ii), both $\mathbb{P}(X > c)$ and $\mathbb{P}(X < -c)$ are bounded above by $1/(1 + c^2)$; hence $\mathbb{P}(|X| > c) \leq 2/(1 + c^2)$, which is sharper than Chebyshev's inequality $\mathbb{P}(|X| > c) \leq 1/c^2$ if $c < 1$.

4. Scarf–Lo bound

The following estimate is obtained in [11].

Theorem 4.1. (Scarf) Let c, m, σ be strictly positive real numbers. The infimum of the function $X \mapsto \mathbb{E}(X \wedge c)$ on the set $\mathcal{X}_{m,\sigma}^0$ is attained, i.e. it is a minimum, and is given by

$$\min_{X \in \mathcal{X}_{m,\sigma}^0} \mathbb{E}(X \wedge c) = \begin{cases} \frac{m^2}{m^2 + \sigma^2} c & \text{if } c \leq \frac{m^2 + \sigma^2}{2m}, \\ \frac{c + m}{2} - \frac{1}{2}((c - m)^2 + \sigma^2)^{1/2} & \text{if } c \geq \frac{m^2 + \sigma^2}{2m}. \end{cases}$$

Note that the constraint $X \geq 0$ is dictated by the structure of the practical inventory problem considered by Scarf. It is not needed, however, to avoid the infimum being minus infinity. In fact,

$$|\mathbb{E}(X \wedge c)| \leq \mathbb{E}|X \wedge c| \leq \mathbb{E}|X| + |c| \leq (\mathbb{E}X^2)^{1/2} + |c|,$$

where $(\mathbb{E}X^2)^{1/2} = \|X\|_2 = \|X - m + m\|_2 \leq \sigma + |m|$, which implies

$$\inf_{X \in \mathcal{X}_{m,\sigma}^0} \mathbb{E}(X \wedge c) \geq -(\sigma + |m| + |c|).$$

As observed in [7], Lemma 2.2(iii) immediately yields the following result.

Corollary 4.1. (Lo) *Let c, m, σ be strictly positive real numbers. The supremum of the function $X \mapsto \mathbb{E}(X - c)^+$ on the set $\mathcal{X}_{m,\sigma}^0$ is attained, i.e. it is a maximum, and is given by*

$$\max_{X \in \mathcal{X}_{m,\sigma}^0} \mathbb{E}(X - c)^+ = \begin{cases} m - \frac{m^2}{m^2 + \sigma^2}c & \text{if } c \leq \frac{m^2 + \sigma^2}{2m}, \\ \frac{m - c}{2} + \frac{1}{2}((m - c)^2 + \sigma^2)^{1/2} & \text{if } c \geq \frac{m^2 + \sigma^2}{2m}. \end{cases}$$

The remainder of this section is dedicated to showing that Theorem 4.1, and hence also its corollary, are consequences of Theorem 3.1. We shall argue by a sequence of elementary lemmas and propositions. The first is a reduction step that, in spite of its simplicity, considerably reduces the burden of symbolic calculations. Throughout the section we assume that $c, m, \sigma \in \mathbb{R}$, with $\sigma > 0$. Further constraints (that do not imply any loss of generality) will be introduced as needed.

Lemma 4.1. *Let $\tilde{c} := (c - m)/\sigma$. Then*

$$\inf_{X \in \mathcal{X}_{m,\sigma}^0} \mathbb{E}(X \wedge c) = m + \sigma \inf_{X \in \mathcal{X}^{-m/\sigma}} \mathbb{E}(X \wedge \tilde{c}).$$

Proof. Since $\mathcal{X}_{m,\sigma}^0 = m + \sigma \mathcal{X}^{-m/\sigma}$ by (2.1), we have

$$\inf_{X \in \mathcal{X}_{m,\sigma}^0} \mathbb{E}(X \wedge c) = \inf_{Y \in \mathcal{X}^{-m/\sigma}} \mathbb{E}((m + \sigma Y) \wedge c),$$

where, by Lemma 2.2,

$$\mathbb{E}((m + \sigma Y) \wedge c) = m + \sigma \mathbb{E}\left(Y \wedge \frac{c - m}{\sigma}\right),$$

which immediately yields the conclusion. □

The lemma implies that it suffices to study the problem of minimizing the function $X \mapsto \mathbb{E}(X \wedge c)$ over the set \mathcal{X}^{-m} . We shall first study the minimization problem without the lower-boundedness constraint, i.e. on \mathcal{X} rather than on \mathcal{X}^{-m} . We shall need some more notation: the subset of $\mathcal{X}_{m,\sigma}$ such that $\mathbb{P}(X \leq c) = p$ is denoted by $\mathcal{X}_{m,\sigma}(p; c)$. Note that, in view of Corollary 3.1, these sets are nonempty only for certain combinations of the parameters p and c . Let $L_c :]0, 1[\mapsto \mathbb{R}$ be the function defined by $L_c : p \mapsto -(p - p^2)^{1/2} + c(1 - p)$.

Lemma 4.2. $\inf_{X \in \mathcal{X}(c;p)} \mathbb{E}(X \wedge c) \geq L_c(p)$.

Proof. Lemma 2.2 (iii) implies $\mathbb{E}(X - c)^+ = -\mathbb{E}(X \wedge c)$ for any X with mean zero; hence, by Theorem 3.1,

$$\inf_{X \in \mathcal{X}(c;p)} \mathbb{E}(X \wedge c) = - \sup_{X \in \mathcal{X}(c;p)} \mathbb{E}(X - c)^+ \geq -(p - p^2)^{1/2} + c(1 - p). \quad \square$$

We are now going to show that the infimum in Lemma 4.2 is achieved, and that the minimizer is a two-point distributed random variable. We shall only consider, without loss

of generality, those values of p such that $\mathcal{X}(p;c)$ is nonempty, that is, by Corollary 3.1, setting

$$\Pi_c := \begin{cases} \left] 0, \frac{1}{1+c^2} \right] & \text{if } c < 0, \\ \left[\frac{c^2}{1+c^2}, 1 \right[& \text{if } c \geq 0, \end{cases}$$

to $p \in \Pi_c$.

Lemma 4.3. *Let $p \in \Pi_c$, and $X_0 \in \mathcal{X}$ be the two-point distributed random variable with parameter p . Then $\mathbb{E}(X_0 \wedge c) = L_c(p)$, and hence*

$$\inf_{X \in \mathcal{X}(c;p)} \mathbb{E}(X \wedge c) = \min_{X \in \mathcal{X}(c;p)} \mathbb{E}(X \wedge c) = \mathbb{E}(X_0 \wedge c) = L_c(p).$$

Proof. Since $p \in \Pi_c$, Lemma 2.1 implies that the random variable X_0 , taking the values x and y as defined in (2.2), is such that $x \leq c \leq y$. In particular, $\mathbb{P}(X_0 \leq c) = p$, that is, $X_0 \in \mathcal{X}(p;c)$, and an elementary computation finally shows that $\mathbb{E}(X_0 \wedge c) = L_c(p)$. \square

The following result essentially shows that Theorem 3.1 implies Theorem 4.1 in the unconstrained case (i.e. without assuming that the minimizer should be bounded from below).

Proposition 4.1. $\inf_{X \in \mathcal{X}} \mathbb{E}(X \wedge c) = \inf_{p \in \Pi_c} L_c(p)$.

Proof. The decomposition $\mathcal{X} = \bigcup_{p \in \Pi_c} \mathcal{X}(p;c)$ implies (see, e.g., [3, p. III.11])

$$\inf_{X \in \mathcal{X}} \mathbb{E}(X \wedge c) = \inf_{p \in \Pi_c} \inf_{X \in \mathcal{X}(p;c)} \mathbb{E}(X \wedge c),$$

so that Lemma 4.3 implies the claim. \square

This clearly indicates that the next step should be to find the minimum of the function L_c .

Lemma 4.4. *The function L_c satisfies the following properties:*

- (i) *it is decreasing on the interval $\left] 0, \frac{1}{2} + \frac{1}{2} \frac{c}{(1+c^2)^{1/2}} \right]$;*
- (ii) *it is increasing on the interval $\left[\frac{1}{2} + \frac{1}{2} \frac{c}{(1+c^2)^{1/2}}, 1 \right[$;*
- (iii) *it admits a unique minimum point p_* defined by $p_* := \frac{1}{2} + \frac{1}{2} \frac{c}{(1+c^2)^{1/2}}$.*

Moreover, p_ belongs to Π_c and $L_c(p_*) = \frac{1}{2}c - \frac{1}{2}(1+c^2)^{1/2}$.*

Proof. The argument is elementary, so it will be sketched only (the details can be found in [8]). The function L_c is smooth and its derivative is the function

$$L'_c: p \mapsto -\frac{1}{2}(p(1-p))^{-1/2}(1-2p) - c;$$

hence, standard calculus yields the claims (i)–(iii). It only remains to show that $p_* \in \Pi_c$; if $c \geq 0$, this is equivalent to

$$p_* = \frac{1}{2} + \frac{1}{2} \frac{c}{(1+c^2)^{1/2}} \geq \frac{c^2}{1+c^2}.$$

Setting $x := c/(1 + c^2)^{1/2} \in [0, 1]$, this reduces to $1 + x \geq 2x^2$, which is satisfied if $x \in [0, 1]$. If $c < 0$, p_* belongs to Π_c if and only if

$$p_* = \frac{1}{2} + \frac{1}{2} \frac{c}{(1 + c^2)^{1/2}} = \frac{1}{2} - \frac{1}{2} \frac{|c|}{(1 + c^2)^{1/2}} \leq \frac{1}{1 + c^2},$$

that is, setting $\langle c \rangle := (1 + c^2)^{1/2}$ for convenience, if and only if $\langle c \rangle |c| \geq \langle c \rangle^2 - 2$. This inequality is easily seen to be satisfied for every $c \in [-1, 0]$. If $c \leq -1$, the inequality is equivalent to $\langle c \rangle^2 c^2 \geq (c^2 - 1)^2$, which simplifies to $3c^2 \geq 1$, i.e. it is verified for every $|c| \geq 1/\sqrt{3} \vee 1$, that is for every $c \leq -1$. Finally, the expression for $L_c(p_*)$ follows by elementary algebra. \square

We have thus solved the problem of minimizing the function $X \mapsto \mathbb{E}(X \wedge c)$ on \mathcal{X} .

Proposition 4.2. *Let p_* be defined as in Lemma 4.4, and*

$$x_* := -\left(\frac{1 - p_*}{p_*}\right)^{1/2}, \quad y_* := \left(\frac{p_*}{1 - p_*}\right)^{1/2}.$$

The two-point distributed random variable X_0 with $\mathbb{P}(X_0 = x_) = p_*$, $\mathbb{P}(X_0 = y_*) = 1 - p_*$ is a minimizer of $\inf_{X \in \mathcal{X}} \mathbb{E}(X \wedge c)$, i.e.*

$$\inf_{X \in \mathcal{X}} \mathbb{E}(X \wedge c) = L_c(p_*) = \mathbb{E}(X_0 \wedge c).$$

If the parameters of the problem are such that X_0 , as defined in Proposition 4.2, is bounded below by $-m$, the original minimization problem is clearly solved. We shall assume, until further notice, that $m \geq 0$ and $c > -m$. This comes at no loss of generality, as \mathcal{X}^{-m} is empty if $m < 0$, and the problem degenerates if $c \leq -m$, in the sense that $\mathbb{E}(X \wedge c) = c$ for every $X \geq -m$.

Corollary 4.2. *Let X_0 be defined as in Proposition 4.2. Then*

$$\inf_{X \in \mathcal{X}^{-m}} \mathbb{E}(X \wedge c) = \mathbb{E}(X_0 \wedge c)$$

if and only if $c \geq (1 - m^2)/2m$.

Proof. By definition of X_0 , the lower bound $X_0 \geq -m$ holds if and only if $p_* \geq 1/(1 + m^2)$, i.e. if and only if

$$\frac{1}{1 + m^2} \leq \frac{1}{2} + \frac{1}{2} \frac{c}{(1 + c^2)^{1/2}},$$

or, equivalently,

$$2 \frac{1}{1 + m^2} - 1 \leq \frac{c}{(1 + c^2)^{1/2}},$$

where the left-hand side takes values in the interval $] -1, 1]$. Elementary algebra shows that the inequality

$$\beta \leq \frac{c}{(1 + c^2)^{1/2}}, \quad \beta \in] -1, 1],$$

is verified if and only if

$$c \geq \frac{\beta}{(1 - \beta^2)^{1/2}}.$$

Replacing β by $2/(1 + m^2) - 1$ finally implies that $X_0 \geq -m$ if and only if

$$c \geq \frac{1 - m^2}{2m},$$

from which the claim follows immediately. □

In view of the corollary, we only need to consider the problem under the condition

$$c < \frac{1 - m^2}{2m}.$$

Note that this implies $c < 1/m$.

Let us start by observing that, for any $X \geq -m$,

$$\begin{aligned} \mathbb{E}(X \wedge c) &= \mathbb{E}(X\mathbf{1}_{\{X \leq c\}} + c\mathbf{1}_{\{X > c\}}) \\ &\geq -m\mathbb{P}(X \leq c) + c\mathbb{P}(X > c) \\ &= \mathbb{E}(Y \wedge c), \end{aligned} \tag{4.1}$$

where Y is a random variable taking values in $\{-m, y\}$, $y \geq c$, with

$$\mathbb{P}(Y = -m) = \mathbb{P}(X \leq c), \quad \mathbb{P}(Y = y) = \mathbb{P}(X > c).$$

In order for the random variable Y to belong to \mathcal{X} , it is sufficient and necessary, in view of (2.2) and Lemma 2.1, that

$$\mathbb{P}(Y = -m) = \mathbb{P}(X \leq c) = \frac{1}{1 + m^2},$$

and either $c \leq 0$ or $c \geq 0$ and

$$\mathbb{P}(X \leq c) \geq \frac{c^2}{1 + c^2}.$$

In other words, Y belongs to \mathcal{X} and takes values in $\{-m, y\}$ with $y \geq c$ if and only if $c \leq 0$ or $c > 0$ and

$$\frac{1}{1 + m^2} \geq \frac{c^2}{1 + c^2}.$$

As this inequality is satisfied if and only if $cm \leq 1$, which holds by assumption, Y satisfies the abovementioned conditions if and only if $\mathbb{P}(X \leq c) = 1/(1 + m^2)$. Let us then define the random variable $Y_0 \in \mathcal{X}^{-m}$ as the (unique) random variable in \mathcal{X} identified by the parameter $p_m = 1/(1 + m^2)$. We are going to show that Y_0 is in fact the minimizer of the problem.

Proposition 4.3. *If $c < (1 - m^2)/(2m)$, then*

$$\inf_{X \in \mathcal{X}^{-m}} \mathbb{E}(X \wedge c) = \mathbb{E}(Y_0 \wedge c) = c - (m + c) \frac{1}{1 + m^2}.$$

Proof. Let us rewrite (4.1) as $\mathbb{E}(X \wedge c) \geq -c + (m + c)\mathbb{P}(X \leq c)$, which holds for every $X \in \mathcal{X}^{-m}$. Since $m + c > 0$ by assumption, the function $p \mapsto -c + (m + c)p$ is decreasing. Therefore, for every $X \in \mathcal{X}^{-m}$ such that $\mathbb{P}(X \leq c) \leq p_m$, we have

$$\mathbb{E}(X \wedge c) \geq c - (m + c)p_m = \mathbb{E}(Y_0 \wedge c).$$

Let $X \in \mathcal{X}^{-m}$ be such that $p := \mathbb{P}(X \leq c) > p_m$. Then Lemma 4.2 yields

$$\mathbb{E}(X \wedge c) \geq -(p(1-p))^{1/2} + c(1-p) = L_c(p).$$

Since $p_* < p_m$ by assumption and the function L_c is increasing on $]p_*, 1[$ by Lemma 4.4, it follows that $\mathbb{E}(X \wedge c) \geq L_c(p) \geq L_c(p_m) = \mathbb{E}(X_0 \wedge c)$, which concludes the proof. \square

We have therefore proved the main result, which reads as follows.

Theorem 4.2. *Let $c, m, \sigma \in \mathbb{R}$ with $m \geq 0, \sigma > 0$, and $c > -m$. Then*

$$\inf_{X \in \mathcal{X}^{-m}} \mathbb{E}(X \wedge c) = \begin{cases} \mathbb{E}(Y_0 \wedge c) = L_c(p_m) = c - (m+c) \frac{1}{1+m^2} & \text{if } c \leq \frac{1-m^2}{2m}, \\ \mathbb{E}(X_0 \wedge c) = L_c(p_*) = \frac{1}{2}c - \frac{1}{2}(1+c^2)^{1/2} & \text{if } c \geq \frac{1-m^2}{2m}. \end{cases}$$

Using the notation $X(p)$ to denote a two-point distributed random variable in \mathcal{X} with parameter p , we could write, more concisely,

$$\inf_{X \in \mathcal{X}^{-m}} \mathbb{E}(X \wedge c) = \mathbb{E}(X(p_* \vee p_m) \wedge c) = L_c(p_* \vee p_m).$$

The bound by Scarf, and hence the one by Lo, i.e. Theorem 4.1 and its corollary, follow immediately by the previous theorem and Lemma 4.1.

5. Applications to option prices

In order to also consider bounds for put options, we record an easy consequence of Theorem 3.1.

Corollary 5.1. *Under the hypotheses of Theorem 3.1, let $p := \mathbb{P}(X \leq c)$. Then*

$$(c-m)p \leq \mathbb{E}(c-X)^+ \leq (c-m)p + \sigma(p-p^2)^{1/2}.$$

Proof. It immediately follows from the identities $\mathbb{E}(c-X)^+ = \mathbb{E}(X-c)^+ - (m-c)$ and $p-p^2 = p(1-p) = (1-p_0)p_0 = p_0 - p_0^2$. \square

Let us also note that a simple but useful sharpening of the lower bound in Theorem 3.1 can be given: Jensen’s inequality implies that

$$\mathbb{E}(X-c)^+ \geq (\mathbb{E}X-c)^+ = (m-c)^+$$

as well as $\mathbb{E}(c-X)^+ \geq (c-m)^+$.

Assume that the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is equipped with a filtration $(\mathcal{F}_t)_{t \in [0, T]}$ satisfying the so-called usual conditions. Let \widehat{S} and β be the price processes of two traded assets, the latter of which is strictly positive and is used as numéraire, so that $S := \beta^{-1}\widehat{S}$ is the discounted price process of the former asset. We assume that no asset pays dividends. A classical result [6] asserts that a suitable version of no-arbitrage holds (precisely, no free lunch with vanishing risk) if and only if there exists a probability measure \mathbb{Q} equivalent to \mathbb{P} such that S is a σ -martingale with respect to \mathbb{Q} . For simplicity of notation, let us assume that \mathbb{P} is already an equivalent σ -martingale measure that is used for pricing. We are then interested in upper and lower bounds for

$$\pi_c := \mathbb{E}\beta_T^{-1}(\widehat{S}_T - K)^+ = \mathbb{E}(S_T - \beta_T^{-1}K)^+, \quad \pi_p := \mathbb{E}\beta_T^{-1}(K - \widehat{S}_T)^+ = \mathbb{E}(\beta_T^{-1}K - S_T)^+.$$

We first establish an easy consequence of the σ -martingale property of S .

Lemma 5.1.

- (i) If S is a supermartingale (in particular, if S is bounded from below), then $\pi_p \geq (K\mathbb{E}\beta_T^{-1} - S_0)^+$.
- (ii) If S is a martingale, then $\pi_c \geq (S_0 - K\mathbb{E}\beta_T^{-1})^+$.

Proof. Recall first that a σ -martingale bounded from below is a local martingale thanks to the Ansel–Stricker lemma [1, p. 309], thus also a supermartingale by an application of Fatou’s lemma. This implies that $\mathbb{E}S_T \leq S_0$, and hence, by Jensen’s inequality,

$$\pi_p = \mathbb{E}(\beta_T^{-1}K - S_T)^+ \geq (K\mathbb{E}\beta_T^{-1} - \mathbb{E}S_T)^+ \geq (K\mathbb{E}\beta_T^{-1} - S_0)^+,$$

which proves (i). The proof of (ii) is entirely analogous. □

Remark 5.1. The lower bound for π_c is in general not true without the hypothesis that S is a martingale. This is essentially equivalent to the failure of put–call parity for asset prices with so-called bubbles (cf., e.g., [4]).

It is clear that the estimates of Theorem 3.1 and Corollary 4.1 cannot be directly applied, as β_T is a random variable. In some cases, however, this is indeed possible. For instance, apart from the trivial case where the price process β of the numéraire is non-random, if the random variables β_T^{-1} and \widehat{S}_T^+ belong to L^2 , and β_T^{-1} and $(S_T - K)^+$ are uncorrelated, so that

$$\mathbb{E}\beta_T^{-1}(\widehat{S}_T - K)^+ = \mathbb{E}\beta_T^{-1}\mathbb{E}(\widehat{S}_T - K)^+,$$

estimates on π_c can be obtained by Corollary 4.1 in terms of the mean of β_T^{-1} and the mean and variance of \widehat{S}_T . In order to apply Theorem 3.1, the value of the distribution function of \widehat{S}_T at K is also needed. In the following we make the stronger assumption that the random variables β_T and \widehat{S}_T are independent. Furthermore, we assume that S is a supermartingale. By independence,

$$S_0 \geq \mathbb{E}S_T = \mathbb{E}\beta_T^{-1}\widehat{S}_T = \mathbb{E}\beta_T^{-1}\mathbb{E}\widehat{S}_T,$$

and hence $\mathbb{E}\widehat{S}_T \leq S_0/\mathbb{E}\beta_T^{-1}$. Setting $k := K\mathbb{E}\beta_T^{-1}$ and $\widehat{\sigma} := \|\widehat{S}_T - \mathbb{E}\widehat{S}_T\|_2$, we thus have the bounds

$$\begin{aligned} (k - S_0)^+ &\leq \pi_p \leq (k - \mathbb{E}S_T)p + \mathbb{E}\beta_T^{-1}\widehat{\sigma}(p - p^2)^{1/2}, \\ (\mathbb{E}S_T - k)^+ &\leq \pi_c \leq (S_0 - k)p_0 + \mathbb{E}\beta_T^{-1}\widehat{\sigma}(p_0 - p_0^2)^{1/2}. \end{aligned}$$

If S is a martingale, then $\mathbb{E}S_T = S_0$, so the bounds for π_p and π_c assume a more symmetric form.

Adapting Lo’s estimate to option prices can be done along the same lines. It does not seem possible, though, to exploit the supermartingale property of S to obtain bounds involving S_0 rather than $\mathbb{E}S_T$. On the other hand, if S is a martingale, and if the constraint $\widehat{S}_T \geq 0$ is not enforced, it follows from the proofs in Section 4 and elementary computations that

$$\pi_c \leq \frac{S_0 - k}{2} + \frac{1}{2}((S_0 - k)^2 + (\mathbb{E}\beta_T^{-1})^2\widehat{\sigma}^2)^{1/2},$$

from which a corresponding upper bound for put options can be obtained by put–call parity.

References

- [1] ANSEL, J. P. AND STRICKER, C. (1994). Couverture des actifs contingents et prix maximum. *Ann. Inst. H. Poincaré Prob. Statist.* **30**, 303–315.
- [2] BERTSIMAS, D. AND POPESCU, I. (2002). On the relation between option and stock prices: A convex optimization approach. *Operat. Res.* **50**, 358–374.
- [3] BOURBAKI, N. (1970). *Théorie des ensembles*. Hermann, Paris.
- [4] COX, A. M. G. AND HOBSON, D. G. (2005). Local martingales, bubbles and option prices. *Finance Stoch.*, **9**, 477–492.
- [5] DE LA PEÑA, V. H., IBRAGIMOV, R. AND JORDAN, S. (2004). Option bounds. *J. Appl. Prob.* **41A**, 145–156.
- [6] DELBAEN, F. AND SCHACHERMAYER, W. (1998). The fundamental theorem of asset pricing for unbounded stochastic processes. *Math. Ann.* **312**, 215–250.
- [7] LO, A. W. (1987). Semi-parametric upper bounds for option prices and expected payoffs. *J. Financial Econometrics* **19**, 373–387.
- [8] MARINELLI, C. (2023). On some semi-parametric estimates for European option prices. Preprint, [arXiv:2306.10929](https://arxiv.org/abs/2306.10929).
- [9] PINELIS, I. (2011). Exact lower bounds on the exponential moments of truncated random variables. *J. Appl. Prob.* **48**, 547–560.
- [10] PINELIS, I. AND MOLZON, R. (2009). Berry–Esseen bounds for general nonlinear statistics, with applications to Pearson’s and non-central Student’s and Hotelling’s. Preprint, [arXiv:0906.0177](https://arxiv.org/abs/0906.0177).
- [11] SCARF, H. (1958). A min–max solution of an inventory problem. In *Studies in the Mathematical Theory of Inventory and Production*, eds K. J. Arrow, S. Karlin, and H. Scarf. Stanford University Press, pp. 201–209.