

## ON ASYMPTOTIC VALUES OF SLOWLY GROWING ALGEBROID FUNCTIONS

JUNJI SUZUKI

1. Let  $f(z)$  be a  $k$ -valued algebroid function in  $|z| < \infty$  and

$$(1) \quad F(z, f) \equiv A_0(z)f^k + A_1(z)f^{k-1} + \cdots + A_k(z) = 0$$

be its defining equation such that the coefficients  $A_i(z)$  ( $i = 0, 1, \dots, k$ ) are entire functions without any common zero and the left hand side is irreducible. We denote by  $\mathfrak{X}$  the  $k$ -sheeted covering surface over  $|z| < \infty$  generated by  $f(z)$  and by  $\mathfrak{X}(r)$  and  $\Gamma(r)$  the part of  $\mathfrak{X}$  over  $|z| \leq r$  and the curves on  $\mathfrak{X}$  over  $|z| = r$ , respectively. We use the standard notations of the Nevanlinna-Selberg theory [4]:

$$m(r, a) = \frac{1}{2k\pi} \int_{\Gamma(r)} \log^+ \left| \frac{1}{f(re^{i\theta}) - a} \right| d\theta, \quad m(r, f) = \frac{1}{2k\pi} \int_{\Gamma(r)} \log^+ |f(re^{i\theta})| d\theta$$

$$N(r, a) = \frac{1}{k} \int_0^r \frac{n(t, a) - n(0, a)}{t} dt + \frac{n(0, a)}{k} \log r, \quad N(r, \infty) = N(r, f)$$

$$T(r, f) = m(r, f) + N(r, f), \quad \delta(a, f) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N(r, a)}{T(r, f)},$$

where  $n(r, a)$  is the number of zeros of  $f(z) - a$  on  $\mathfrak{X}(r)$  and  $n(r, \infty) = n(r, f)$ .

From now on, we consider the functions with the slow growth:

$$(2) \quad T(r, f) = O[(\log r)^2].$$

For such functions both of the number of deficient values and that of asymptotic values are at most  $k$  (Valiron [7], [9] and Tumura [5]). Especially, when  $k=1$  i.e. the function is single-valued and meromorphic, it can possess no deficient value without that value being an asymptotic value (Valiron [9] and Anderson-Clunie [1]).

For an algebroid function  $f(z)$ , a value  $\alpha$  is an asymptotic value, if there exists a path  $L_{\mathfrak{X}}$  on  $\mathfrak{X}$  stretching to the point at infinity such that  $f(z)$

---

Received November 27, 1969.

tends to  $\alpha$  along  $L_x$ , in other words, if there exists a path  $L$  on the  $z$ -plane stretching to the point at infinity such that at least one branch of  $f(z)$  can be continued analytically along  $L$  and the value taken by the branch tends to  $\alpha$  along  $L$ .

Our main aim in this note is to give an extension of the above result of Anderson-Clunie to the case of an algebroid function:

**THEOREM 1.** *Let  $f(z)$  be a  $k$ -valued algebroid function in  $|z| < \infty$  satisfying (2). If  $f(z)$  has  $k$  deficient values  $\alpha_i$  ( $i=1, 2, \dots, k$ ), then each of  $\alpha_i$  ( $i=1, 2, \dots, k$ ) is an asymptotic value of  $f(z)$ .*

This theorem will be obtained as an immediate corollary of Theorem 2 stated in §5. In the last section, we shall give a condition for a deficient value to be an asymptotic value without the restriction that  $f(z)$  has  $k$  deficient values.

2. First we shall give some lemmas. To prove them, we use the following results.

I. (Valiron [6]) *If  $f(z)$  is a  $k$ -valued algebroid function in  $|z| < \infty$ , then*

$$(3) \quad \left| T(r, f) + \frac{1}{k} \log |C_1| - \mu(r, A) \right| < \log 2,$$

where  $\mu(r, A) = \frac{1}{2k\pi} \int_0^{2\pi} \log A(re^{i\theta}) d\theta$  with  $A(z) = \max_{0 \leq i \leq k} |A_i(z)|$  and  $C_1 z^1$  is the first non-zero term of the Taylor development of  $A_0(z)$  at the origin.

II. (Valiron [9]) *If  $f(z)$  is a  $k$ -valued algebroid function in  $|z| < \infty$  satisfying (2), and if  $a_i$  ( $i=1, 2, \dots, k+1$ ) are  $k+1$  distinct complex numbers (may be infinity), then we have*

$$\lim_{r \rightarrow \infty} \frac{N(r, a_1, a_2, \dots, a_{k+1})}{kT(r, f)} = 1$$

where  $N(r, a_1, a_2, \dots, a_{k+1}) = \max_{1 \leq i \leq k+1} N\left(r, \frac{1}{F(z, a_i)}\right)$  for each  $r > 0$ .

III. (Valiron [8]) *If  $g(z)$  is an entire function of order zero with  $g(0) = 1^1$ , then*

$$\log M(r, g) = N\left(r, \frac{1}{g}\right) + \theta(r)W\left(r, \frac{1}{g}\right) \quad (0 < \theta(r) < 1),$$

<sup>1)</sup> This condition is not essential to obtain (4).

where  $M(r, g) = \max_{|z|=r} |g(z)|$  and  $W\left(r, \frac{1}{g}\right) = r \int_0^\infty n\left(t, \frac{1}{g}\right) \frac{dt}{t^2}$ .

In particular, if  $\log M(r, g) = O[(\log r)^2]$ , then

$$\log M(r, g) < K(\log r)^2 \quad (K: \text{constant})$$

$$n\left(r, \frac{1}{g}\right) \log r = \int_r^{r^2} n\left(t, \frac{1}{g}\right) \frac{dt}{t} \leq \int_r^{r^2} n\left(t, \frac{1}{g}\right) \frac{dt}{t} < K(\log r^2)^2$$

$$= K'(\log r)^2$$

$$W\left(r, \frac{1}{g}\right) < K'r \int_0^\infty \frac{\log t}{t^2} dt = K'r \frac{\log r + 1}{r} = O(\log r),$$

so that we have

$$(4) \quad \log M(r, g) \sim N\left(r, \frac{1}{g}\right) \quad (r \rightarrow \infty).$$

IV. (Hayman [3]) If an entire function  $g(z)$  satisfies

$$\log M(r, g) = O[(\log r)^2],$$

then

$$(5) \quad \log M(r, g) \sim \log |g(z)|,$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set.

Here we call an  $\mathcal{E}$ -set any countable set of circles not containing the origin and subtending angles at the origin whose sum  $s$  is finite. We note the following two facts about  $\mathcal{E}$ -sets.

a) The union of two  $\mathcal{E}$ -sets is again an  $\mathcal{E}$ -set.

b) Given any  $\mathcal{E}$ -set then for almost all fixed  $\theta$  and any  $r > r_0(\theta)$ , where  $r_0(\theta)$  depends only on  $\theta$ ,  $z = re^{i\theta}$  lies outside the  $\mathcal{E}$ -set.

We consider a system  $\mathfrak{S}(z) = (S_0(z), S_1(z), \dots, S_k(z))$  of  $k + 1$  entire functions  $S_i(z)$  ( $i = 0, 1, \dots, k$ ) having no common zero and satisfying

$$(6) \quad \log M(r, S_i) = O[(\log r)^2] \quad (i = 0, 1, \dots, k).$$

We define  $\mu(r, S)$  by

$$\mu(r, S) = \frac{1}{2k\pi} \int_0^{2\pi} \log S(re^{i\theta}) d\theta,$$

where  $S(z) = \max_{0 \leq i \leq k} |S_i(z)|$  for each  $z$  and set

$$1 - \overline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{S_i}\right)}{k\mu(r, S)} = \delta_i(\mathfrak{S}) \quad (i = 0, 1, \dots, k).$$

Particularly, when  $\lim_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{S_i}\right)}{k\mu(r, S)}$  exists, we set

$$1 - \lim_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{S_i}\right)}{k\mu(r, S)} = \bar{\delta}_j(\mathfrak{S}).$$

Then we have  $0 \leq \delta_i(\mathfrak{S}) \leq 1$  ( $i = 0, 1, \dots, k$ ), since by Jensen’s formula

$$\begin{aligned} N\left(r, \frac{1}{S_i}\right) &= \frac{1}{2\pi} \int_0^{2\pi} \log |S_i(re^{i\theta})| d\theta - \log |S_i(0)|^{(2)} \\ &\leq \frac{k}{2k\pi} \int_0^{2\pi} \log S(re^{i\theta}) d\theta + O(1) = k\mu(r, S) + O(1). \end{aligned}$$

LEMMA 1. For a system  $\mathfrak{S}(z) = (S_0(z), S_1(z), \dots, S_k(z))$ , if  $\delta_j(\mathfrak{S}) > 0$  for some  $j(0 \leq j \leq k)$ , then

$$\lim_{r \rightarrow \infty} \frac{-\log \frac{|S_i(z)|^2}{\sum_0^k |S_i(z)|^2}}{2k\mu(r, S)} \geq \delta_j(\mathfrak{S}) > 0,$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set.

Proof. From our hypothesis, we have

$$N\left(r, \frac{1}{S_i}\right) < (1 - \delta_j(\mathfrak{S}) + o(1))k\mu(r, S).$$

Since  $\mathfrak{S}(z)$  satisfies (6), we can apply (4) and (5) to  $S_j(z)$  and have

$$(7) \quad \log |S_i(z)| < (1 - \delta_j(\mathfrak{S}) + o(1))k\mu(r, S),$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set.

By Cauchy’s inequality, we have for all  $\nu$  ( $\nu = 0, 1, \dots, k$ )

$$\begin{aligned} \log \left( \sum_{i=0}^k |S_i(z)|^2 \right) &\geq \log \left\{ \frac{1}{k+1} \left( \sum_{i=0}^k |S_i(z)|^2 \right) \right\} = 2\log \left( \sum_{i=0}^k |S_i(z)| \right) + \log \frac{1}{k+1} \\ &\geq 2\log |S_\nu(z)| + \log \frac{1}{k+1}. \end{aligned}$$

<sup>2)</sup> We assume that  $S_i(0) \neq 0, \infty$ .

Applying (5) to  $S_\nu(z)$ , we have for all  $\nu(\nu = 0, 1, \dots, k)$

$$\log \left( \sum_{i=0}^k |S_i(z)|^2 \right) \geq 2(1 + o(1)) \log M(r, S_\nu) + \log \frac{1}{k+1},$$

and hence

$$\log \left( \sum_{i=0}^k |S_i(z)|^2 \right) \geq 2(1 + o(1)) \max_{0 \leq \nu \leq k} \log M(r, S_\nu) + \log \frac{1}{k+1},$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set.

On the other hand, by definition of  $S(z)$ ,

$$S(z) \leq \max_{0 \leq \nu \leq k} M(r, S_\nu) \quad (|z| = r)$$

so that  $\mu(r, S) = \frac{1}{2k\pi} \int_0^{2\pi} \log S(re^{i\theta}) d\theta \leq \frac{1}{k} \max_{0 \leq \nu \leq k} \log M(r, S_\nu)$ . Thus we have

$$(8) \quad \log \left( \sum_{i=0}^k |S_i(z)|^2 \right) \geq 2k(1 + o(1))\mu(r, S),$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside the  $\mathcal{E}$ -set.

We combine (7) and (8) and have from the property a) of  $\mathcal{E}$ -sets,

$$\begin{aligned} \log \frac{|S_j(z)|^2}{\sum_{i=0}^k |S_i(z)|^2} &= 2 \log |S_j(z)| - \log \left( \sum_{i=0}^k |S_i(z)|^2 \right) \\ &\leq 2k(-\delta_j(\mathbb{C}) + o(1))\mu(r, S) \end{aligned}$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set. Thus we obtain the desired result.

By using the property b) of  $\mathcal{E}$ -sets and the fact that the function  $\mu(r, S)$  of  $r$  is unbounded, we have that

$$\frac{|S_j(z)|^2}{\sum_{i=0}^k |S_i(z)|^2} \rightarrow 0$$

as  $z = re^{i\theta} \rightarrow \infty$  for almost all fixed  $\theta$  ( $0 \leq \theta < 2\pi$ ).

**3.** Before giving the next lemma, we shall state some about the distance between two systems, which was introduced by Dufresnoy [2].

We consider only the systems consisting of  $k + 1$  complex numbers, all of which are not zero simultaneously. Here if two systems

$$w^{(1)} = (w_0^{(1)}, w_1^{(1)}, \dots, w_k^{(1)}) \text{ and } w^{(2)} = (w_0^{(2)}, w_1^{(2)}, \dots, w_k^{(2)})$$

are proportional i.e.  $w_i^{(1)} = cw_i^{(2)}$  ( $i = 0, 1, \dots, k$ ) for some constant  $c(c \neq 0)$ , we identify  $w^{(1)}$  with  $w^{(2)}$ .

We set

$$(9) \quad [[w^{(1)}, w^{(2)}]] = \left\{ \frac{\sum_{i>j} |w_i^{(1)}w_j^{(2)} - w_j^{(1)}w_i^{(2)}|^2}{\sum_{i=0}^k |w_i^{(1)}|^2 \sum_{i=0}^k |w_i^{(2)}|^2} \right\}^{\frac{1}{2}}$$

Then this satisfies three axioms for distances. According to Dufresnoy [2] we call  $[[w^{(1)}, w^{(2)}]]$  the distance between two systems  $w^{(1)}$  and  $w^{(2)}$ . We can easily see that an inequality

$$(10) \quad [[w^{(1)}, w^{(2)}]]^2 \leq \frac{\sum_{i=0}^k |w_i^{(1)} - w_i^{(2)}|^2}{\left\{ \sum_{i=0}^k |w_i^{(1)}|^2 \sum_{i=0}^k |w_i^{(2)}|^2 \right\}^{1/2}}$$

holds. This shows how our distance relates to the distance in ordinary sense between  $w^{(1)}$  and  $w^{(2)}$ .

Now we consider a non-degenerate, linear and homogeneous substitution of the elements of the system  $w = (w_0, w_1, \dots, w_k)$ ;

$$(11) \quad W_i = \sum_{j=0}^k a_{ij}w_j \quad (i = 0, 1, \dots, k).$$

Then we have a new system  $W = (W_0, W_1, \dots, W_k)$ . Let

$$W^{(1)} = (W_0^{(1)}, W_1^{(1)}, \dots, W_k^{(1)}) \text{ and } W^{(2)} = (W_0^{(2)}, W_1^{(2)}, \dots, W_k^{(2)})$$

be the systems obtained by the substitution (11) of the elements of systems  $w^{(1)}$  and  $w^{(2)}$ , respectively. Then, using the inequality (10) we have an important property about the distance (9) which is stated as follows;

LEMMA 2. (Dufresnoy [2]) *Under such a substitution, two systems being close to each other correspond to two systems also being close to each other i.e. there exists a constant  $c$ ,  $0 < c < 1$ , depending only on  $a_{ij}$  ( $i, j = 0, 1, \dots, k$ ) such that*

$$c[[w^{(1)}, w^{(2)}]] < [[W^{(1)}, W^{(2)}]] < c^{-1}[[w^{(1)}, w^{(2)}]].$$

Let 
$$p(z) = a_0z^k + a_1z^{k+1} + \dots + a_k = 0$$

$$p^*(z) = a_0^*z^k + a_1^*z^{k-1} + \dots + a_k^* = 0$$

be two algebraic equations whose coefficients make systems  $a = (a_0, a_1, \dots, a_k)$  and  $a^* = (a_0^*, a_1^*, \dots, a_k^*)$ , respectively. By means of distance (9), the well

known theorem on continuity of roots of algebraic equations is described as follows;

LEMMA 3. (Dufresnoy [2]) Let  $z_1, z_2, \dots, z_k$  and  $z_1^*, z_2^*, \dots, z_k^*$  be the roots of the equations  $p(z) = 0$  and  $p^*(z) = 0$ , respectively. If  $[[a, a^*]]$  is sufficiently small, then we can associate each  $z_i (i = 0, 1, \dots, k)$  with some  $z_j^* (1 \leq j \leq k)$ , say  $z_i$  with  $z_{\alpha_i}^*$ , such that

$$[z_i, z_{\alpha_i}^*] < 8e[[a, a^*]]^{\frac{1}{k}} \quad (i = 1, 2, \dots, k),$$

where  $[ \quad , \quad ]$  denotes the chordal distance.

The next lemma is an immediate consequence of Lemma 3.

LEMMA 4. (Dufresnoy [2]) If

$$\frac{\sum_{i=0}^p |a_i|^2}{\sum_{j=0}^k |a_j|^2} \quad (0 \leq p \leq k - 1)$$

is sufficiently small, then an algebraic equation

$$p(z) = a_0 z^k + a_1 z^{k-1} + \dots + a_k = 0$$

has at least  $p + 1$  roots whose chordal distances from the point at infinity are less than

$$8e \left\{ \frac{\sum_{i=0}^p |a_i|^2}{\sum_{j=0}^k |a_j|^2} \right\}^{\frac{1}{2k}}.$$

For the sake of the later discussion, we shall give a proof following Dufresnoy [2].

*Proof.* We consider one more equation

$$p^*(z) = a_0^* z^k + a_1^* z^{k-1} + \dots + a_k^* = 0$$

with  $a_i^* = 0 (i = 0, 1, \dots, p)$  and  $a_j^* = a_j (j = p + 1, \dots, k)$ . Then we have

$$[[a, a^*]] = \left\{ \frac{\sum_{i=0}^p |a_i|^2}{\sum_{j=0}^k |a_j|^2} \right\}^{\frac{1}{2}}$$

We may consider that the equation  $p^*(z) = 0$  has  $k$  roots,  $p + 1$  of them lying at the point at infinity. Thus our Lemma is obtained from Lemma 3. Here we note that each of the other  $k - p - 1$  roots  $z_i (i = 1, 2, \dots, k - p - 1)$  of  $p(z) = 0$  is associated with one of the  $k - p - 1$  roots  $z_i^* (i = 1, 2, \dots, k - p - 1)$  of  $p^*(z) = 0$ , say  $z_l$  with  $z_{\alpha_l}^*$ , in such a way that

$$[z_l, z_{\alpha_l}^*] < 8e \left\{ \frac{\sum_{i=0}^p |a_i|^2}{\sum_{j=0}^k |a_j|^2} \right\}^{\frac{1}{2k}} \quad (l = 1, 2, \dots, k - p - 1).$$

4. LEMMA 5. Let  $\mathfrak{S}(z) = (S_0(z), S_1(z), \dots, S_k(z))$  be a system such that  $S_i(z) (j = 0, 1, \dots, k)$  have no common zero and satisfy (6). If  $\delta_\lambda(\mathfrak{S}) = 0$  for only one  $\lambda (0 \leq \lambda \leq k)$  and  $\delta_\nu(\mathfrak{S}) > 0$  for other all  $\nu \neq \lambda (0 \leq \nu \leq k)$ , then

$$[[\mathfrak{S}(z_1), \mathfrak{S}(z_2)]] \rightarrow 0$$

uniformly in  $\theta_m$  as  $z_m = r_m e^{i\theta_m} \rightarrow \infty$  outside an  $\mathcal{E}$ -set ( $m = 1, 2$ ).

*Proof.* For any pair  $(i, j) (i \neq j; i, j = 0, 1, \dots, k)$ ,

$$\begin{aligned} \frac{|S_i(z_1)S_j(z_2) - S_j(z_1)S_i(z_2)|}{\left\{ \sum_{h=0}^k |S_h(z_1)|^2 \sum_{h=0}^k |S_h(z_2)|^2 \right\}^{\frac{1}{2}}} &\leq \frac{|S_i(z_1)S_j(z_2)|}{\left\{ \sum_{h=0}^k |S_h(z_1)|^2 \sum_{h=0}^k |S_h(z_2)|^2 \right\}^{\frac{1}{2}}} \\ + \frac{|S_j(z_1)S_i(z_2)|}{\left\{ \sum_{h=0}^k |S_h(z_1)|^2 \sum_{h=0}^k |S_h(z_2)|^2 \right\}^{\frac{1}{2}}} &\leq \min_{i=j} \left\{ \frac{|S_i(z_1)|}{\left( \sum_{h=0}^k |S_h(z_1)|^2 \right)^{\frac{1}{2}}} + \frac{|S_i(z_2)|}{\left( \sum_{h=0}^k |S_h(z_2)|^2 \right)^{\frac{1}{2}}} \right\}. \end{aligned}$$

By Lemma 1 and our hypotheses, we have for all  $\nu (\neq \lambda)$

$$\frac{|S_\nu(z)|}{\left( \sum_{h=0}^k |S_h(z)|^2 \right)^{\frac{1}{2}}} \rightarrow 0$$

uniformly in  $\theta$  as  $z = r e^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set, and hence

$$\frac{|S_i(z_1)S_j(z_2) - S_j(z_1)S_i(z_2)|}{\left( \sum_{h=0}^k |S_h(z_1)|^2 \sum_{h=0}^k |S_h(z_2)|^2 \right)^{\frac{1}{2}}} \rightarrow 0.$$

uniformly in  $\theta_m$  as  $z_m = r_m e^{i\theta_m} \rightarrow \infty$  outside an  $\mathcal{E}$ -set ( $m = 1, 2$ ). Thus our lemma is obtained.



**COROLLARY.** *Let  $f(z)$  be a  $k$ -valued algebroid function in  $|z| < \infty$  satisfying (2). Suppose that  $f(z)$  has  $k$  deficient values  $\alpha_i (i = 1, 2, \dots, k)$ . Then for the system  $\mathfrak{A}(z) = (A_0(z), A_1(z), \dots, A_k(z))$ , we have the same assertion as that in the above lemma.*

*Proof.* We take a value  $\alpha_0$  which is different from  $\alpha_i (i = 1, 2, \dots, k)$  and set

$$(12) \quad F(z, \alpha_i) = A_0(z)\alpha_i^k + A_1(z)\alpha_i^{k-1} + \dots + A_k(z) = B_i(z) \quad (i = 0, 1, 2, \dots, k).$$

Now we shall prove that for the system  $\mathfrak{B}(z) = (B_0(z), B_1(z), \dots, B_k(z))$ , all the conditions of Lemma 5 are satisfied. At first, entire functions  $B_i(z) (i = 0, 1, \dots, k)$  have no common zero. In fact, suppose that  $B_i(z) (i = 0, 1, \dots, k)$  have a common zero  $a$ . We solve the equation (12) with respect to  $A_i(z) (i = 0, 1, \dots, k)$  and have

$$(13) \quad A_i(z) = \beta_{i0}B_0(z) + \beta_{i1}B_1(z) + \dots + \beta_{ik}B_k(z) \quad (i = 0, 1, \dots, k; \beta_{ij}; \text{ constants})$$

so that  $a$  is also a common zero of  $A_i(z) (i = 0, 1, \dots, k)$ , which is absurd. Further, we have from (12) and (13),

$$(14) \quad \mu(r, A) = \mu(r, B) + O(1)$$

so that  $B_i(z) (i = 0, 1, \dots, k)$  satisfy (6) by (2) and (3).

Next, since  $N\left(r, \frac{1}{f - \alpha_i}\right) = \frac{1}{k} N\left(r, \frac{1}{B_i}\right) (i = 0, 1, \dots, k)$  and  $\alpha_i (i = 1, 2, \dots, k)$  are deficient values of  $f(z)$ , we have by (3)

$$(15) \quad \delta_j(\mathfrak{B}) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{B_j}\right)}{kT(r, f)} = \delta(\alpha_j, f) > 0 \quad (j = 1, 2, \dots, k).$$

On the other hand, the value  $\alpha_0$  is normal by II in §2, i.e.

$$(16) \quad \bar{\delta}_0(\mathfrak{B}) = 1 - \lim_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{B_0}\right)}{kT(r, f)} = \delta(\alpha_0, f) = 0.$$

Now Lemma 5 applied to the system  $\mathfrak{B}(z) = (B_0(z), B_1(z), \dots, B_k(z))$  shows that

$$[[\mathfrak{B}(z_1), \mathfrak{B}(z_2)]] \rightarrow 0$$

uniformly in  $\theta_m$  as  $z_m = r_m e^{i\theta_m} \rightarrow \infty$  outside an  $\mathcal{E}$ -set ( $m = 1, 2$ ).

Since we can take (12) as a non-degenerate, linear and homogeneous substitution of the elements  $A_i(z)$  of the system  $\mathfrak{A}(z) = (A_0(z), A_1(z), \dots, A_k(z))$ , we obtain the desired result by Lemma 2.

**5. THEOREM 2.** *Let  $f(z)$  be a  $k$ -valued algebroed function  $|z| < \infty$  of arbitrary order. Suppose that there exists a path  $L$  on the plane stretching to the point at infinity such that*

$$(17) \quad \frac{|A_0(z)|}{\left(\sum_{i=0}^k |A_i(z)|\right)^{\frac{1}{2}}} \rightarrow 0$$

$$(18) \quad [[\mathfrak{A}(z_1), \mathfrak{A}(z_2)]] \rightarrow 0$$

as  $z, z_1$  and  $z_2$  tend to infinity along  $L$ . Then the infinity is an asymptotic value of  $f(z)$ .

*Proof.* We denote by  $K(\delta)$  the spherical disk with center at the point at infinity and with chordal radius  $\delta > 0$ , and denote by  $f_i(z)$  ( $i = 1, 2, \dots, k$ )  $k$  roots of  $F(z, f) = 0$  for any  $z$  counting with their proper multiplicities. We express the curve  $L$  by

$$L : z = z(t) \quad (0 < t < \infty); \quad z(t) \rightarrow \infty \text{ as } t \rightarrow \infty.$$

Given a sufficiently small  $\varepsilon > 0$ , we can find from (17) and (18)  $t_0^{(n)}$  ( $n = 1, 2, \dots$ ) depending on  $\varepsilon$  such that for any  $t \geq t_0^{(n)}$ ,

$$(19) \quad 8e \left\{ \frac{|A_0(z)|^2}{\sum_{i=0}^k |A_i(z)|^2} \right\}^{\frac{1}{2k}} < \frac{\varepsilon}{2(k+1)^n} \quad (z = z(t))$$

and for any pair  $t_1$  and  $t_2; t_1, t_2 \geq t_0^{(n)}$ ,

$$(20) \quad 8e [[\mathfrak{A}(z_1), \mathfrak{A}(z_2)]]^{\frac{1}{k}} < \frac{\varepsilon}{2(k+1)^n} \quad (z_i = z(t_i); i = 1, 2).$$

First we take whole branches  $f_i$  ( $i = 1, 2, \dots, k$ ) as our candidates and let  $z$  go to infinity along  $L$ . Then we drop from the list of candidates branches  $f_i$ , if any, with  $f_i(z(t_0^{(1)})) \notin K(\varepsilon)$ . The disk  $K\left(\frac{\varepsilon}{2(k+1)}\right)$  contains

at least one root of the equation  $F(z(t_0^{(1)}), f) = 0$  because of Lemma 4 and (19) and so there remains at least one  $f_j$  in our list. Next we drop  $f_i$ , if any, with  $f_i(z(t_0^{(2)})) \notin K\left(\frac{\varepsilon}{k+1}\right)$  from our 2nd list and still have a list containing at least one  $f_j$  by the same reason as above. Then we see that, for any  $f_j$  in the list, the curve  $f_j(z(t))$ ,  $t_0^{(1)} \leq t \leq t_0^{(2)}$ , is contained in  $K(\varepsilon)$ . In fact, if not, the curve  $f_j(z(t))$ ,  $t_0^{(1)} \leq t \leq t_0^{(2)}$ , can not be covered by any  $k$  disks with radii  $\frac{\varepsilon}{2(k+1)}$  and so there exists at least one point  $z^* = z(t^*)$ ,  $t_0^{(1)} < t^* < t_0^{(2)}$ , such that

$$[f_j(z^*), f_i(z(t_0^{(1)}))] > \frac{\varepsilon}{2(k+1)} \quad (i = 1, 2, \dots, k),$$

which contradicts Lemma 3 and (20). We repeat the above procedures and, at the  $n$ -th step, we drop  $f_i$ , if any, with  $f_i(z(t_0^{(n)})) \notin K\left[\frac{\varepsilon}{(k+1)^{n-1}}\right]$  from our  $n$ -th list, and have the  $(n+1)$ -th list containing at least one  $f_j$ . For any  $f_j$  in this list, the curve  $f_j(z(t))$ ,  $t_0^{(n-1)} \leq t \leq t_0^{(n)}$ , is contained in  $K\left[\frac{\varepsilon}{(k+1)^{n-2}}\right]$ . Since we have only a finite number of branches  $f_i$ , there is at least one  $f_j$ , say  $f_1$ , which belongs to the  $n$ -th list for  $n = 1, 2, \dots$ . Thus  $f_1$  satisfies

$$f_1(z(t)) \in K\left[\frac{\varepsilon}{(k+1)^{n-2}}\right], \quad t \geq t_0^{(n-1)},$$

so that  $f_1(z)$  tends to infinity as  $z$  goes to infinity along  $L$ . The proof is now complete.

*Proof of Theorem 1.* When  $\alpha_i \neq \infty$ , we consider  $\frac{1}{f - \alpha_i}$  instead of  $f$ . Then  $\frac{1}{f - \alpha_i}$  is an algebroid function satisfying (2) and has  $k$  deficient values, one of which is the infinity, so that we may assume that  $\alpha_i = \infty$ . From Lemma 1 and Corollary of Lemma 5, the coefficients  $A_0(z), A_1(z), \dots, A_k(z)$  of the defining equation of  $f(z)$  satisfying the conditions (17) and (18) outside an  $\mathcal{E}$ -set, consequently on any half-line  $L = re^{i\theta} (r > 0)$  for almost every  $\theta$ . Applying Theorem 2, we conclude that  $\alpha_i$  is an asymptotic value of  $f$  along  $L$ .

*Remark.* As we saw in the above proof, we can take any half-line  $L$  for almost every  $\theta$  as an asymptotic path of  $\alpha_i$  and hence an  $L$  commonly to all  $\alpha_i$ ;  $i = 1, 2, \dots, k$ .

**6. LEMMA 6.** (*Dufresnoy* [2]) *Let*  $p(z) = a_0z^\nu + a_1z^{\nu-1} + \dots + a_\nu = 0$  *be an algebraic equation with*

$$\frac{|a_0|^2}{\sum_{i=0}^{\nu} |a_i|^2} = \frac{\nu}{1 + M^2} \quad (M > 0).$$

*Then*  $p(z) = 0$  *has no root of modulus larger than*  $M$ .

From this, we can see that if

$$\frac{|a_0|^2}{\sum_{i=0}^{\nu} |a_i|^2} = \nu d^2 \quad (d > 0),$$

every root of  $p(z) = 0$  lies outside a spherical disk  $K(d)$  with center at the point at infinity and with chordal radius  $d$ . Using this lemma, we can prove

**THEOREM 3<sup>3)</sup>.** *Let*  $f(z)$  *be a*  $k$ -*valued algebroid function in*  $|z| < \infty$  *which is defined by (1) and satisfies (2). Suppose that, for some*  $n(0 < n \leq k)$ , *the system*  $\mathfrak{A}(z) = (A_0(z), A_1(z), \dots, A_k(z))$  *satisfies*

$$\delta_j(\mathfrak{A}) > 0 \quad (j = 0, 1, \dots, n-1), \quad \bar{\delta}_n(\mathfrak{A}) = 0.$$

*Then the infinity is an asymptotic value of*  $f(z)$ .

*Proof.* From our hypothesis  $\bar{\delta}_n(\mathfrak{A}) = 0$  and (3), we have  $\lim_{r \rightarrow \infty} \frac{N(r, \frac{1}{A_n})}{kT(r, f)} = 1$ . Hence we have by (4) and (5)

$$\log |A_n(z)|^2 = (1 + o(1))2kT(r, f),$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set. Further, we have

$$\begin{aligned} \log \left( \sum_{i=0}^k |A_i(z)|^2 \right) &\leq \log \left( \sum_{i=0}^k |A_i(z)|^2 \right) \leq 2 \log A(z) + \log(k+1) \\ &\leq 2 \max_{0 \leq \nu \leq k} \log M(r, A_\nu) + \log(k+1) = 2(1 + o(1)) \max_{0 \leq \nu \leq k} N\left(r, \frac{1}{A_\nu}\right) \\ &\leq (1 + o(1))2kT(r, f). \end{aligned}$$

Thus

<sup>3)</sup> As for notations used in this theorem, see § 2.

$$\log \frac{|A_n(z)|^2}{\sum_{i=n}^k |A_i(z)|^2} = o[T(r, f)]$$

and hence for any small  $\varepsilon > 0$ ,

$$e^{-\varepsilon T(r, f)} < \left( \frac{1}{k-n} \frac{|A_n(z)|^2}{\sum_{i=n}^k |A_i(z)|^2} \right)^{\frac{1}{2}} < e^{\varepsilon T(r, f)}$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside the  $\mathcal{E}$ -set. Since  $\delta_j(\mathfrak{A}) > 0$  ( $j = 0, 1, \dots, n-1$ ), we see from Lemma 1,

$$\log \frac{|A_j(z)|^2}{\sum_{i=0}^k |A_i(z)|^2} < (-\delta_j(\mathfrak{A}) + o(1))2kT(r, f) \quad (j = 0, 1, \dots, n-1)$$

and hence

$$(22) \quad 8e \left\{ \frac{\sum_{j=0}^{n-1} |A_j(z)|^2}{\sum_{i=0}^k |A_i(z)|^2} \right\}^{\frac{1}{2k}} < e^{(-\delta + \varepsilon)T(r, f)}$$

uniformly in  $\theta$  as  $z = re^{i\theta} \rightarrow \infty$  outside an  $\mathcal{E}$ -set, where  $\delta = \min_{0 \leq j \leq n-1} \delta_j(\mathfrak{A}) > 0$ .

We take  $\varepsilon < \delta/3$  and a path  $L: z = z(r) = re^{i\theta}$  ( $r_0 < r < \infty$ ) such that (21) and (22) hold on  $L^4$ , and set

$$d_1(r) = e^{(-\delta + \varepsilon)T(r, f)}$$

$$d_2(r) = e^{-\varepsilon T(r, f)}.$$

We consider on  $L$  the following equation

$$A_n(z)f^{*k-n} + A_{n+1}(z)f^{*k-n-1} + \dots + A_k(z) = 0.$$

Recall (21). Then we see from Lemma 6 that the roots  $f_i^*(z)$  ( $i = 1, 2, \dots, k-n$ ) lie outside  $K(d_2(r))$ . The equation  $F(z, f) = 0$  has  $k-n$  roots, say  $f_i(z)$  ( $i = 1, 2, \dots, k-n$ ), such that

$$[f_i^*(z), f_i(z)] < d_1(r),$$

because of the comment given just after Lemma 4 and (22). Thus the values  $f_i(z)$  ( $i = 1, 2, \dots, k-n$ ) lie outside  $K(d_2(r) - d_1(r))$ . On the other

<sup>4</sup>) We can find such a path  $L$  because (21) and (22) hold as  $z \rightarrow \infty$  outside an  $\mathcal{E}$ -set.

hand, we see from Lemma 4 that the remainder  $f_j(z)$  ( $j = k - n + 1, \dots, k$ ) satisfies

$$[f_j(z), \infty] < d_1(r).$$

Since  $d_1(r)/d_2(r) = e^{(-\delta+2\varepsilon)T(r,f)} \rightarrow 0$  as  $r \rightarrow \infty$ , we see that  $K(d_1(r))$  is disjoint with the complement of  $K(d_2(r) - d_1(r))$  for every sufficiently large  $r \geq r_1$ , whence we can conclude that the branches  $f_j(z)$  ( $j = k - n + 1, \dots, k$ ) with  $f_j(z(r_1)) \in K(d_1(r_1))$  draw a curve  $f_j(z(t))$ ,  $t \geq r \geq r_1$ , in  $K(d_1(r))$ . In fact, if the curve  $f_j(z(t))$ ,  $t \geq r$  ( $\geq r_1$ ), invades the zone;  $\{w; d_2(r) - d_1(r) < [w, \infty] < d_1(r)\}$ , we have at least one point  $z^* = z(t^*)$ ,  $t^* > r$ , on the curve such that

$$\begin{aligned} f_j(z^*) &\notin K(d_1(t^*)), \\ f_j(z^*) &\notin \text{complement of } K(d_2(t^*) - d_1(t^*)), \end{aligned}$$

which contradicts the fact that any root of the equation  $F(z^*, f) = 0$  must be contained in  $K(d_1(t^*))$  or the complement of  $K(d_2(t^*) - d_1(t^*))$ . Since  $d_1(r) \rightarrow 0$  ( $r \rightarrow \infty$ ), we see that the branches  $f_j(z)$  tend to infinity as  $z \rightarrow \infty$  along  $L$ . Thus our theorem has been established.

#### REFERENCES

- [ 1 ] J.M. Anderson-J. Clunie: Slowly growing meromorphic functions. *Comment. Math. Helv.*, **40**, pp. 267–280 (1966).
- [ 2 ] J. Dufresnoy: Théorie nouvelle des familles complexes normales. *Ann. Éc. Norm. sup.*, pp. 1–44 (1944).
- [ 3 ] W.K. Hayman: Slowly growing integral and subharmonic functions. *Comment. Math. Helv.*, **34**, pp. 75–84 (1960).
- [ 4 ] H.L. Selberg: Algebroiden Funktionen und Umkehrfunktionen der Abelscher Integrale. *Avh. Norske Vid. Akad., Oslo*, **8**, pp. 1–74 (1934).
- [ 5 ] Y. Tumura: Sur les théorèmes de M. Valiron et les singularités transcendentes indirectement critiques. *Proc. Imp. Acad. Japan*, **17**, pp. 65–69 (1941).
- [ 6 ] G. Valiron: Sur la dérivée des fonctions algébroides. *Bull. Soc. Math. France*, **59**, pp. 17–39 (1931).
- [ 7 ] —————: Sur le nombre des singularités transcendentes des fonctions inverses d'une classe d'algébroides. *C.R. Acad. Sci. Paris*, **200**, pp. 713–715 (1935).
- [ 8 ] —————: *Lectures on the general theory of integral functions*. Chelsea, (1949).
- [ 9 ] —————: Sur les valeurs déficientes des fonctions algébroides méromorphes d'ordre nul. *J.d'analyse Math.* **1**, pp. 28–42 (1951).

*Mathematical Institute  
Nagoya University*