
CORRIGENDUM

Adolescents from upper middle class communities: Substance misuse and addiction across early adulthood—CORRIGENDUM

SUNIYA S. LUTHAR, PHILLIP J. SMALL, AND LUCIA CICIOILLA

Arizona State University

doi: 10.1017/S0954579417000645, published by Cambridge University Press, 31 May 2017

Table 5 in the original online article contained labeling errors in the 2nd, 4th, 5th, 9th, and 10th columns. The entire cor-

rected table is reprinted herein. We regret these errors and any problems they may have caused.

Reference

Luthar, S. S., Small, P. J., & Cicicolla, L. (2017). Adolescents from upper middle class communities: Substance misuse and addiction across early

adulthood. *Development and Psychopathology*. Advance online publication. doi:10.1017/S0954579417000037

Address correspondence and reprint requests to: Suniya S. Luthar, Department of Psychology, Arizona State University, 950 South McAllister Avenue, Tempe, AZ 85287-1104; E-mail: suniya.luthar@asu.edu.

Table 5. Lifetime DSM-IV substance use diagnoses in the NESSY-Y (age 22 years) and NESSY-O (age 26 years) cohorts versus NCS-R for participants aged 22 years and 26 years

		Women					Men				
		NESSY ^a % (n)	NCS-R ^a % (n)	Ratio	<i>z</i>	NESSY ^b % (n)	NESSY ^a % (n)	NCS-R % (n)	Ratio	<i>z</i>	NESSY ^b % (n)
Any substance dependence	NESSY-Y	11.4 (9)	12.0 (9)	0.95	-0.12	15.8 (15)	19.2 (14)	8.1 (5)	2.37	2.04*	26.9 (25)
	NESSY-O	18.5 (15)	5.6 (5)	3.30	2.68**	24.2 (23)	22.6 (21)	11.0 (9)	2.05	1.94*	40.0 (44)
Any substance abuse	NESSY-Y	10.1 (8)	21.0 (17)	0.48	-1.94*	15.8 (15)	17.8 (13)	23.7 (14)	0.75	-0.91	30.1 (28)
	NESSY-O	22.2 (18)	15.9 (13)	1.40	1.07	30.5 (29)	26.9 (25)	20.2 (14)	1.33	0.99	51.8 (57)
Alcohol dependence	NESSY-Y	8.9 (7)	9.9 (8)	0.89	-0.22	12.6 (12)	13.7 (10)	8.1 (5)	1.69	1.13	19.4 (18)
	NESSY-O	14.8 (12)	4.1 (4)	3.60	2.48**	22.1 (21)	17.4 (16)	10.9 (9)	1.60	1.16	31.8 (35)
Alcohol abuse	NESSY-Y	7.6 (6)	18.2 (15)	0.41	-2.04**	12.6 (12)	15.1 (11)	20.5 (12)	0.74	-0.88	25.8 (24)
	NESSY-O	18.5 (15)	12.7 (11)	1.46	1.07	27.4 (26)	20.4 (19)	20.2 (14)	1.01	0.03	46.4 (51)
Drug dependence	NESSY-Y	2.5 (2)	5.7 (4)	0.44	-1.01	4.2 (4)	9.7 (7)	1.8 (1)	5.39	2.17*	16.1 (15)
	NESSY-O	7.4 (7)	2.3 (2)	3.22	1.61*	8.4 (8)	8.6 (8)	4.8 (4)	1.79	0.94	19.1 (21)
Drug abuse	NESSY-Y	3.8 (3)	11.2 (10)	0.34	-1.80*	6.3 (6)	9.7 (7)	10.3 (7)	0.94	-0.12	21.5 (20)
	NESSY-O	11.1 (9)	12.9 (10)	0.86	-0.38	12.6 (12)	11.8 (11)	18.5 (12)	0.64	-1.19	26.4 (29)

Note: NESSY-Y, New England Study of Suburban Youth younger cohort; NESSY-O, NESSY older cohort; NCS-R, National Comorbidity Survey Replication. Sampling weights were used to calculate percentages for NCS-R data according to Kessler et al. (2004). NCS-R age 22 women: *n* = 95, men: *n* = 84; NCS-R age 26 women: *n* = 96, men: *n* = 71. Comparisons of population proportions using *z* scores were weighted for sample size.

^aNESSY rates based on final assessment point only: ages 22 and 27 for NESSY-Y and NESSY-O, respectively. Ratio = %NESSY/%MTF. NESSY-Y: *n* = 78 women, *n* = 73 men; NESSY-O: *n* = 82 women, *n* = 89 men.

^bNESSY rates based on all annual assessments. Cumulative NESSY-O: *n* = 95 women, *n* = 110 men; cumulative NESSY-Y: *n* = 95 women, *n* = 93 men.

z* ≥ 1.6, *p* < .05. *z* ≥ 2.5, *p* < .01.