

Methods: We conducted a cross-sectional comparative experimental study. Subjects were 50 patients exhibiting behavioral disorders aged 9 to 18 years, with a diagnosis of attention-deficit/hyperactivity disorder (ADHD), oppositional defiant disorder (ODD), and conduct disorder (CD). Participants were assigned to ECBT and control groups. ECBT group contributed in 12-h weekly sessions within ECBT-inspired program. The control group received standard care. To assess emotion-related competencies, children were administered the emotion regulation questionnaire (ERQ-CA) and the alexithymia questionnaire for children (AQC). Parents completed the Child Behavior Checklist (CBCL) to measure youth externalizing problems. Tests were administered in pre- and post-test to all subjects. Both groups were matched for age, sex and educational level. Comparison of pre- and post-test results was performed using the Student's t-test.

Results: ECBT demonstrated a significant difference in the reduction of behavioral problems. ECBT effectively increased adaptive emotion regulation strategies (cognitive reappraisal) in the post-test. ECBT also reduced alexithymia scores, particularly difficulty identifying feelings, and externally oriented thinking.

Conclusions: ECBT demonstrated promising initial effectiveness in addressing emotion regulation deficits of children with externalizing behaviors

Disclosure: No significant relationships.

Keywords: externalizing behaviors; Children; emotion regulation

EPV0214

Descriptive analysis of patients admitted to a new adolescent inpatient unit in Madrid

M. Taracena Cuerda*, M. Esperesate Pajares, M. Feito Garcia, C. Arranz Martin, E. Sánchez Sampedro, A.M. Jiménez Bidón, R. Puente García and C. Pastor Jordá

Hospital 12 de Octubre, Psychiatry, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1125

Introduction: Adolescent mental health problems may have increased after COVID-19 worldwide pandemic. Therefore it seems necessary to study the state of mental health inpatient adolescent units.

Objectives: Adolescent mental health problems may have increased after COVID-19 worldwide pandemic. Therefore it seems necessary to study the state of mental health inpatient adolescent units.

Methods: An observational and descriptive analysis of the sample of patients between 12 and 17 years-old, that were admitted to the inpatient mental health unit since its opening on April 2021.

Results: A total of 205 patients were admitted from April 2021 until October 2021. We have observed sex differences within patients admitted, as the 82.9% of them were female. The mean age was 14.7, being 14.6 for girls and 15.3 for boys. The most common reason for admission (RFA) were suicidal ideation/attempt, eating disorders, affective disorders, conduct disorders/challenging behaviors and psychosis. Suicidal ideation/attempt was the most common RFA (57.07%) in both sexes, being higher among females (60.3%) than males (42.9%). Eating disorders were the second most common RFA in girls (17.7%) while psychosis (17.1%) and mood disorders (17.1%) were the second most common RFA within boys.

Conclusions: Findings on how COVID-19 affected adolescents mental health are controversial in the literature, our data suggest

that there is a need of developing quality studies that analyse how the pandemic might be influencing adolescents suicidal ideation/attempt and its protective and risk factors.

Disclosure: No significant relationships.

Keywords: Adolescents; inpatient unit; suicidal

EPV0215

Tension-Type Headache in Early Adolescents: Exploring the Predictive Role of Anxiety and Alexithymia

F. Bizzi¹, S. Charpentier Mora¹, A. Riva², M. Tironi^{1*} and R. Nacinovich²

¹University of Genoa, Department Of Educational Sciences (disfor), Genoa, Italy and ²San Gerardo Hospital, Child Neuropsychiatry Clinic, Monza, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1126

Introduction: Primary Headache, including Tension-Type Headache (TTH), represents one of the most common somatic disorders in children and adolescents with a strong impact on quality of life. Several risk factors, as environmental, familiar, and psychological features, including personality traits, are related to the development of Primary Headache. However, studies on specific subgroups of TTH are relatively few in early adolescents.

Objectives: Therefore, this cross-sectional pilot study aims at exploring the role of anxiety and alexithymia in early adolescents with and without TTH.

Methods: A sample of 70 early adolescents ($M_{age}=14.59$, $SD=1.85$; 71% females) consisting of a clinical group (31 with TTH) enrolled in an Italian Child Neuropsychiatry Clinic and a comparison group (38 without TTH) enrolled in schools, matched on gender and age, completed: 1) Multidimensional Anxiety Scale for Children (MASC) to detect the Total levels of Anxiety, also in their factor of Physical Symptoms, Social Anxiety, Harm Avoidance, and Separation Anxiety; 2) the Toronto Alexithymia Scale (TAS-20) to detect the Total levels of Alexithymia, also in their factor of Difficulty to Identifying and to Describing Feelings and Externally Oriented Thinking.

Results: TTH outcome positively correlated with Harm Avoidance ($\rho=.68$, $p<.001$) and Total Alexithymia ($\rho=.72$, $p<.001$). In a logistic regression, Harm Avoidance and Total Alexithymia predicted 69% of the variance in TTH outcome ($p<.032$).

Conclusions: This disorder may be a maladaptive strategy to cope with problems and feeling emotions, then early adolescents could be fostered in the acquisition of more adaptive emotion regulation abilities.

Disclosure: No significant relationships.

Keywords: Tension-Type Headache; Early Adolescents; alexithymia; Anxiety

EPV0217

Psychoeducation of Parents of Children with Autism Spectrum Disorders

M. Ivanov^{1*}, O. Bogacheva¹ and N. Simashkova^{1,2}

¹Federal State Budgetary Scientific Institution, Department Of Child Psychiatry, Moscow, Russian Federation and ²Mental Health Research