

Comparison of BMI, triponderal mass index, and paediatric body adiposity Index for predicting body fat and screening obesity in preschool children

Yimin Wang^{1 2}; Ke Xu^{1 2}; Miyuan Wang^{1 2}; Paiziyetia^{1 2}; Wenli Dong^{1 2};
Mengna Wei^{1 2}; Yanfen Jiang^{1 2}; Wenqi Xia^{1 2} Jiameng Zhou^{1 2}; Jianduan
Zhang^{1 2}

¹School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, China

²Key Laboratory of Environment and Health, Ministry of Education and Ministry of Environmental Protection, State Key Laboratory of Environmental Health (incubating), School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, 13 Hangkong Road, Wuhan, Hubei, 430030, People's Republic of China.

Address correspondence to: Jianduan Zhang, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, China [jd_zh@hust.edu.cn], 027-83657956.

Short version of the title: Comparing BMI, TMI, and BAIP for Predicting Body Fat in Preschool Children



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114524002046

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Several novel anthropometric indices, including paediatric body adiposity index (BAIp) and triponderal mass index (TMI), have emerged as potential tools for estimating body fat in preschool children. However, their comparative validity and accuracy, particularly when compared to established indicators such as Body Mass Index (BMI), have not been thoroughly investigated. This cross-sectional study enrolled 2869 preschoolers aged 3-6 years in Wuhan, China. The nonparametric Bland–Altman analysis was employed to evaluate the agreement between BMI, BAIp, and TMI with percentage of body fat (PBF), determined by bioimpedance analysis (BIA), serving as the reference measure of adiposity. Additionally, Receiver Operating Characteristic Curve (ROC) analysis was conducted to assess the effectiveness of BMI, BAIp, and TMI in screening for obesity. BAIp demonstrated the least bias in estimating PBF, showing discrepancies of 3.64% (95%CI: 3.40% to 4.12%) in boys and 3.95% (95%CI: 3.79% to 4.23%) in girls. Conversely, BMI underestimated PBF by 3.89% (95%CI: 3.70% to 4.37%) in boys and 4.81% (95%CI: 4.59% to 5.09%) in girls, while TMI also underestimated PBF by 5.15% (95%CI: 4.90% to 5.52%) in boys and 5.68% (95%CI: 5.30% to 5.91%) in girls. BAIp exhibited the highest area under the curve (AUC) values (AUC=0.867-0.996) in boys, whereas in girls, there was no statistically significant difference between BMI (AUC = 0.936, 95% CI: 0.921-0.948) and BAIp (AUC = 0.901, 95% CI 0.883-0.916) in girls (P=0.054). In summary, when considering the identification of obesity, BAIp shows promise as a screening tool for both boys and girls.

Abbreviations

TMI Triponderal Mass Index

BIA Bioelectrical impedance analysis

BAIp Paediatric Body Adiposity Index

HC Hip circumference

BMI Body Mass Index

PBF Percentage of body fat

DEXA Dual-energy X-ray absorptiometry

GAMLSS Generalized additive model

LMS Lambda-Mu-Sigma

LoA limits of agreement

ROC Receiver Operating Characteristic Curve

AUC Area under the curve

INTRODUCTION

Childhood obesity is a global concern, with approximately 39 million children under five being overweight or obese worldwide In 2020⁽¹⁾. China has seen a significant rise in preschool obesity, increasing from 3.1% in 2013 to 10.4% in 2020⁽²⁾. Childhood obesity often persists into adulthood, elevating the risk of chronic illnesses and premature death⁽³⁾. Early detection of childhood obesity is vital for preventing future health issues⁽⁴⁾, necessitating accurate, user-friendly, and cost-effective assessment tools.

For decades, BMI has widely served as a valuable tool for tracking obesity prevalence⁽⁵⁾. It has also linked obesity status with an increased risk for cardiovascular disease⁽⁶⁾, type 2 diabetes⁽⁷⁾, and mortality⁽⁸⁾. However, as an indirect measure of body fat mass, Body Mass Index (BMI)'s association with body fatness is not entirely accurate⁽⁹⁾. Besides, BMI does not offer insight into body fat distribution, a crucial aspect of health assessment⁽⁹⁾. For example, individuals with the same BMI but higher proportions of visceral adipose tissue and ectopic fat depots face an elevated risk of cardiovascular disease^(10, 11). The Triponderal Mass Index (TMI), recently introduced by Peterson et al., offers a potentially more accurate alternative, calculated as weight (kg)/height (m³)⁽¹²⁾. It has been suggested that this measurement may be a more effective predictor of percent body fat and metabolic syndrome than BMI⁽¹³⁾. Moreover, TMI is observed to be fairly consistent throughout childhood and adolescence⁽¹⁴⁾. Therefore, utilizing a single cutoff value for TMI can be a practical and convenient approach⁽¹²⁾. However, it has remained unclear whether TMI is superior to BMI in predicting body fat and identifying obesity in preschool children. An alternative measurement, the Body Adiposity Index (BAI), has also been proposed. It is calculated as hip circumference (HC) in centimeters divided by height in meters (HM) to the 1.5th power, minus 18⁽¹⁵⁾ ($BAI = HC / (HM)^{1.5} - 18$), primarily for adults. Some studies have suggested that BAI outperformed BMI in estimating body fat percentage in young adults⁽¹⁶⁾. However, inconsistent results have been found in Chinese children and adolescents, with BMI often being considered a better tool for estimating whole-body and central body fat⁽¹⁷⁾. Building on the concept of BAI, the Paediatric Body Adiposity Index (BAIp) has recently been developed specifically as a screening tool for childhood obesity^(18, 19), calculated as HC divided by height in HM to the 0.8th power, minus 38 ($BAIp = HC / (HM)^{0.8} - 38$)⁽²⁰⁾. However, if BAIp is suitable for the

Chinese population remains unexamined. Furthermore, although previous research has shown promise for BAIp in epidemiological research, Filgueiras M et al., have raised concerns about its accuracy compared to body fat⁽²¹⁾. Therefore, it remains unclear which index is the optimal one for evaluating body composition, necessitating a comparative analysis of the reliability of each indicator.

Understanding the relationships between adiposity markers and body fat is essential for enhancing the clinical identification of childhood obesity, guiding research efforts to comprehend its association with related diseases, and developing precise interventions. Therefore, this investigation aims to verify the validity of BMI, BAIp, and TMI in predicting the percentage of body fat(PBF) and screening obesity in a sample of Chinese preschool children, using bioelectrical impedance analysis (BIA) as the reference method.

METHODS

Participants

Children aged 3 to 6 years old in Wuhan were enrolled in this study. Wuhan was chosen due to its demographic mix and urban characteristics. The data were obtained from two cross-sectional surveys as part of the Wuhan Healthy Start Project for Preschool Children. To optimize resources and ensure regional diversity, a total of 35 kindergartens from Jiangnan and Hanyang (two representative districts in Wuhan) were included in this study. The first survey was carried out from 2021 to 2022, utilizing a cluster random sampling method to select all children from nine kindergartens in the Jiangnan District of Wuhan for investigation. The second survey was conducted in 2023, and 30 kindergartens were randomly selected by cluster sampling from Jiangnan and Hanyang districts, with all children in these kindergartens included in the survey (4 kindergartens overlapped with the previous survey, but with no duplication of participants). Initially, 3227 children were surveyed. After excluding cases of parental refusal and children's absence due to illness or other factors, a final sample of 2869 children aged 3-6 years was included for analysis (Figure 1).

In our study, informed consent was obtained from legal guardians of all participating children. We adhered to ethical standards by providing detailed information about the study objectives, procedures, potential risks, and benefits to the participants or their guardians. Consent forms

were written in clear and understandable language, ensuring that the guardians had adequate time to review the information and ask questions before providing consent.

Anthropometric index

Children's height, weight, waist circumference, and hip circumference were collected by uniformly trained researchers with a standard procedure. Kangwa WS-RT-2U(Wuhan, China) physical examination instrument and non-elastic tape measure were employed for measurements. To reduce the measurement error, all indicators were measured twice. If the differences between the two consecutive values exceeded 5mm for height or 100 g for weight, a third measurement was conducted until the difference fell within these thresholds. The final value used was the average of the two closest measurements. BMI, Paediatric Body Adiposity Index (BAIp), and Triponder Mass Index (TMI, kg/m³) were calculated as: $BMI(kg/m^2) = \text{Weight (kg)} / [\text{Height (m)}]^2$; $BAIp(0.01 m^{-0.5}) = \text{Hip Circumference} / (\text{Height})^{0.8} - 38$; $TMI(kg/m^3) = \text{Weight(kg)} / [\text{Height (m)}]^3$.

Assessment of PBF using BIA

Body fat was determined by multifrequency BIA using an InBody 230 analyzer (Inbody230 system, InBody Corp, Seoul, Korea), with tetra-polar 8-point tactile electrodes. Measurements were taken at two different frequencies (20 and 100 kHz) on each segment (right arm, left arm, trunk, right leg, and left leg), with participants bearing feet and wearing light clothing. The measurement was taken two hours after a meal, and children were instructed not to exercise excessively before the measurement to minimize the possible influence on the BIA values. During the measurement, children stood on the device for weight measurement, and then their identification number, age, sex, and height were entered into the device for PBF calculation. Impedance measurements were obtained with the children standing still while holding hand grips that were slightly abducted. The device then used the manufacturer's algorithm to calculate and output data, including fat mass, fat-free mass, and PBF .

Definition of obesity

Currently, there is no widely recognized cut-off value for preschool children's PBF to determine obesity. The study by Williams et al.⁽²²⁾ suggests using 25% and 30% as cutoff values to define obesity in boys and girls, respectively. However, preschool children

experience rapid growth and development, leading to significant fluctuation in their body composition with age. Using the same cut-off value across different age groups may compromise diagnostic accuracy⁽²³⁾. Therefore, many studies have adopted the 95th percentile of PBF (P_{95}) as the diagnostic criterion for obesity. For example, Mi et al. used P_{95} of PBF as the diagnostic standard for obesity based on Dual-energy X-ray absorptiometry (DEXA) measurement data⁽²⁴⁾. However, this standard cannot be widely applied, including in our study, due to the difficulty in obtaining DEXA data, especially for epidemiology study. Previous studies have shown that BIA and DEXA methods can be used interchangeably at the population level⁽²⁵⁾. Therefore, in this study, we established the reference values of PBF based on BIA data from our study population, defining $PBF > 95$ th percentile as obesity.

Statistical analysis

MedCalc version 20.027 and R 4.1.5 were used for statistical analysis, with a p-value < 0.05 considered statistically significant. Kolmogorov–Smirnov tests were applied to check for the assumption of normal distribution ($P < 0.05$). Continuous variables are presented as median (P_{25} , P_{75}), and group comparisons were conducted with non-parametric test. Generalized additive model (GAMLSS) was used to establish percentile reference curves of PBF in children. GAMLSS is an extension of the Lambda-Mu-Sigma (LMS) method, which uses four distribution parameters: median, coefficient of variation, skewness, and kurtosis. The P_5 , P_{10} , P_{25} , P_{50} , P_{75} , P_{80} , P_{85} , and P_{95} of sex - and age-specific PBF were estimated, and the percentile curves were drawn. Gamlss package in R 4.1.2 was used for analysis. The nonparametric Bland-Altman plots were used to assess the agreement between BMI, BAIP, and TMI with standard body fat measure. The studentized bootstrap method were utilized to establish the limits of agreement (LoA), defined as the 2.5th and 97.5th percentiles of differences. The receiver operating characteristic curve (ROC) was used to evaluate the performance of BMI, BAIP, and TMI in correctly classifying children as obesity. The P_{95} of PBF was established as the standard reference value of obesity, and MedCalc 20.027 software was used to draw and analyze the ROC curve and calculate the area under the curve (AUC). To ensure data accuracy, the dataset was refined by applying specific exclusion criteria. Entries with a variance of more than 25 units between each index (BMI, BAIP, and TMI) and PBF were deemed unsuitable and subsequently removed.

RESULTS

Characteristics of subjects

The 2869 participants consist of 1471 (51.27%) boys and 1398 (48.73%) girls, with means ages of 4.33 years and 4.32, respectively. Boys generally exhibit greater height, weight, WC, HC, BMI, and TMI than girls. Conversely, girls typically present a higher PBF ($P < 0.05$). There is no sex difference in age and BAIP. Demographic information is presented in Table 1.

Percentile reference value of PBF in children of different ages and sexes

The reference values of percentile PBF (P_5 , P_{10} , P_{25} , P_{50} , P_{75} , P_{85} , P_{90} , P_{95}) of children of different ages and sexes are shown in Table 2. The PBF percentile curves for boys and girls are shown in Figure 2. The PBF of boys and girls showed a steady decline with the increase of age. However, a significant decline in PBF was evident among boys aged 3 to 6, compared to girls.

Nonparametric Bland-Altman analysis of the differences between BMI, BAIP, and TMI with PBF.

The nonparametric Bland-Altman plot in Figure 3 depicts the concordance and median of the differences of BMI, BAIP, and TMI with PBF. Among the three measures, BAIP demonstrated the least mean bias in both boys 3.64% (95%CI: 3.40% to 4.12%) and girls 3.95% (95%CI: 3.79% to 4.23%) regarding PBF. This indicates that BAIP underestimated PBF by 3.64% in boys and 3.95% in girls. In contrast, BMI and TMI showed more significant bias. BMI underestimated PBF by 3.89% (95%CI: 3.70% to 4.37%) in boys and 4.81% (95%CI: 4.59% to 5.09%) in girls. Similarly, TMI underestimated PBF by 5.15% (95%CI: 4.90% to 5.52%) and 5.68% (95%CI: 5.30% to 5.91%) for boys and girls, respectively. The median of differences for BMI, BAIP, and TMI is displayed in Table 3.

ROC curves for obesity screening using BMI, BAIP, and TMI

We evaluated the effectiveness of BMI, BAIP, and TMI in identifying obesity by plotting ROC curves for the three indexes (Figure 4) and calculating corresponding AUC values, optimal cut-off values, specificity, and sensitivity (Table 4). Pairwise comparisons of the AUCs for these indexes were performed to identify the optimal indices for obesity screening. The AUCs of the indexes were all greater than 0.8, indicating acceptable accuracy and

predictive ability for obesity screening. In boys, BAIp demonstrated the highest AUC value (AUC=0.950, 95% CI: 0.937-0.961) among all indexes, followed by BMI (AUC=0.875, 95% CI: 0.857-0.892), and TMI (AUC=0.799, 95% CI: 0.777-0.819). Both of BMI and TMI has a significantly smaller AUC than BMI ($P < 0.05$). In girls, BMI had the highest ability to recognize obesity (AUC = 0.936, 95% CI: 0.921-0.948), followed by BAIp (AUC = 0.901, 95% CI: 0.883-0.916), and TMI (AUC = 0.866, 95% CI: 0.846-0.883). However, there was no statistically significant difference between BMI and BAIp in girls ($P=0.054$).

DISCUSSION

The present study evaluated the efficacy of BMI, BAIp, and TMI as screening tools for obesity and predicting PBF among preschool children in China. The nonparametric Bland–Altman plots showed that BAIp had better agreement with PBF than BMI and TMI in both boys and girls. In terms of obesity screening, BAIp has demonstrated higher accuracy values and greater effectiveness among boys as compared to BMI and TMI. However, for girls, the AUC value of BMI appeared to be higher than that of BAIp; however, this difference did not reach statistical significance.

This study used the GAMLSS method to fit the percentile reference values of PBF for preschool children aged 3-6 years in Wuhan. From the age of 3 to 6 years, the PBF showed a downward trend with age in both boys and girls and the decline of PBF is generally more noticeable in boys compared to girls. Previous research observed that children experience a peak in their fat content within the first 9-12 months after birth, owing to their energy requirements for growth. This fat content gradually decreases due to the development of their body structure and metabolism, reaching its minimum between the ages of 3-8 years old. After this age, fat content starts to increase again, a phenomenon known as adiposity rebound⁽²⁶⁾. Our findings indicate that older children have a lower PBF than 3-year-old children, signifying that preschool children are approaching the lowest level of PBF, which occurs before the onset of adiposity rebound. This trend is consistent with the findings of Dong et al. who measured the PBF of Chinese children using DEXA⁽²⁴⁾. In addition, our study also found sex differences in the PBF across preschool years. Specifically, at the age of 3 years, boys had a higher body fat percentage than girls, while at the age of 4 to 6 years, girls had a higher body fat percentage than boys. This finding is consistent with the studies

conducted by Zhao⁽²⁷⁾ and Kiumars⁽²⁸⁾ which also found differences in PBF between boys and girls aged 3-17 years. However, the factors that contribute to the differences in PBF between boys and girls in preschool age have yet to be fully clarified. While it is believed that sex steroids, leptin, and insulin-like growth factor I⁽²⁸⁾ may be involved, further scientific inquiry is necessary to better understand the complex interplay of these factors.

The result of the nonparametric Bland-Altman plot indicated that BAIP had better agreement with PBF than BMI and TMI in both boys and girls. Previous study have indicated that BMI has only a fair correlation with body fat⁽²⁹⁾, and Vanderwall et al. reported that in children under 9 years, BMIz demonstrate a weak to moderate predictive effect for both total fat mass and PBF⁽³⁰⁾. Compared to BMI, BAI gives more consideration to the characteristics of body shape and fat distribution, rather than solely focusing on the proportion between weight and height⁽³¹⁾. According to Aarbaoui, the disparity in PBF assessed by BIA compared to the estimation derived from BAIP did not exhibit statistical significance nor practical clinical relevance. Therefore, utilizing BAIP could be a simpler and more reliable method for determining PBF in paediatric populations⁽²⁰⁾. However, inconsistent with studies involving school-age children⁽³²⁾ and adolescents⁽³³⁾, TMI did not demonstrate superior performance in predicting PBF in this research. This might be attributed to the younger age of the participants, as the predictive ability of TMI for PBF may be influenced by children's growth and developmental characteristics⁽³⁴⁾. Our results demonstrated that the BAIP proves to be a more reliable indicator of obesity in boys compared to BMI and TMI, whereas BMI has similar screening performances to BAIP in girls. This contrasts with findings from Yu, who conducted a cross-sectional study in the Chinese population aged 6–60 years and found that BMI was more highly correlated with PBF than BAI⁽¹⁷⁾. Similarly, Ye reported that BAI is not an ideal index for obesity screening and that it is less closely related to PBF than BMI and TMI in Chinese children and adolescents⁽³⁴⁾. The different conclusions may have arisen due to the utilization of BAIP rather than BAI as the assessment index for childhood obesity in our study. Aarbaoui indicated that BAI, which was developed using an adult sample, is not valid for children, as it may overestimate the PBF of young people⁽³⁴⁾. The observed difference in the effectiveness of BMI, BAIP, and TMI for obesity screening in preschool children between sexes could stem from physiological disparities in body fat distribution and

composition. In the case of girls, similar to BAIp, BMI also exhibited a favorable effect in obesity screening. This may be attributed to the observed tendency for girls to accumulate fat at a faster rate at this age. Therefore, the favorable performance of BMI in girls could potentially be linked to the higher PBF of girls at this particular age. Freedman indicated that the accuracy of BMI varies according to the degree of body fatness and BMI tends to be a reliable indicator of excess adiposity in relatively fat children⁽³⁵⁾.

As for the limitations of the study, firstly, we used the BIA method to measure body fat content rather than more accurate techniques such as DXA and CT, which might introduce measurement bias. Additionally, the cross-sectional nature of this study could have limited our ability to assess the accuracy of BMI, BAIp, and TMI in tracking changes in body fat over time. Furthermore, while our study focused on children aged 3-6 in Wuhan, the results may not be directly extrapolated to the entire population of children in that age group across China. Factors such as regional variations in lifestyle, diet, and socioeconomic status could influence the generalizability of our findings. Therefore, caution should be exercised when applying our results to the broader population of children aged 3-6 in China.

CONCLUSION

In conclusion, this study indicated that compared with BMI and TMI, BAIp showed better agreement with PBF. BAIp could potentially serve as a promising alternative screening tool for obesity in both boys and girls. This finding suggests that BAIp could serve as a valuable screening tool for identifying obesity risk with greater accuracy, thereby providing a reliable foundation for early intervention. Consequently, it is recommended that the public health sector consider incorporating BAIp into routine child health surveillance programs to enhance the precision of screening and the effectiveness of interventions.

Acknowledgements: We wish to thank all the participants and the healthcare staff in the kindergartens for making this critical study possible.

Author contributions: JDZ conceptualized and designed the study, supervised the overall study implementation, and reviewed and revised the manuscript. YMW drafted the initial manuscript, conducted the initial analyses, and revised the manuscript. KX conceptualized and designed the study, collected data. PZYT and WLD, data collection. MYW, MNW, YFJ, WQX and JMZ reviewed the manuscript for important intellectual content.

Ethical statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology (Project identification code: (2020) IEC (A179))

Funding: This study was funded by the National Natural Science Foundation of China (NSFC 82073567).

Declaration of competing interest: The authors declare that they have no competing interests.

Data availability: Data will be available upon request to the corresponding author.

References

1. Okunogbe A, Nugent R, Spencer G, et al. Economic impacts of overweight and obesity: current and future estimates for 161 countries. *BMJ global health*. 2022;7(9).
2. Zeng Q, Li N, Pan XF, et al. Clinical management and treatment of obesity in China. *The lancet Diabetes & endocrinology*. 2021;9(6):393-405.
3. Simmonds M, Llewellyn A, Owen CG, et al. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev*. 2016;17(2):95-107.
4. Smith JD, Fu E, Kobayashi MA. Prevention and Management of Childhood Obesity and Its Psychological and Health Comorbidities. *Annual review of clinical psychology*. 2020;16:351-78.
5. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017;390(10113):2627-42.
6. Attard SM, Herring AH, Howard AG, et al. Longitudinal trajectories of BMI and cardiovascular disease risk: the national longitudinal study of adolescent health. *Obesity (Silver Spring)*. 2013;21(11):2180-8.
7. Mi B, Wu C, Gao X, et al. Long-term BMI change trajectories in Chinese adults and its association with the hazard of type 2 diabetes: evidence from a 20-year China Health and Nutrition Survey. *BMJ open diabetes research & care*. 2020;8(1).
8. Global BMIMC, Di Angelantonio E, Bhupathiraju Sh N, et al. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet*. 2016;388(10046):776-86.
9. Bray GA. Beyond BMI. *Nutrients*. 2023;15(10).
10. Piché ME, Poirier P, Lemieux I, et al. Overview of Epidemiology and Contribution of Obesity and Body Fat Distribution to Cardiovascular Disease: An Update. *Progress in cardiovascular diseases*. 2018;61(2):103-13.
11. Abraham TM, Pedley A, Massaro JM, et al. Association between visceral and subcutaneous adipose depots and incident cardiovascular disease risk factors. *Circulation*. 2015;132(17):1639-47.

12. Peterson CM, Su H, Thomas DM, et al. Tri-Ponderal Mass Index vs Body Mass Index in Estimating Body Fat During Adolescence. *JAMA Pediatr.* 2017;171(7):629-36.
13. Wang X, Dong B, Ma J, et al. Role of tri-ponderal mass index in cardio-metabolic risk assessment in children and adolescents: compared with body mass index. *Int J Obes (Lond).* 2020;44(4):886-94.
14. Ashley-Martin J, Ensenaer R, Maguire B, et al. Predicting cardiometabolic markers in children using tri-ponderal mass index: a cross-sectional study. *Arch Dis Child.* 2019;104(6):577-82.
15. Bergman RN, Stefanovski D, Buchanan TA, et al. A better index of body adiposity. *Obesity (Silver Spring).* 2011;19(5):1083-9.
16. Fedewa MV, Nickerson BS, Esco MR. Associations of body adiposity index, waist circumference, and body mass index in young adults. *Clin Nutr.* 2019;38(2):715-20.
17. Yu Y, Wang L, Liu H, et al. Body mass index and waist circumference rather than body adiposity index are better surrogates for body adiposity in a Chinese population. *Nutr Clin Pract.* 2015;30(2):274-82.
18. Mahmoud I, Al-Wandi AS, Gharaibeh SS, et al. Concordances and correlations between anthropometric indices of obesity: a systematic review. *Public Health.* 2021;198:301-6.
19. Alves Junior CA, Mocellin MC, Gonçalves ECA, et al. Anthropometric Indicators as Body Fat Discriminators in Children and Adolescents: A Systematic Review and Meta-Analysis. *Adv Nutr.* 2017;8(5):718-27.
20. El Aarbaoui T, Samouda H, Zitouni D, et al. Does the body adiposity index (BAI) apply to paediatric populations? *Ann Hum Biol.* 2013;40(5):451-8.
21. De Santis Filgueiras M, Cecon RS, de Faria ER, et al. Agreement of body adiposity index (BAI) and paediatric body adiposity index (BAIp) in determining body fat in Brazilian children and adolescents. *Public Health Nutr.* 2019;22(1):132-9.
22. Williams DP, Going SB, Lohman TG, et al. Body fatness and risk for elevated blood pressure, total cholesterol, and serum lipoprotein ratios in children and adolescents. *Am J Public Health.* 1992;82(3):358-63.
23. Neovius M, Rasmussen F. Evaluation of BMI-based classification of adolescent overweight and obesity: choice of percentage body fat cutoffs exerts a large influence. *The*

COMPASS study. *Eur J Clin Nutr.* 2008;62(10):1201-7.

24. Dong H, Yan Y, Liu J, et al. Reference centiles for evaluating total body fat development and fat distribution by dual-energy x-ray absorptiometry among children and adolescents aged 3-18 years. *Clin Nutr.* 2021;40(3):1289-95.

25. Chen M, Liu J, Ma Y, et al. Association between Body Fat and Elevated Blood Pressure among Children and Adolescents Aged 7-17 Years: Using Dual-Energy X-ray Absorptiometry (DEXA) and Bioelectrical Impedance Analysis (BIA) from a Cross-Sectional Study in China. *Int J Environ Res Public Health.* 2021;18(17).

26. Rolland-Cachera MF, Deheeger M, Maillot M, et al. Early adiposity rebound: causes and consequences for obesity in children and adults. *Int J Obes (Lond).* 2006;30 Suppl 4:S11-7.

27. Zhao Y, Gong JX, Ji YT, et al. Cross-sectional study of characteristics of body composition of 24,845 children and adolescents aged 3-17 years in Suzhou. *BMC Pediatr.* 2023;23(1):358.

28. Arfai K, Pitukcheewanont PD, Goran MI, et al. Bone, muscle, and fat: sex-related differences in prepubertal children. *Radiology.* 2002;224(2):338-44.

29. Nevill AM, Duncan MJ, Myers T. BMI is dead; long live waist-circumference indices: But which index should we choose to predict cardio-metabolic risk? *Nutrition, Metabolism and Cardiovascular Diseases.* 2022;32(7):1642-50.

30. Vanderwall C, Randall Clark R, Eickhoff J, et al. BMI is a poor predictor of adiposity in young overweight and obese children. *BMC Pediatr.* 2017;17(1):135.

31. Feng J, He S, Chen X. Body Adiposity Index and Body Roundness Index in Identifying Insulin Resistance Among Adults Without Diabetes. *The American journal of the medical sciences.* 2019;357(2):116-23.

32. Niu Y, Zhang Y, Sheng J, et al. Tri-ponderal mass index as a screening tool for obesity prediction in children aged 6-9 years. *Front Endocrinol (Lausanne).* 2023;14:1277125.

33. Sun J, Yang R, Zhao M, et al. Tri-Ponderal Mass Index as a Screening Tool for Identifying Body Fat and Cardiovascular Risk Factors in Children and Adolescents: A Systematic Review. *Front Endocrinol (Lausanne).* 2021;12:694681.

34. Ye XF, Dong W, Tan LL, et al. Identification of the most appropriate existing anthropometric index for home-based obesity screening in children and adolescents. *Public*

Health. 2020;189:20-5.

35. Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness and risk among children. *Pediatrics*. 2009;124 Suppl 1:S23-34.

Conflict of Interest Disclosures (includes financial disclosures): The authors declare no competing interests.

Funding/Support: This study was funded by the National Natural Science Foundation of China (NSFC 82073567)

Ethical Approval:

We obtained ethical approval from the Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology.

Data Availability Statement:

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Authorship:

JDZ conceptualized and designed the study, and reviewed and revised the manuscript. YMW drafted the initial manuscript, conducted the initial analyses, and revised the manuscript. KX conceptualized and designed the study, collected data, and reviewed and revised the manuscript. PZYT, WLD coordinated and supervised data collection. MYW, MNW, WQX, JMZ and YFJ reviewed the manuscript for important intellectual content.

Table 1

Selected characteristics of the study population

Variables	Total(N=2817)	Boys (N=1444)	Girls (N=1373)	P-value
Age, years, (P ₂₅ ,P ₇₅)	4.31(3.90,5.04)	4.31(3.90,5.10)	4.31(3.89,5.00)	0.373
Age, years, n (%)				0.697
3	873 (31.97)	442 (30.57)	431 (31.39)	
4	1208 (42.89)	612 (42.39)	596 (43.41)	
5	572 (20.29)	306 (21.16)	266 (19.37)	
6	165 (5.85)	85 (5.88)	80 (5.83)	
Height, cm, (P ₂₅ ,P ₇₅)	106.55(102.30,111.70)	107.25(103.00,112.50)	105.65(101.40,111.20)	<0.001
Weight, kg, (P ₂₅ ,P ₇₅)	17.55(15.80,20.02)	17.98 (16.18,20.74)	17.10(15.51,19.24)	<0.001
WC, cm, (P ₂₅ ,P ₇₅)	50.70(48.65,53.25)	51.33(49.25,53.85)	50.05(48.10,52.60)	<0.001
HC, cm, (P ₂₅ ,P ₇₅)	57.00(54.10,60.30)	57.25(54.40,60.50)	56.75(53.90,60.05)	0.007
BMI, kg/m ² , (P ₂₅ ,P ₇₅)	15.50(14.66,16.40)	15.69(14.86,16.60)	15.30(14.48,16.20)	<0.001
BAIp, 0.01 m ^{-0.5} , (P ₂₅ ,P ₇₅)	16.11(14.13,18.21)	15.96 (14.10,18.18)	16.23(14.14,18.26)	0.397
TMI, kg/m ³ , (P ₂₅ ,P ₇₅)	14.55(13.66,15.43)	14.57(13.79,15.40)	14.50(13.53,15.47)	0.016
PBF, %, (P ₂₅ ,P ₇₅)	19.80(16.40,23.70)	19.60(16.30,23.50)	20.00(16.60,23.90)	0.041

Notes: WC waist circumference, HC hip circumference, BMI body mass index, BAIp pediatric body adiposity index, TMI triponderal mass index, PBF percentage body fat.

Table2. Percentiles for percentage of body fat (PBF) by age in boys and girls aged 3-6 years

Sex	Year	PBF(%)							
		5th	10th	25th	50th	75th	85th	90th	95th
Boys	3.0	13.28	14.84	17.65	21.18	25.38	28.05	30.08	33.50
	4.0	12.26	13.70	16.30	19.56	23.44	25.90	27.77	30.93
	5.0	11.19	12.51	14.88	17.85	21.40	23.65	25.36	28.23
	6.0	10.00	11.18	13.30	15.96	19.12	21.13	22.66	25.24
Girls	3.0	12.78	14.45	17.36	20.88	24.93	27.44	29.32	32.47
	4.0	12.27	13.88	16.67	20.05	23.94	26.35	28.15	31.17
	5.0	11.68	13.21	15.87	19.09	22.78	25.08	26.80	29.67
	6.0	10.93	12.36	14.84	17.85	21.31	23.46	25.07	27.75

Note: PBF: percentage of body fat.

Indexes	Median of the differences	Lower LoA	Upper LoA	p Value
Boys				
BMI	3.89(3.70,4.37)	-3.99(-4.32,-3.19)	14.80(13.48,15.40)	<0.001
BAIp	3.64(3.40,4.12)	-6.20(-6.87,-5.42)	15.38(13.76,16.11)	<0.001
TMI	5.15(4.90,5.52)	-3.49(-3.99,-2.98)	17.28(15.62,18.32)	<0.001
Girls				
BMI	4.81(4.59,5.09)	-3.35(-4.00,-2.92)	15.27(14.57,16.90)	<0.001
BAIp	3.95(3.79,4.23)	-2.85(-3.38,-2.59)	12.99(12.32,14.63)	<0.001
TMI	5.68(5.30,5.91)	-2.34(-3.18,-1.88)	16.84(16.02,18.87)	<0.001

Table3. Agreement, and proportional bias assessment between BMI, BAIp, and TMI for PBF.

Notes: BMI: body mass index; BAIp: pediatric body adiposity index; TMI: triponderal mass index; SD: standard deviation; LoA: limits of agreement.

TABLE 4

Comparison of the Receivers Operator Characteristic curves for various anthropometric indices in predicting obesity

Sex	Variable	AUC (95% CI)	Comparison of AUC				Sensitivity	Specificity	Optimal cut-offs
			BAIp		TMI				
			Z	P	Z	P			
Boys	BMI	0.875(0.857~0.892)	2.324	0.020	4.180	<0.001	77.3	93.0	17.776
	BAIp	0.950(0.937~0.961)			4.180	<0.001	92.0	93.3	20.410
	TMI	0.799(0.777~0.819)					73.3	88.7	16.005
Girls	BMI	0.936(0.921~0.948)	1.924	0.054	3.935	<0.001	86.2	90.5	16.924
	BAIp	0.901(0.883~0.916)			1.463	0.143	87.7	84.9	19.130
	TMI	0.866(0.846~0.883)					80.0	80.3	15.619

Notes: BMI: body mass index; BAIp: pediatric body adiposity index; TMI: triponderal mass index




Figure1. Flow chart showing inclusion and exclusion of children from Wuhan healthy start project.




Figure2. Percentile curves of percentage of body fat(%) for boys and girls aged 3 to 5 years




Figure 3. Nonparametric Bland-Altman plots comparing the agreement between PBF estimated by BMI, BAIp, and the TMI with PBF estimated by BIA in boy and girl.




Figure 4. ROC curves of BMI, BAIP, and TMI for screening obesity by sex. BMI, body mass index; BAIP, pediatric body adiposity index; TMI, tri-ponderal mass index