

Research

Cite this article: Amegbor PM, Kuuire VZ, Bisung E, Braimah JA. (2019) Modern or traditional health care? Understanding the role of insurance in health-seeking behaviours among older Ghanaians. *Primary Health Care Research & Development* 20(e71): 1–8. doi: [10.1017/S1463423619000197](https://doi.org/10.1017/S1463423619000197)

Received: 26 September 2018

Revised: 4 January 2019

Accepted: 24 February 2019


Key words:

health-seeking behaviour; health utilization; health insurance; ageing; health care access; Ghana

Author for Correspondence: Prince Michael Amegbor, Department of Geography and Planning, Queen's University, Mackintosh-Corry Hall, E208, Kingston, ON K7L 3N6, Canada. E-mail: 11pma4@queensu.ca

© The Author(s) 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Modern or traditional health care? Understanding the role of insurance in health-seeking behaviours among older Ghanaians

Prince M. Amegbor¹ , Vincent Z. Kuuire^{2,3}, Elijah Bisung⁴ and Joseph A. Braimah¹

¹Department of Geography and Planning, Queen's University, Kingston, ON, Canada; ²Department of Geography, University of Toronto Mississauga, Mississauga, ON, Canada; ³Social & Behavioural Health Sciences Division, Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada and ⁴School of Kinesiology and Health, Queen's University, Kingston, ON, Canada

Abstract

Aim: This paper examined the association between wealth and health insurance status and the use of traditional medicine (TM) among older persons in Ghana. **Background:** There have been considerable efforts by sub-Saharan African countries to improve access to primary health care services, partly through the implementation of risk-pooling community or national health insurance schemes. The use of TM, which is often not covered under these insurance schemes, remains common in many countries, including Ghana. Understanding how health insurance and wealth influence the use of TM, or otherwise, is essential to the development of equitable health care policies. **Methods:** The study used data from the first wave of the World Health Organisation's Study of Global Ageing and Adult Health conducted in Ghana in 2008. Descriptive statistics and negative loglog regression models were fitted to the data to examine the influence of insurance and wealth status on the use of TM, controlling for theoretically relevant factors. **Findings:** Seniors who had health insurance coverage were also 17% less likely to frequently seek treatment from a TM healer relative to the uninsured. For older persons in the poorest income quintile, the odds of frequently seeking treatment from TM increased by 61% when compared to those in the richest quintile. This figure was 46%, 62% and 40% for older persons in poorer, middle and richer income quintiles, respectively, compared to their counterparts in the richest income quintile. **Conclusion:** The findings indicate that TM was primarily used by the poor and persons who were not enrolled in the National Health Insurance Scheme. TM continues to be a vital health care resource for the poor and uninsured older adults in Ghana.

Introduction

There have been considerable efforts by sub-Saharan African (SSA) countries to improve access to primary health care services, partly through the implementation of risk-pooling community or national health insurance schemes (Jowett *et al.*, 2004; Witter & Garshong, 2009; Robyn *et al.*, 2013). Universal health insurance coverage programmes, such as Ghana's National Health Insurance Scheme (NHIS), are vital mechanisms for ensuring equity in access to and use of health care services, via the removal of financial barriers associated with seeking treatment in health care facilities (Bennett, 2004; Agyepong & Adjei, 2008; Blanchet *et al.*, 2012). Community-based and national health insurance schemes also offer additional funds for health care financing (Jowett *et al.*, 2004; Gnawali *et al.*, 2009). Yet, evidence from existing studies in the sub-region shows payment of health insurance premiums remains a significant challenge and impacts the utilization of health care services (Dixon *et al.*, 2011; Jehu-Appiah *et al.*, 2011; Aryeetey *et al.*, 2013). As a consequence, direct out-of-pocket payment for health care services still remains a predominant practice (McIntyre *et al.*, 2006; Gnawali *et al.*, 2009; Saksena *et al.*, 2011; Kusi *et al.*, 2015). In Ghana, this expenditure accounts for nearly half of the total expenditure on health care (Akazili *et al.*, 2012). At the household level, inability to pay for insurance premiums excludes many people, particularly those with low socioeconomic status, from enrolling in insurance schemes (Dixon *et al.*, 2011; Jehu-Appiah *et al.*, 2011; Kusi *et al.*, 2015). Thus, access to basic health care services in the formal health care system still remains a significant challenge for many poor and vulnerable populations in Ghana and SSA.

Despite the growing availability of modern or Western health care services, significant sections of populations in Ghana and other SSA countries continue to rely on indigenous healing practices (traditional medicine – TM) for their health care needs (Pouliot, 2011; Sato, 2012a; Thorsen and Pouliot, 2015). TM refers to healing knowledge, skills and practices based on the traditions, cultural beliefs and experiences indigenous to different communities or cultures (WHO, 2002). It comprises of the use of animal-based, plant-based and mineral-based

medicine, as well as, the use of spiritual therapies in the prevention, diagnosis, improvement and treatment of physical and psychosocial illness (World Health Organisation, 2013). Traditional medical practitioners include herbalist, bone setters, faith healers and spiritualist (Amegbor 2014; World Health Organisation, 2013; Lakshmi *et al.*, 2015). Evidence from existing studies show that TM is effective in treating various illness and common health conditions such as malaria, cholera and typhoid, as well as complex health issues such as infertility (O'Brien *et al.*, 2012; Thorsen and Pouliot, 2015; Towns and Van Andel, 2016). Reports indicate that between 60% and 80% of the population in SSA rely on indigenous medical remedies, including herbal medicine for their primary health care needs (World Health Organisation, 2013; McFarlane, 2015; Thorsen and Pouliot, 2015).

The vital role of TM in health and health care delivery is particularly evident among rural residents, the poor and vulnerable populations (Aikins & Marks, 2007; Powell-Jackson *et al.*, 2014). Among these populations, barriers to Western health care services include unavailability and inadequate health care facilities and practitioners, cost of care and other sociocultural beliefs (Tabi *et al.*, 2006; Pouliot, 2011; Sato, 2012b; Kuuire *et al.*, 2015). On the contrary, TM is readily available, accessible and affordable for these segments of the population (Macha *et al.*, 2012; Sato, 2012b). Though TM use is often associated with basic health problems, studies in Ghana reveals its use is particularly higher among patients with multiple acute and chronic health conditions (Sato, 2012b; Aikins *et al.*, 2014). Likewise, studies report that patients with non-communicable diseases (NCDs), such as hypertension and diabetes, view Western services for their health care needs as expensive and inaccessible, thereby resorting to the use of TMs (Aikins *et al.*, 2014; Aikins *et al.*, 2015). The surge in NCDs in SSA coupled with the demand for TM has led to the emergence of quack TM healers whose practices pose a health danger to the poor and vulnerable that patronize their services (Abdullahi, 2011; Kigen *et al.*, 2013).

The objective of this paper is to examine the effects of wealth and health insurance status on the use of TM among older persons (persons aged 60 years and above) in Ghana. While there is a plethora of literature on the effect of wealth and health insurance on health care-seeking behaviour in Ghana (see Kuuire *et al.*, 2017; Jehu-Appiah *et al.*, 2011; Dixon *et al.*, 2011), our understanding of how these factors influence the decision around seeking Western or traditional health care remain limited among older people. Existing studies on health care-seeking behaviour among Ghana's older adult population focus on the effects of socioeconomic factors on NHIS enrolment (Kuiure *et al.*, 2017) and the impacts of NHIS status on health care utilization (Fenny *et al.*, 2016; Dalinjong *et al.*, 2017; van der Wielen *et al.*, 2018). Moreover, the discussions in these papers ignore the vital role TM play in health and health care-seeking behaviour of people, especially the poor and vulnerable in society – including the older persons. TM utilization is predominant among Ghanaians with chronic health conditions (Sato, 2012b; Aikins *et al.*, 2015), and research shows that older persons are more susceptible to chronic health conditions compared to younger persons (Aikins *et al.*, 2014; Prince *et al.*, 2015). Currently, the NHIS provides limited coverage for chronic conditions (such as hypertension) and does not cover treatment outside the biomedical care setting, including TM (Barimah, 2013; Aikins *et al.*, 2014).

Ghana has an estimated 1.6 million older persons (6.7% of the country's total population) and this is expected to quadruple to about 4.8 million in 2050 (8.9% of the country's estimated

population in 2050) (Amegbor *et al.*, 2018; Minicuci *et al.*, 2014; World Health Organisation, 2014). Understanding the health-seeking behaviour of this rapidly growing section of the population is useful for providing relevant policy direction that contributes to improved health and well-being of older persons. Importantly also, prevalence of NCDs is usually considerably higher among older persons (Aikins *et al.*, 2010; Amegbor *et al.*, 2018; Minicuci *et al.*, 2014), although such conditions are not currently covered under the NHIS and could potentially influence health-seeking behaviours, including the use of TM, due to associated cost of treatment. This context provides the impetus for examining how health insurance and wealth status impact the use of TM among older Ghanaians. Thus, controlling for other factors, the paper assumes that (1) individuals with health insurance coverage and (2) individuals in higher income/wealth categories will be less likely to use traditional healers as their most frequent source of treatment during illness.

Materials and methods

Data

Data for the study comes from the World Health Organisation's Study of Global Ageing and Adult Health (SAGE). SAGE has a longitudinal design and involves collecting data on the health needs of older persons in five countries including Ghana. Irrespective of the focus on persons who are 50 and older, a small sample of persons aged between 18 and 49 were also included for comparison. We used the first wave of the SAGE data collected in Ghana between January 2007 and December 2008 through face-to-face interviews with respondents who were selected via a stratified multi-stage sampling strategy. While the survey for Wave 2 has been completed, the data are not currently available, hence our decision to use data from Wave 1. The first stage involved stratifying the sample by the 10 administrative regions in Ghana and by rural/urban location. From this, primary sampling units consisting of 235 enumeration areas (EAs) were selected. All EAs that did not have residents who are 50 years or older were excluded. A total of 5269 households were surveyed with some 5573 individuals sampled from these households. An estimated response rate of 86% and 80% were achieved at the household and individual levels respectively. However, this study uses a sub-sample of individuals who are 60 years or older because this is the official retirement age in Ghana and the age from which individuals are categorized as older persons. The final analytical samples used in the study were 2256.

Measures

Dependent variable: The dependent variable (frequently used health care type) for this study was a measure of the type of health care frequently used by older persons during the last three years. We derived this variable from the following question: "Thinking about health care you needed in the last three years, where did you go most often when you felt sick or needed to consult someone about your health?" Response categories were as follows: private doctor's office, private clinic or health care facility, private hospital, public clinic or health care facility, public hospital, charity or church-run clinic, charity or church-run hospital, traditional healer, pharmacy or dispensary and other. Except respondents who used a traditional healer (recoded "1 = traditional medicine"), all responses were recoded as "0 = modern health facility".

Focal independent variables: Health insurance coverage, the focal independent variable in the study was obtained from the question ‘do [you] have health insurance coverage?’ with the following response options: ‘Yes, mandatory insurance’, ‘Yes, voluntary insurance’, ‘Yes, both mandatory and voluntary insurance’ and ‘No, none’. The responses were operationalized to indicate whether respondents had health insurance or not (coded 0 = uninsured and 1 = insured). The second focal independent variable in this study, wealth status, was measured by income quintile which is a household asset-based measure of wealth provided in the data. Income quintile was coded as follows: 0 = richest, 1 = richer, 2 = middle, 3 = poorer and 4 = poorest.

Other independent variables: We included theoretically relevant demographic and socioeconomic variables in the study based on existing literature on health-seeking behaviours. The demographic variables included in the study were age; sex (coded 1 = male, 2 = female); ethnicity (coded 1 = Akan, 2 = Ewe/Ga-Adangbe, 3 = Gruma/Mole-Dagbani, 4 = other ethnic group); religion (coded 1 = Christian, 2 = Muslim, 3 = Traditional, 4 = Other religion/none); education (coded 0 = no formal education, 1 = primary, 2 = secondary/higher); sector of current or sector of former occupation if currently retired (coded 1 = self-employed, 2 = public/private sector, 3 = informal sector); current employment status (coded 0 = unemployed/retired, 1 = employed); marital status (coded 1 = married, 2 = separated/divorced, 3 = widowed) and location of residence (coded 1 = urban, 2 = rural). We also included respondents’ self-rated health (coded 1 = good, 2 = moderate, 3 = bad), and whether they had ever been diagnosed with at least one NCD (hypertension, stroke, angina, diabetes and asthma) as control variables.

Analytical strategy

We employed descriptive statistics, bivariate and multivariate regression techniques in analysing the data. For the regression models, we adopted the negative loglog link function in estimating the relationship between the outcome and independent variables. The choice of this technique was informed by the skewed nature of the distribution between the categories of our outcome variable (modern health facility = 96.8%; TM = 3.2%). Unlike the logit link function, the negative loglog link function accounts for the specific problem of skewed distributions where lower categories of the binary outcome are more probable. This avoids the possibility of obtaining biased parameter estimates. In all bivariate and multivariate analyses, we imposed respondents’ identification number as a cluster variable due to the hierarchical nature of the data. This strategy prevents the violation of the independence assumption in standard regression models. Missing cases in the data were handled by using the multiple imputation technique. All analyses were undertaken following the completion of the multiple imputations in STATA 14. Analyses also accounted for the sampling weight provided in the data.

Results

Descriptive

From [Table 1](#), most respondents (~97%) often relied on private or public modern health facilities, while only a small section of respondents (~3%) used the services of TM practitioners. A little over half (~55%) of seniors in Ghana do not have health insurance coverage. Most respondents were Akan (~48%), Christian (~70%), had no formal education (~63%), were self-employed (~80%),

Table 1. Distribution of study variables (n = 2256)

	Per cent
‘Frequently used health care type’	
Modern facility	96.8
Traditional healer	3.2
Insurance status	
Uninsured	55.2
Insured	44.8
Income quintile	
Richest	21.2
Richer	19.7
Middle	21.6
Poorer	18.6
Poorest	18.9
Age	71.2 (mean)
Gender	
Male	49.2
Female	50.8
Ethnicity	
Akan	48.3
Ewe/Ga-Adangbe	18.2
Gruma/Mole Dagbani	7.9
Other	25.6
Religion	
Christian	70.2
Muslim	15.2
Traditional	8.9
Other/none	5.7
Education	
No formal education	63.2
Primary	17.5
Secondary or higher	19.3
Main activity	
Unemployed	41.7
Working	58.3
Employment sector	
Self-employed	80.0
Private/public sector	13.4
Informal sector	6.6
Marital status	
Married	54.2
Separated/divorced	10.6
Widowed	35.2
Location	
Rural	60.1

(Continued)

Table 1. (Continued)

	Per cent
Urban	39.9
NCD status	
No	62.7
Yes	37.3
Self-rated health	
Good	31.3
Moderate	45.4
Bad	23.3




Figure 1. Bivariate negative loglog regression of ‘frequently used health care type’ (n = 2256)

married (~54%) and resided in rural locations (~60%). Majority of seniors reported they do not have an NCD (~63%) and close to half reported their health status as moderate (~45%).

Bivariate

Figure 1 shows only variables that emerged as predictors of the outcome variable at the bivariate level. The results show that there is a statistically significant relationship between the two focal independent variables and the most frequently used type of health care. Seniors who are insured have a 2% decrease in the odds of seeking treatment from TM healers compared to their counterparts who are uninsured. Overall, seniors who belonged to lower income quintiles had higher odds of seeking TM. For example, compared to the richest, the poorest seniors had a 4% increase odds of frequently seeking TM. Similarly, compared to the richest, seniors who belonged to poorer and middle income quintiles had 2% and 5% increased odds of frequently using TM, respectively.

In addition to the two focal independent variables, two other independent variables were significantly associated with the type of health care (see Table 2). Respondents who indicated they were widowed had 2% decreased odds of frequently using TM compared to seniors who are married. Seniors who indicated they had one or more NCD had a 2% decrease in the odds of seeking TM compared to those who had no NCD.

Table 2. Bivariate negative loglog regression of ‘frequently used health care type’ (n = 2256)

	Coef.	(Std. Err.)
Insurance status (ref: uninsured)		
Insured	-0.02	(0.01) *
Income quintile (ref: richest)		
Richer	0.02	(0.01)
Middle	0.05	(0.01)**
Poorer	0.02	(0.01)*
Poorest	0.04	(0.01)**
Age		
	0.00	(0.00)
Gender (ref: male)		
Female	-0.01	(0.01)
Ethnicity (ref: Akan)		
Ewe/Ga-Adangbe	-0.01	(0.01)
Gruma/Mole-Dagbani	-0.02	(0.01)
Other	0.00	(0.01)
Religion (ref: Christian)		
Muslim	0.03	(0.02)
Traditional	-0.01	(0.02)
Other/None	0.01	(0.02)
Education (ref: no formal education)		
Primary	0.00	(0.01)
Secondary/Higher	0.00	(0.01)
Current employ (ref: unemployed)		
Working	0.01	(0.01)
Sector of employment (ref: informal)		
Public/private	-0.01	(0.01)
Informal sector	-0.01	(0.01)
Marital status (ref: married)		
Separated/divorced	0.00	(0.02)
Widowed	-0.02	(0.01)**
Location (ref: rural)		
Urban	-0.02	(0.01)
NCD status (ref: no)		
Yes	-0.02	(0.01)**
Self-Rated Health (ref: good)		
Moderate	-0.01	(0.01)
Bad	0.01	(0.01)

*** p < 0.001, ** p < 0.01, * p < 0.05

Multivariate

It is important to note that there was a general significant increase in the size of the parameter estimates in the multivariate results (see Table 3). Figure 2 shows the visual comparison in the changes in the coefficient estimates for the focal independent variables between Models 1 and 2. In Model 1, only income quintile was

Table 3. Multivariate negative loglog regression of ‘frequently used health care type’ (n = 2256)

	Model 1		Model 2	
	Coef.	(Std. Err.)	Coef.	(Std. Err.)
Insurance status (ref: uninsured)				
Insured	-0.14	(0.09)	-0.17	(0.09)*
Income quintile (ref: richest)				
Richer	0.37	(0.19)	0.40	(0.18)*
Middle	0.56	(0.19)**	0.62	(0.19)**
Poorer	0.37	(0.20)	0.46	(0.20)*
Poorest	0.48	(0.18)**	0.61	(0.19)***
Age				
			0.00	(0.01)
Gender (ref: male)				
Female			0.06	(0.11)
Ethnicity (ref: Akan)				
Ewe/Ga-Adangbe			-0.15	(0.13)
Gruma/Mole-Dagbani			-0.38	(0.22)
Other			-0.17	(0.13)
Religion (ref: Christian)				
Muslim			0.36	(0.14)*
Traditional			-0.08	(0.19)
Other/none			0.10	(0.17)
Education (ref: no formal education)				
Primary			0.06	(0.10)
Secondary/higher			0.17	(0.15)
Current employ (ref: unemployed)				
Working			0.05	(0.10)
Sector of employment (ref: informal)				
Public/private			-0.03	(0.18)
Informal sector			-0.18	(0.16)
Marital status (ref: married)				
Separated/divorced			0.01	(0.12)
Widowed			-0.29	(0.12)*
Location (ref: rural)				
Urban			-0.01	(0.09)
NCD status (ref: no)				
Yes	-0.16	(0.09)	-0.15	(0.09)
Self-rated health (ref: good)				
Moderate	-0.10	(0.10)	-0.08	(0.09)
Bad	0.08	(0.12)	0.12	(0.12)

***p < 0.001, ** p < 0.01, * p < 0.05

significantly associated with ‘frequently used health care type’ in the last three years. Further investigation revealed that income quintile suppressed the relationship between insurance status




Figure 2. Multivariate negative loglog regression of ‘frequently used health care type’ (n = 2256)

and ‘frequently used health care type’. The findings show that seniors who belonged in the middle income quintile had 56% increased odds of seeking TM compared to the richest. Compared to the richest, the odds for the poorest seniors frequently seeking TM were 48% higher.

In Model 2, while the relationship between income quintile and ‘frequently used health care type’ remained robust, the relationship between insurance status and ‘frequently used health care type’ also became statistically significant. Further analysis revealed that the relationship between insurance status and ‘frequently used health care type’ was mediated by marital status. Seniors who had health insurance coverage were also 17% less likely to frequently seek treatment from a TM healer relative to the uninsured. The odds of richer seniors seeking treatment from a TM increased by 40% compared to the richest. Seniors in the poorest income quintile observed the highest percentage increase in coefficient estimates between Model 1 and Model 2 (see Figure 2). For this group, the odds of frequently seeking TM increased by 61% when compared to those in the richest quintile.

The results also show that two of the selected socioeconomic variables emerged as significant predictors of ‘frequently used health care type’ in the last three years. While religion had a positive relationship with ‘frequently used health care type’, marital status had a negative relationship. Specifically, seniors who were identified as Muslims had 36% increased odds of frequently seeking TM healers compared to Christians. On the other hand, seniors who indicated they were widowed had 29% decreased odds of frequently seeking TM relative to those who were married.

Discussion and Conclusion

Our study sought to examine the effect of health insurance and wealth status on health care-seeking behaviour of older people in Ghana. It is a widely held belief that TM is a popular form of health care – in terms of access, cost, availability and acceptance – in Ghana and other SSA countries (Pouliot, 2011; Sato, 2012a; Powell-Jackson *et al.*, 2014). Recent evidence in the literature questions this widely acclaimed popularity of TM in Ghana and SSA, particularly without adequate consideration for the context of utilization (Amegbor, 2017a, 2017b; Sato, 2012b). The findings of this study show that the majority of Ghanaian older adults (96.8%) reported using modern health care, which is consistent

with evidence from other recent studies (Sato, 2012b). The popularity of modern care is restricted not only to health care facilities but also in self-care managing of common diseases such as malaria (Fenny *et al.*, 2015). In cases of recurring diseases and chronic diseases, the use of TM tends to be relatively higher than biomedicine (Aikins, 2005; Sato, 2012b). The literature further indicates that in remote and rural areas where there are deficiencies in modern health facilities, the population are more likely to use TM to address health care needs (Aikins & Marks, 2007; Powell-Jackson *et al.*, 2014). Findings in this paper demonstrate the need to contextualize TM utilization with prevailing health care cost and means of payment.


Health insurance coverage remains a crucial enabling resource for accessing modern health care services (Fenny *et al.*, 2016). It provides a higher degree of financial risk protection against cost associated with seeking treatment. It also reduces health care expenditure significantly for insured persons (Saksena *et al.*, 2011; Kusi *et al.*, 2015; Kansanga *et al.*, 2018). However, these risk-pooling and financial protection do not cover TM or other alternative forms of health care. Among the poor and vulnerable, such as older persons in Ghana, inability to pay for modern health care results in delays in seeking treatment or forgoing treatment (Robyn *et al.*, 2012). It is also important to note that a considerable proportion of Ghana's older adult population is uninsured despite health premium age-exemption criterion for persons 70 years or older (Parmar *et al.*, 2014). In view of this, we argue that uninsured older persons do not have the financial cushion associated with seeking modern health care; hence, they may resort to less expensive forms of treatment such as using TM or self-medicating with TM (Amegbor, 2017a, 2017b).

Income remains an important determinant of health care-seeking behaviour and health insurance enrolment. A considerable number of persons of lower income status are unenrolled in the NHIS, a situation largely attributable to limited financial capabilities to meet the insurance premium (Dixon *et al.*, 2011; Jehu-Appiah *et al.*, 2011; Aryeetey *et al.*, 2013; Kusi *et al.*, 2015). In the midst of widespread poverty, individuals often have to make a budgetary trade-off for vital goods such as food, in order to afford the health insurance premiums (Dixon *et al.*, 2011; 2014; Fenny *et al.*, 2016). Economic vulnerability of persons of lower income, as well as their inability to afford health insurance premiums, means this section of the Ghanaian population have limited access to and use of modern health care services. As evident in this study, older persons in lower income quintiles were more likely to report using TM for their regular health care needs. This may be due to their inability to afford the cost associated with seeking modern care. TM offers a cheaper alternative form of health care for the poor and vulnerable; and it is also readily available in rural communities where most low income older adults live (Parmar *et al.*, 2014). As shown in this paper and others, persons with higher incomes are less reliant on TM for their regular health care needs (Pouliot, 2011; Sato, 2012a; Thorsen & Pouliot, 2015), partly because they can afford the cost associated with seeking modern health care as well as pay premiums for health insurance coverage to guarantee access to modern care when needed (Jehu-Appiah *et al.*, 2011; Sato, 2012a; Fenny *et al.*, 2016; Dalinjong *et al.*, 2017). The rich turn to TM only in cases of chronic health where biomedical care fails to provide a cure (Aikins, 2005; Sato, 2012b). Even in such situations, they rely on the services of licensed and commercial TM practitioners compared to the poor and disadvantaged who mostly rely on TM for self-treatment or rely on non-commercial TM healers (Amegbor, 2017a; Pouliot, 2011).

The findings of our study have several policy implications. Although numerous studies have examined the effect of income and health insurance status on access to modern care, our understanding of how these two variables influence choice between TM and modern care use remain limited. Findings from this study demonstrate that TM is a vital health care resource for poor and uninsured older persons in Ghana. However, the existence of quack healers, untested or counterfeit TM remedies pose a danger to the uninsured and lower income older adults who depend on TM for primary health care and other health needs. Regulation of TM practices and products in Ghana, like most SSA countries, remains a challenge – providing justification for their current exclusion from public health delivery under the Ministry of Health. TM products and remedies in Ghana are largely untested and do not have approval from state regulatory institutions such as the Centre for Plant Medicine Research (CSRPM), the Food and Drugs Authority and the Ghana Standards Authority. Consequently, some TM products may pose health dangers to users due to their potential toxicities (Pouliot, 2011; Sato, 2012b). Thus, there is the need to fast track the inclusion of TM practitioners, drugs and health centres under the Ministry of Health. Including TM under the Ministry of Health must only be undertaken after ascertaining, on a case-by-case basis, efficacy and establishing best practices. This will help minimize the risk of harm for the public, especially for the uninsured and the poor who frequently rely on TM during times of illness.

The findings of this study ought to be considered in view of some limitations. TM, as used in this study does not incorporate the use of TM in self-care context. While a substantial majority of older adults use modern health care for their regular health care needs, the literature demonstrates a high level of TM use in self-care context (Danso-Appiah *et al.*, 2010; Pouliot, 2011; Chipwaza *et al.*, 2014). Respondents of the SAGE survey may have limited their response to TM to professional care as the response categories only reference the use of professional TM service. Also, the WHO SAGE wave 1 is a cross-sectional data; hence, we cannot draw causal relationships between health insurance status and the 'frequently used health care type' or income status and the 'frequently used health care type'. Future studies can use subsequent waves of SAGE in addition to Wave 1 to examine the temporal effect of health insurance and wealth status on the use of TM among older Ghanaians. Our study also focused on frequent health care-seeking behaviour which may not be similar to health care-seeking behaviour for specific illnesses or disease episodes (Amegbor, 2017b).

Concerns around efficacy and dosage of TM remain important discussion points that drive the conversation towards incorporating TM in mainstream health care delivery in Ghana. While such ongoing discussions remain important, it is imperative to understand TM utilization patterns, particularly in the context of Ghana's NHIS health equity mandate. In this paper, we provided evidence of the factors associated with TM use among older Ghanaians. This understanding is important because it provides useful information that can be incorporated in the broader discussion of strategies to improve primary health care access from the perspective of older people's health utilization behaviours. The findings from our study demonstrate that TM is a popular source of access to health care among uninsured and older persons with lower incomes. This reiterates the need to improve NHIS coverage for poor older persons and calls for the need to regulate TM and incorporate it into the broader health care system. This will go a long way to protect the poor and uninsured older people from seeking treatment from unlicensed TM practitioners.

Author ORCIDs.  Prince M. Amegbor 0000-0001-7701-2585

Acknowledgements. We are grateful for the helpful comments from the editors and anonymous reviewers. We are also grateful to the WHO SAGE programme and the Inter-University Consortium for Political and Social Research (ICPSR) for the data used in this study.

Conflict of Interest. None

Ethical Standards. We used data from the World Health Organisation (WHO)'s Study on Global Ageing and Adult Health (SAGE) Wave 1. Our study used a secondary survey data; hence, ethical approval was not required. We accessed the data through Queen's University library from the Inter-University Consortium for Political and Social Research (ICPSR) platform.

References

- Abdullahi AA** (2011) Trends and challenges of traditional medicine in Africa. *African Journal of Traditional, Complementary and Alternative Medicines* 8 (5 Suppl.), 115–123. doi: [10.4314/ajtcam.v8i5.S.5](https://doi.org/10.4314/ajtcam.v8i5.S.5).
- Agyepong IA and Adjei S** (2008) Public social policy development and implementation: a case study of the Ghana National Health Insurance scheme. *Health Policy and Planning* 23, 150–160. doi: [10.1093/heapol/czn002](https://doi.org/10.1093/heapol/czn002).
- Aikins de-GA** (2005) Healer shopping in Africa: new evidence from rural-urban qualitative study of Ghanaian diabetes experiences. *BMJ (Clinical Research ed.)* 331, 737. doi: [10.1136/bmj.331.7519.737](https://doi.org/10.1136/bmj.331.7519.737).
- Aikins de-GA, Awuah RB, Pera TA, Mendez M and Ogedegbe G** (2015) Explanatory models of diabetes in urban poor communities in Accra, Ghana. *Ethnicity & Health* 20, 391–408. doi: [10.1080/13557858.2014.921896](https://doi.org/10.1080/13557858.2014.921896).
- Aikins de-GA, Boynton P and Atanga LL** (2010) Developing effective chronic disease interventions in Africa: insights from Ghana and Cameroon. *Globalization and Health* 6, 6. doi: [10.1186/1744-8603-6-6](https://doi.org/10.1186/1744-8603-6-6).
- Aikins de-GA, Kushitor M, Koram K, Gyamfi S and Ogedegbe G** (2014) Chronic non-communicable diseases and the challenge of universal health coverage: insights from community-based cardiovascular disease research in urban poor communities in Accra, Ghana. *BMC Public Health* 14 (Suppl 2), S3. doi: [10.1186/1471-2458-14-S2-S3](https://doi.org/10.1186/1471-2458-14-S2-S3).
- Aikins de-GA and Marks DF** (2007) Health, disease and healthcare in Africa. *Journal of Health Psychology* 12, 387–402. doi: [10.1177/1359105307076228](https://doi.org/10.1177/1359105307076228).
- Akazili J, Garshong B, Aikins M, Gyapong J, McIntyre D** (2012) Progressivity of health care financing and incidence of service benefits in Ghana. *Health Policy and Planning* 27 (Suppl 1), i13–i22. doi: [10.1093/heapol/czs004](https://doi.org/10.1093/heapol/czs004).
- Amegbor PM** (2014) *Health seeking behaviour in Asikuma-Odoben-Brakwa district: A pluralistic health perspective*. Oslo, Norway: University of Oslo.
- Amegbor PM** (2017a) An assessment of care-seeking behavior in Asikuma-Odoben-Brakwa district: a triple pluralistic health sector approach. *SAGE Open* 7, 1–17. doi: [10.1177/2158244017710688](https://doi.org/10.1177/2158244017710688).
- Amegbor PM** (2017b) Understanding usage and preference for health care therapies in a Ghanaian context: a pluralistic perspective. *Norwegian Journal of Geography* 71, 288–300. doi: [10.1080/00291951.2017.1340908](https://doi.org/10.1080/00291951.2017.1340908).
- Amegbor PM, Kuire VZ, Robertson H and Kuffuora OA** (2018) Predictors of basic self-care and intermediate self-care functional disabilities among older adults in Ghana. *Archives of Gerontology and Geriatrics* 77, 81–88. doi: [10.1016/j.archger.2018.04.006](https://doi.org/10.1016/j.archger.2018.04.006).
- Aryeetey GC, Jehu-Appiah C, Kotoh AM, Spaan E, Arhinful DK, Baltussen R, van der Geest S and Agyepong IA** (2013) Community concepts of poverty: an application to premium exemptions in Ghana's National Health Insurance Scheme. *Globalization and Health* 9, 12. doi: [10.1186/1744-8603-9-12](https://doi.org/10.1186/1744-8603-9-12).
- Barimah KB** (2013) Traditional healers as service providers in Ghana's National Health Insurance Scheme: the wrong way forward? *Global Public Health* 8, 202–208. doi: [10.1080/17441692.2012.761262](https://doi.org/10.1080/17441692.2012.761262).
- Bennett S** (2004) The role of community-based health insurance within the health care financing system: a framework for analysis. *Health Policy and Planning* 19, 147–158. doi: [10.1093/heapol/czh018](https://doi.org/10.1093/heapol/czh018).
- Blanchet NJ, Fink G and Osei-Akoto I** (2012) The effect of Ghana's National Health Insurance Scheme on health care utilisation. *Ghana Medical Journal* 46, 76–84.
- Chipwaza B, Mugasa JP, Mayumana I, Amuri M, Makungu C and Gwakisa PS** (2014) Self-medication with anti-malarials is a common practice in rural communities of Kilosa district in Tanzania despite the reported decline of malaria. *Malaria Journal* 13, 1–11. doi: [10.1186/1475-2875-13-252](https://doi.org/10.1186/1475-2875-13-252).
- Dalinjong PA, Welaga P, Akazili J, Kwarteng A, Bangha M, Oduro A, Sankoh O and Goudge J** (2017) The association between health insurance status and utilization of health services in rural Northern Ghana: evidence from the introduction of the National Health Insurance Scheme. *Journal of Health, Population and Nutrition* 36, 1–10. doi: [10.1186/s41043-017-0128-7](https://doi.org/10.1186/s41043-017-0128-7).
- Danso-Appiah A, Stolk WA, Bosompem KM, Otchere J, Looman CW, Habbema JD and de Vlas SJ** (2010) Health seeking behaviour and utilization of health facilities for schistosomiasis-related symptoms in Ghana. *PLoS Neglected Tropical Diseases* 4, e867. doi: [10.1371/journal.pntd.0000867](https://doi.org/10.1371/journal.pntd.0000867).
- Dixon J, Luginaah I and Mkandawire P** (2014) The national health insurance scheme in Ghana's upper west region: a gendered perspective of insurance acquisition in a resource-poor setting. *Social Science & Medicine* 122, 103–112.
- Dixon J, Tenkorang EY and Luginaah I** (2011) Ghana's national health insurance scheme: helping the poor or leaving them behind?. *Environment and Planning C: Government and Policy* 29, 1102–1115.
- Fenny AP, Asante FA, Arhinful DK, Kusi A, Parmar D and Williams G** (2016) Who uses outpatient healthcare services under Ghana's health protection scheme and why?. *BMC Health Services Research* 16, 1–8. doi: [10.1186/s12913-016-1429-z](https://doi.org/10.1186/s12913-016-1429-z).
- Fenny AP, Asante FA, Enemark U and Hansen KS** (2015) Malaria care seeking behavior of individuals in Ghana under the NHIS: are we back to the use of informal care? *BMC Public Health* 15, 370. doi: [10.1186/s12889-015-1696-3](https://doi.org/10.1186/s12889-015-1696-3).
- Gnawali DP, Pokhrel S, Sié A, Sanon M, De Allegri M, Souares A, Dong H and Sauerborn R** (2009) The effect of community-based health insurance on the utilization of modern health care services: evidence from Burkina Faso. *Health Policy (Amsterdam, Netherlands)* 90, 214–222. doi: [10.1016/j.healthpol.2008.09.015](https://doi.org/10.1016/j.healthpol.2008.09.015).
- Jehu-Appiah C, Aryeetey G, Spaan E, de Hoop T, Agyepong I and Baltussen R** (2011) Equity aspects of the National Health Insurance Scheme in Ghana: who is enrolling, who is not and why? *Social Science and Medicine* 72, 157–165.
- Jowett M, Deolalikar A and Martinsson P** (2004) Health insurance and treatment seeking behaviour: evidence from a low-income country. *Health Economics* 13, 845–857. doi: [10.1002/hec.862](https://doi.org/10.1002/hec.862).
- Kansanga MM, Asumah Braimah J, Antabe R, Sano Y, Kyeremeh E and Luginaah I** (2018) Examining the association between exposure to mass media and health insurance enrolment in Ghana. *The International Journal of Health Planning and Management* 33, e531–e540. doi: [10.1002/hpm.2505](https://doi.org/10.1002/hpm.2505).
- Kigen GK, Ronoh HK, Kipkore WK and Rotich JK** (2013) Current trends of traditional herbal medicine practice in Kenya: a review. *African Journal of Pharmacology and Therapeutics* 2, 32–37.
- Kusi A, Enemark U, Hansen KS and Asante FA** (2015) Refusal to enrol in Ghana's National Health Insurance Scheme: is affordability the problem? *International Journal for Equity in Health* 14, 2.
- Kuire VZ, Bisung E, Rishworth A, Dixon J and Luginaah I** (2015) Health-seeking behaviour during times of illness: a study among adults in a resource poor setting in Ghana. *Journal of Public Health* 38(4), e545–e553. doi: [10.1093/pubmed/fdv176](https://doi.org/10.1093/pubmed/fdv176)
- Kuire VZ, Tenkorang EY, Rishworth A, Luginaah I and Yawson AE** (2017) Is the pro-poor premium exemption policy of Ghana's NHIS reducing disparities among the elderly? *Population Research and Policy Review* 36, 231–249.
- Lakshmi JK, Nambiar D, Narayan V, Sathyanarayana TN, Porter J and Sheikh K** (2015) Cultural consonance, constructions of science and co-existence: a review of the integration of traditional, complementary and alternative medicine in low- and middle-income countries. *Health Policy and Planning* 30, 1067–1077. doi: [10.1093/heapol/czu096](https://doi.org/10.1093/heapol/czu096).
- Macha J, Harris B, Garshong B, Ataguba JE, Akazili J, Kuwawenaruwa A and Borghi J** (2012) Factors influencing the burden of health care financing and the distribution of health care benefits in Ghana, Tanzania and South Africa. *Health Policy and Planning* 27 (Suppl. 1), 46–54. doi: [10.1093/heapol/czs024](https://doi.org/10.1093/heapol/czs024).

- McFarlane C** (2015) South Africa: the rise of traditional medicine. *Insight on Africa* 7, 60–70. doi: [10.1177/0975087814554070](https://doi.org/10.1177/0975087814554070).
- McIntyre D, Thiede M, Dahlgren G and Whitehead M** (2006) What are the economic consequences for households of illness and of paying for health care in low- and middle-income country contexts? *Social Science and Medicine* 62, 858–865. doi: [10.1016/j.socscimed.2005.07.001](https://doi.org/10.1016/j.socscimed.2005.07.001).
- Minicuci N, Biritwum RB, Mensah G, Yawson AE, Naidoo N, Chatterji S and Kowal P** (2014) Sociodemographic and socioeconomic patterns of chronic non-communicable disease among the older adult population in Ghana. *Global Health Action* 7, 21292. doi: [10.3402/gha.v7.21292](https://doi.org/10.3402/gha.v7.21292).
- O'Brien KS, Soliman AS, Annan K, Lartey RN, Awuah B and Merajver SD** (2012) Traditional herbalists and cancer management in Kumasi, Ghana. *Journal of Cancer Education* 27, 573–579. doi: [10.1007/s13187-012-0370-z](https://doi.org/10.1007/s13187-012-0370-z).
- Parmar D, Williams GA, Dkhimi F, Ndiaye AI, Asante FA, Arhinful DK and Mladovsky P** (2014) Enrolment of older people in social health protection programs in West Africa – Does social exclusion play a part? *Social Science and Medicine* 119, 36–44.
- Pouliot M** (2011) Relying on nature's pharmacy in rural Burkina Faso: empirical evidence of the determinants of traditional medicine consumption. *Social Science and Medicine* 73, 1498–1507. doi: [10.1016/j.socscimed.2011.08.037](https://doi.org/10.1016/j.socscimed.2011.08.037).
- Powell-Jackson T, Hanson K, Whitty CJM and Ansah EK** (2014) Who benefits from free healthcare? Evidence from a randomized experiment in Ghana. *Journal of Development Economics* 107, 305–319. doi: [10.1016/j.jdeveco.2013.11.010](https://doi.org/10.1016/j.jdeveco.2013.11.010).
- Prince MJ, Wu F, Guo Y, Gutierrez Robledo LM, O'Donnell M, Sullivan R and Yusuf S** (2015) The burden of disease in older people and implications for health policy and practice. *The Lancet* 385, 549–562. doi: [10.1016/S0140-6736\(14\)61347-7](https://doi.org/10.1016/S0140-6736(14)61347-7).
- Robyn PJ, Fink G, Sié A and Sauerborn R** (2012) Health insurance and health-seeking behavior: evidence from a randomized community-based insurance rollout in rural Burkina Faso. *Social Science and Medicine* 75, 595–603. doi: [10.1016/j.socscimed.2011.12.018](https://doi.org/10.1016/j.socscimed.2011.12.018).
- Robyn PJ, Sauerborn R and Barnighausen T** (2013) Provider payment in community-based health insurance schemes in developing countries: a systematic review. *Health Policy and Planning* 28, 111–122. doi: [10.1093/heapol/czs034](https://doi.org/10.1093/heapol/czs034).
- Saksena P, Antunes AF, Xu K, Musango L and Carrin G** (2011) Mutual health insurance in Rwanda: evidence on access to care and financial risk protection. *Health Policy* 99, 203–209. doi: [10.1016/j.healthpol.2010.09.009](https://doi.org/10.1016/j.healthpol.2010.09.009).
- Sato A** (2012a) Does socio-economic status explain use of modern and traditional health care services? *Social Science and Medicine* 75, 1450–1459. doi: [10.1016/j.socscimed.2012.05.032](https://doi.org/10.1016/j.socscimed.2012.05.032).
- Sato A** (2012b) Revealing the popularity of traditional medicine in light of multiple recourses and outcome measurements from a user's perspective in Ghana. *Health Policy and Planning* 27, 625–637. doi: [10.1093/heapol/czs010](https://doi.org/10.1093/heapol/czs010).
- Tabi M, Powell M and Hodnicki D** (2006) Use of traditional healers and modern medicine in Ghana. *International Nursing Review* 53, 52–58. doi: [10.1111/j.1466-7657.2006.00444.x](https://doi.org/10.1111/j.1466-7657.2006.00444.x).
- Tagoe HA** (2012) Household burden of chronic disease in Ghana. *Ghana Medical Journal* 46, 54–58.
- Thorsen RS and Pouliot M** (2015) Traditional medicine for the rich and knowledgeable: challenging assumptions about treatment-seeking behaviour in rural and peri-urban Nepal. *Health Policy and Planning* 31, 314–324. doi: [10.1093/heapol/czv060](https://doi.org/10.1093/heapol/czv060).
- Towns AM and Van Andel T** (2016) Wild plants, pregnancy, and the food-medicine continuum in the southern regions of Ghana and Benin. *Journal of Ethnopharmacology* 179, 375–382. doi: [10.1016/j.jep.2016.01.005](https://doi.org/10.1016/j.jep.2016.01.005).
- van der Wielen N, Channon AA and Falkingham J** (2018) Does insurance enrolment increase healthcare utilisation among rural-dwelling older adults? Evidence from the National Health Insurance Scheme in Ghana. *BMJ Global Health* 3, e000590. doi: [10.1136/bmjgh-2017-000590](https://doi.org/10.1136/bmjgh-2017-000590).
- Witter S and Garshong B** (2009) Something old or something new? Social health insurance in Ghana. *BMC International Health and Human Rights* 9, 20. doi: [10.1186/1472-698X-9-20](https://doi.org/10.1186/1472-698X-9-20).
- World Health Organisation** (2002) *Traditional medicine strategy 2002–2005*. World Health Organization: Programme on traditional medicine. Geneva, Switzerland: WHO.
- World Health Organisation** (2013) *WHO traditional medicine strategy 2014–2023*. Geneva, Switzerland: WHO.
- World Health Organisation** (2014) *Ghana country assessment report on ageing and health*. Geneva, Switzerland: WHO.