

# THE MARINE BIOLOGICAL ASSOCIATION OF THE UNITED KINGDOM

## TERMS OF MEMBERSHIP FOR 2018

MBA MEMBERSHIP (prices per annum)	Price from 1st March 2018
Young Marine Biologist (YMB)	£12.50
Student Member	£25.00
Associate Member	£40.00 (direct debit discount £5)
Professional Member, Mem.MBA	£45.00 (direct debit discount £5)
MBA Fellow, FMBA	£120.00 (direct debit discount £5)
Institutional Member	£340.00

JOURNAL SUBSCRIPTION (MBA price added to the above membership fees, with the exception of YMB as no Journal option for this category)	
Online Journal	£20.00
Print & Online Journal	£80.00

The Association is a learned society founded in 1884 and incorporated by Royal Charter in 2013. The Marine Biological Association (MBA) provides a unified, clear, independent voice on behalf of the marine biological community. In 1888, the MBA opened the Plymouth laboratory at Citadel Hill.

The MBA is also a registered charity with its key aims being to promote scientific research into all aspects of life in the sea, including the environment on which it depends, and to disseminate to the public the knowledge gained. The Patron of the Association is His Royal Highness the Duke of Edinburgh, who is also an Honorary Fellow. The other Honorary Fellows are His Serene Highness Prince Albert of Monaco, Dr Sylvia Earle, Sir Tim Hunt and Professor James Lovelock.

The work of the Association is controlled by the Council elected annually by its subscribing members.

The Marine Biological Association, under the direction of its Council, has undertaken research in all branches of marine science and many of the results have been published in this Journal. An account of the history of the first hundred years of the Association and Journal may be found in Volume 67 (pp. 463 and 465).

Since its foundation the Association has been supported by grants from Research Councils and by subscriptions and donations

from private members, universities, learned societies, the Fishmongers' Company and other public bodies.

The Marine Biological Association is a very active scientific learned society with a large international membership. The MBA also manages an internationally renowned research programme.

Subscriptions can be paid by cash/cheque, direct debit from a UK bank account or credit card. \*Subscriptions are reduced by £5 per annum if paying by direct debit.

Members of the Association have a wide range of rights and privileges including the use of post-nominals (FMBA for MBA Fellows, Mem.MBA for Professional Members) and being entitled to receive The Marine Biologist Magazine. The Commissioners of Inland Revenue have approved the Association for the purposes of Section 16, Finance Act, 1958, and that the whole of the annual subscriptions paid by a member who qualifies for relief under the section will be allowable as a deduction from his emoluments assessable to income tax.

All correspondence should be addressed to: The Membership Secretary, Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK (Tel. +44 (0) 1752 426493; e-mail: [membership@mba.ac.uk](mailto:membership@mba.ac.uk)).

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE  
The Pitt Building, Trumpington Street, Cambridge, CB2 1RP, United Kingdom

This journal issue has been printed on FSC-certified paper and cover board. FSC is an independent, non-governmental, not-for-profit organization established to promote the responsible management of the world's forests. Please see [www.fsc.org](http://www.fsc.org) for information.

Printed by Bell and Bain Limited, Glasgow

CAMBRIDGE UNIVERSITY PRESS

Journals Fulfillment Department, UPH, Shaftesbury Road, Cambridge CB2 8BS, United Kingdom  
1 Liberty Plaza, Floor 20, New York, NY 10006, USA  
477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
C/ Orense, 4, planta 13, 28020, Madrid, Spain  
Dock House, The Waterfront, Cape Town 8001, South Africa

## CONTENTS

Lewis, J. Editorial	645
David, A.A. and Loveday, B.R. The role of cryptic dispersal in shaping connectivity patterns of marine populations in a changing world	647
Yamaguchi, H., Tanimoto, Y., Hayashi, Y., Suzuki, S., Yamaguchi, M. and Adachi, M. Bloom dynamics of noxious <i>Chattonella</i> spp. (Raphidophyceae) in contrastingly enclosed coastal environments: a comparative study of two coastal regions	657
Wang, Z., Xu, G., Zhao, L., Gao, Y., Al Mamun, A. and Xu, H. A community-based approach to identifying defence of microalgae against protozoan grazing	665
Du, P., Liao, Y.-B., Jiang, Z.-B., Wang, K., Zeng, J.-N., Shou, L., Xu, X.-Q., Xu, X.-D., Liu, J.-J., Huang, W. and Zhang, D.-M. Responses of mesozooplankton communities to different anthropogenic activities in a subtropical semi-enclosed bay	673
Pocklington, J.B., Jenkins, S.R., Bellgrove, A., Keough, M.J., O'Hara, T.D., Masterson-Algar, P.E. and Hawkins, S.J. Disturbance alters ecosystem engineering by a canopy-forming alga	687
Quillien, N., Nordström, M.C., Le Bris, H., Bonsdorff, E. and Grall, J. Green tides on inter- and subtidal sandy shores: differential impacts on infauna and flatfish	699
Rocha, C.P., Quadros, M.L.A., Maciel, M., Maciel, C.R. and Abrunhosa, F.A. Morphological changes in the structure and function of the feeding appendages and foregut of the larvae and first juvenile of the freshwater prawn <i>Macrobrachium acanthurus</i>	713
Erdogan Sağlam, N., Demir Sağlam, Y. and Sağlam, C. A study on some population parameters of mantis shrimp ( <i>Squilla mantis</i> L., 1758) in Izmir Bay (Aegean Sea)	721
Pescinelli, R.A., Davanzo, T.M., Pantaleão, J.A.F., Carvalho-Batista, A., Bauer R.T. and Costa, R.C. Population dynamics, relative growth and sex change of the protandric simultaneous hermaphrodite <i>Exhippolysmata oplophoroides</i> (Caridea: Lysmatidae) close to an upwelling area	727
Yamada, K., Miyazaki, K., Tomiyama, T., Kanaya, G., Miyama, Y., Yoshinaga, T., Wakui, K., Tamaoki, M. and Toba, M. Impact of sea spider parasitism on host clams: susceptibility and intensity-dependent mortality	735
Aviz, D., Pinto, A.P.A., Ferreira, M.A.P., Rocha, R.M. and Filho, J.S.R. Reproductive biology of <i>Sabellaria wilsoni</i> (Sabellariidae: Polychaeta), an important ecosystem engineer on the Amazon coast	743
Lucas, Y., Sikorski, A. and Martin, G.S. Syllidae (Annelida) from the Boreal and sub-Arctic seas off Norway (North Sea, Norwegian Sea, Barents Sea)	755
Vasileiadou, K., Pavlou, C., Camisa, F., Tsikopoulou, I., Fragopoulou, N., Kotoulas, G. and Arvanitidis, C. Genetic diversity of <i>Nephtys hombergii</i> (Phyllodoce, Polychaeta) associated with environmental factors in a highly fluctuating ecosystem	777
Perry, O., Sapir, Y., Perry, G., Ten Hove, H. and Fine, M. Substrate selection of Christmas tree worms ( <i>Spirobranchus</i> spp.) in the Gulf of Eilat, Red Sea	791
Yu, T. and Xu, K. Two new oxystominid species (Nematoda: Enoplida) from an abyssal plain in the southern Philippine Sea	801
Terrón-Sigler, A., León-Muez, D., Peñalver-Duque, P. and Espinosa Torre, F. A method for increasing the survival of shallow-water populations of the endemic coral <i>Astroides calycularis</i>	811
Jawad, L.A., Hoedemakers, K., Ibáñez, A.L., Ahmed, Y.A., El-Regal, M.A.A. and Mehanna, S.F. Morphology study of the otoliths of the parrotfish, <i>Chlorurus sordidus</i> (Forsskal, 1775) and <i>Hipposcarus harid</i> (Forsskal, 1775) from the Red Sea coast of Egypt (Family: Scaridae)	819
Ventura, D., Machado, L.F., Bonifazi, A., Gravina, M.F. and Arduzzone, G. Cleaning interactions between the cleaner wrasse <i>Symphodus melanocercus</i> (Osteichthyes: Labridae) and brown meagre <i>Sciaena umbra</i> (Osteichthyes: Sciaenidae)	829
Hernández-Aguilar, S.B., Zenteno-Savín, T., De-Anda-Montañez, J.A. and Méndez-Rodríguez, L.C. Temporal variation in oxidative stress indicators in liver of totoaba ( <i>Totoaba macdonaldi</i> ) Perciformes: Sciaenida	833
Compaire, J.C., Casademont, P., Cabrera, R., Gómez-Cama, C. and Sorriquer, M.C. Feeding of <i>Scorpaena porcus</i> (Scorpaenidae) in intertidal rock pools in the Gulf of Cadiz (NE Atlantic)	845
Avigliano, E., Carvalho, B., Velasco, G., Tripodi, P. and Volpedo, A.V. Inter-annual variability in otolith chemistry of catfish <i>Genidens barbuis</i> from South-western Atlantic estuaries	855
Zappes, C.A., Gama, R.M., Domit, C., Gatts, C.E.N. and Di Benedetto, A.P.M. Artisanal fishing and the franciscana ( <i>Pontoporia blainvillei</i> ) in Southern Brazil: ethnoecology from the fishing practice	867
Kara, A., Sağlam, C., Acarli, D. and Cengiz, Ö. Length-weight relationships for 48 fish species of the Gediz estuary, in Izmir Bay (Central Aegean Sea, Turkey)	879
Muro-Torres, V.M., Amezcua, F., Lara-Mendoza, R.E., Buszkiewicz, J.T. and Amezcua-Linares, F. Trophic ecology of the chihuil sea catfish ( <i>Bagre panamensis</i> ) in the south-east Gulf of California, México	885
Abid, N., Laglaoui, A., Arakrak, A. and Bakkali, M. The role of fish in the diet of swordfish ( <i>Xiphias gladius</i> ) in the Strait of Gibraltar	895
Metochis, C.P., Carmona-Antoñanzas, G., Kousteni, V., Damalas, D. and Megalofonou, P. Population structure and aspects of the reproductive biology of the blackmouth catshark, <i>Galeus melastomus</i> Rafinesque, 1810 (Chondrichthyes: Scyliorhinidae) caught accidentally off the Greek coasts	909
Estupiñán-Montaño, C., Pacheco-Triviño, F., Cedeño-Figueroa, L.G., Galván-Magaña, F. and Estupiñán-Ortiz, J.F. Diet of three shark species in the Ecuadorian Pacific, <i>Carcharhinus falciformis</i> , <i>Carcharhinus limbatus</i> and <i>Nasolamia velox</i>	927
Anastasopoulou, A., Mytilineou, C., Makantasi, P., Smith, C.J., Kavadas, S., Lefkaditou, E. and Papadopoulou, K.N. Life history aspects of two species of the <i>Squalus</i> genus in the Eastern Ionian Sea	937
Lara-Lizardi, F., Hoyos-Padilla, M., Ketchum, J.T. and Galván-Magaña F. Range expansion of the whitenose shark, <i>Nasolamia velox</i> , and migratory movements to the oceanic Revillagigedo Archipelago (west Mexico)	949
Tseng, C.-T., Leu, J.-H. and Cheng, I.-J. On the genetic diversity of two species of the genus <i>Ozobranchus</i> (Hirudinida; Ozobranchidae) from the Atlantic and Pacific oceans	955
Nishizawa, H., Joseph, J., Chew, V.Y.-C., Liew, H.-C. and Chan, E.-H. Assessing tag loss and survival probabilities in green turtles ( <i>Chelonia mydas</i> ) nesting in Malaysia	961

Cambridge Core

For further information about this journal please go to the journal web site at: [cambridge.org/mbi](http://cambridge.org/mbi)



MIX  
Paper from  
responsible sources  
FSC® C007785

CAMBRIDGE  
UNIVERSITY PRESS