

Standard Paper

A new species and a new record of the genus *Squamulea* (Teloschistaceae, lichenized Ascomycota) from Pakistan

Najam-ul-Sehar Afshan , Iram Fayyaz , Fatima Iftikhar , Maria Jabeen  and Abdul Nasir Khalid 

Fungal Biology and Systematics Laboratory, Institute of Botany, University of the Punjab, Quaid-e-Azam Campus-54590, Lahore, Pakistan

Abstract

A novel species in the genus *Squamulea*, *S. chikarensis*, is described from Himalayan moist temperate forest in Pakistan. The morphology, chemistry and ITS sequences support its distinction from other species of this genus. The taxon is characterized by a pale green to yellow thallus, large apothecia up to 0.8–1.8 mm wide, pale yellow to yellow-orange apothecial discs, a hymenium 70–110 µm high, large ascospores (12–20 × 5–11 µm) and a narrow ascospore septum (1.5–3 µm). In addition, *S. flakusii* is reported as new to Pakistan and Eurasia.

Key words: *Caloplacoideae*, Himalayas, *Huriella*, lichen diversity, *Xanthorioideae*

(Accepted 11 December 2022; first published online 20 March 2023)

Introduction

In addition to the *Teloschistes*-type asci, most species of the lichen family *Teloschistaceae* are characterized by the presence of anthraquinones, which impart a yellowish orange colour to the thallus and/or apothecia. The widespread and well-delimited family currently comprises *c.* 110 genera and is estimated to contain over 1500 species (Arup *et al.* 2013; Kondratyuk *et al.* 2013, 2014, 2015a, b, c, d, 2016, 2017a; Søchting *et al.* 2014a, b; Wilk *et al.* 2021). The family is divided into three subfamilies, *Caloplacoideae*, *Teloschistoideae* and *Xanthorioideae* (Gaya *et al.* 2012; Arup *et al.* 2013) or four subfamilies (Kondratyuk *et al.* 2015d). The genus *Squamulea* is most similar to *Huriella* and is characterized by small, squamulose or areolate thalli containing anthraquinones, a saxicolous habit, and by producing apothecia with a paraplectenchymatous proper margin and hypothecium (Wilk 2020). From Pakistan, only one species of the genus *Squamulea* (namely *Squamulea subsoluta* (Nyl.) Arup *et al.* as *Caloplaca irrubescens* (Nyl. ex Arnold) Zahlbr. in Aptroot & Iqbal (2012)) has been previously identified.

During our exploration of the lichen diversity of Pakistan, collections of the genus *Squamulea* were made from Khyber Pakhtunkhwa Province and various sites in Azad Jammu and Kashmir, Pakistan. Using molecular analyses, as well as morphological and chemical characters, we were able to confirm the presence of one new species and one new record of the genus *Squamulea* in Pakistan, which are presented here.

Author for correspondence: Fatima Iftikhar. E-mail: fatimaiftikhar064@gmail.com

Cite this article: Afshan NS, Fayyaz I, Iftikhar F, Jabeen M and Khalid AN (2023) A new species and a new record of the genus *Squamulea* (Teloschistaceae, lichenized Ascomycota) from Pakistan. *Lichenologist* 55, 51–58. <https://doi.org/10.1017/S0024282923000026>

Materials and Methods


Morphological and chemical studies

Collections were made during a lichen survey of Azad Jammu and Kashmir, Kaghan Valley and Khanspur in 2021. The specimens were examined macro and micromorphologically using a stereomicroscope (EMZ-5TR, Meiji Techno, Japan) and compound microscope (SWIFT M4000-D) with a 9MP camera system, respectively. For anatomical investigation, sections of apothecia were made by hand and mounted in water and 10% KOH (K). A minimum of 20 measurements in water were made for each diagnostic feature from five specimens. The collected specimens were deposited in the herbarium of the Institute of Botany, University of the Punjab, Lahore (LAH). Secondary chemistry was analyzed using spot tests which were performed using 10% KOH and sodium hypochlorite solution (C). Thin-layer chromatography was carried out using solvent system C, following standard methods (Orange *et al.* 2010).

DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted directly from a portion of thallus with apothecia from each specimen using a modified 2% CTAB method (Gardes & Bruns 1993). The primer pair ITS1F (Gardes & Bruns 1993) and ITS4 (White *et al.* 1990) was used to amplify the internal transcribed spacer (ITS) regions, following the amplification protocol of Khan *et al.* (2018). PCR products were sent to BGI (Beijing Genomics Institute), China, where both strands were sequenced.

Sequences were assembled using BioEdit (Hall 1999). BLAST analyses (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) were used for initial verification of their identities and to retrieve highly similar ITS sequences. The newly generated sequences and additional sequences retrieved from GenBank were used in an initial



0.03

Fig. 1. Phylogenetic relationships within *Squamulea* based on a maximum likelihood (ML) analysis of the ITS region. The ML bootstrap values ≥ 70 obtained from a RaxML analysis are shown above internal branches. New sequences generated in this study are in bold. *Parvoplaca nigroblastidiata* was used as outgroup.

Table 1. Sequences used in the ITS phylogenetic analysis of *Squamulea* species, with GenBank Accession numbers and voucher information. New sequences generated in this study are in bold.

| Name of species | Country/Origin | Voucher specimen | GenBank Accession no. |
|---------------------------------------|-----------------------|---|-----------------------|
| <i>Parvoplaca nigroblastidiata</i> | USA | 42983 (BG) | KT161982 |
| <i>Squamulea chelonia</i> | Ecuador | Bungartz 6146 (CDS 34358) | MT967452 |
| <i>S. chelonia</i> | Ecuador | Bungartz 9251 (CDS 46069) | MT967451 |
| <i>S. chelonia</i> | Ecuador | Bungartz 4521 (CDS 28607) | MT967448 |
| <i>S. chikarensis</i> sp. nov. | Pakistan | Afshan et al. s. n. (LAH37546, holotype) | OP351700 |
| <i>S. chikarensis</i> sp. nov. | Pakistan | Afshan et al. s. n. (LAH37547) | OP351701 |
| <i>S. chikarensis</i> sp. nov. | Pakistan | Afshan & Iftikhar s. n. (LAH37548) | OP351702 |
| <i>S. flakusii</i> | Ecuador | Bungartz 4131 (CDS 28162) | MT967442 |
| <i>S. flakusii</i> | Ecuador | Bungartz 65261 (CDS 31847) | MT967444 |
| <i>S. flakusii</i> | Peru | Flakus 9263 & Cykowska (KRAM-L-70242) | MN108089 |
| <i>S. flakusii</i> | Pakistan | Afshan & Iftikhar s. n. (LAH37549) | OP341345 |
| <i>S. flakusii</i> | Pakistan | Afshan et al. s. n. (LAH37550) | OP341346 |
| <i>S. galactophylla</i> | USA | Morse 10997 (LD) | KC179122 |
| <i>S. humboldtiana</i> | Saint Kitts and Nevis | Buck 29560 (MIN) | MT967439 |
| <i>S. humboldtiana</i> | Ecuador | Bungartz 9985 (CDS 47354) | MT967441 |
| <i>S. humboldtiana</i> | Ecuador | Bungartz 4711 B (CDS 56235) | MT967440 |
| <i>S. kiamae</i> | Australia | Kondratyuk 20480 (LD, isotype) | KC179123 |
| <i>S. loekoesiana</i> | South Korea | KoLRI 40238 | KY614410 |
| <i>S. loekoesiana</i> | South Korea | KoLRI 40141 | KY614408 |
| <i>S. loekoesiana</i> | South Korea | KoLRI 15423 | KY614406 |
| <i>S. loekoesiana</i> | - | HKAS 102112 | MK499351 |
| <i>S. loekoesiana</i> | South Korea | KoLRI 40236 | KY614409 |
| <i>S. loekoesiana</i> | South Korea | KoLRI 19017 | KY614407 |
| <i>S. oceanica</i> | Ecuador | Bungartz 9857 (CDS 47195) | MT967447 |
| <i>S. oceanica</i> | Ecuador | Bungartz 10152 (CDS 47571) | MT967446 |
| <i>S. oceanica</i> | Ecuador | Bungartz 2023 (CDS 48373) | MT967445 |
| <i>S. osseophila</i> | Ecuador | Bungartz 65489 (CDS 32078) | MT967455 |
| <i>S. phyllidizans</i> | Ecuador | Bungartz 4710 (CDS 28808) | MT967457 |
| <i>S. phyllidizans</i> | Ecuador | Bungartz 65468 (CDS 32057) | MT967456 |
| <i>S. squamosa</i> | Ecuador | Bungartz 7428 (CDS 37915) | MT967465 |
| <i>S. squamosa</i> | Mexico | Moberg 8782 (UPS) | MT967462 |
| <i>S. squamosa</i> | USA | Kärnefelt AM960105 (LD) | KC179125 |
| <i>S. subsoluta</i> | Austria | Arup 97072 | AF353954 |
| <i>S. subsoluta</i> | Italy | Arup L97829 | DQ173238 |
| <i>S. subsoluta</i> | Canada | L89634 (LD) | MT967469 |
| <i>S. subsoluta</i> | Canada | L89634 (LD) | MT967468 |
| <i>S. subsoluta</i> | Ecuador | 11884 (CDS 37243) | MT967467 |

alignment, which was then trimmed and realigned using webPRANK with default settings (Löytynoja & Goldman 2010). On the CIPRES Web Portal (Miller *et al.* 2010), the HYK + G + I model was selected using jModelTest (Posada 2008). A maximum likelihood analysis (ML) was also implemented using

RAXML-HPC2 v. 8.1.11 (Stamatakis 2014) on CIPRES, with 1000 bootstraps for rapid bootstrapping. *Parvoplaca nigroblastidiata* Arup *et al.* (KT161982) was used as outgroup. A tree displaying the phylogeny reconstruction from the ML analysis was generated in FigTree v. 1.4.3 (Rambaut *et al.* 2014).




Fig. 2. *Squamulea chikarensis* (holotype-LAH37546). A, squamulose thallus in natural habitat. B, apothecia. C, transverse section of apothecium showing paraplectenchymatous proper margin and hypothecium. D, ascospore. Scales: A = 1 cm; B = 1.5 mm; C = 100 μ m; D = 10 μ m. In colour online.

Results

Phylogenetic analyses

The final ITS dataset consisted of 35 sequences, representing 12 species of *Squamulea* and *Parvoplaca nigroblastidiata* (see Table 1 for voucher details). The aligned ITS1-5.8S-ITS2 region comprised 565 sites, of which 282 were conserved and 155 variable; 128 sites were parsimony-informative. Our phylogeny recovers *S. chikarensis* in a clade together with *S. loekoesiana* (S.Y. Kondr. & Upreti) Arup *et al.*, (KY614406, KY614407, KY614408, KY614409, KY614410, MK499351), *S. flakusii* (Wilk) Arup *et al.*, (MT967442, MT967444, MN108089, OP341345, OP341346), *S. phyllidizans* (Wetmore) Søchting & Bungartz (MT967456, MT967457) and in a basal position in the clade, *S. osseophila* Søchting & Bungartz (MT967455). This clade contains species from a wide geographical range and diverse substrata, including species with

blastidia, microsquamules and no asexual propagules. Molecular methods are often required in *Squamulea* to distinguish species reliably. The ITS sequence of the holotype of *S. chikarensis* is identical to the sequences of additionally studied specimens (i.e. OP351701, OP351702). Two further newly generated sequences nested within the clade of *S. flakusii* previously known from Ecuador (Galapagos Islands) and Peru. We therefore report *S. flakusii* for the first time from Pakistan and Eurasia.

Taxonomy

Squamulea chikarensis Afshan, Fayyaz, Iftikhar & Khalid sp. nov.

Mycobank No.: MB 845620

The taxon is characterized by a pale green to yellow thallus, large apothecia up to 0.8–1.8 mm wide, pale yellow to yellow-orange

apothecial discs, hymenium 70–110 μm high, large ascospores (12–20 \times 5–11 μm) and a narrow ascospore septum (1.5–3 μm).

Type: Pakistan, Azad Jammu and Kashmir, Chikar, moist temperate forest, 36°23'N, 75°47'E, 3850 m alt., on rock, 2 October 2020, N. S. Afshan, I. Fayyaz & F. Iftikhar (MK-11) (holotype—LAH37546).

(Fig. 2)

Thallus areolate to subsquamulose, areoles/squamules are scattered to crowded, flat to concave, pale green to yellow, slightly pruinose thallus irregular in outline, up to 2–4 cm diam., not delimited by a prothallus. *Cortex* paraplectenchymatous, even, 14–22 μm thick. *Algal layer* continuous, 53–65 μm thick. *Photobiont cells* more or less globose, trebouxoid, 9–15 μm . *Medulla* white, 60–75 μm thick.

Apothecia mostly sparse, erumpent to sessile, circular, (0.8–) 1.0–1.4–1.6(–1.8) mm diam., zeorine. *Disc* concave to flat, pale yellow to yellow-orange, slightly contrasting against thallus, pruinose. *Margin* paler than disc, slightly prominent in young apothecia, then level with disc. *Proper exciple* thin, paraplectenchymatous. *Thalline exciple* thick, slightly/partly reduced, algae abundant, in a continuous layer. *Epihymenium* brownish orange, granular interspersed, K+ purple, 13–18 μm thick. *Hymenium* hyaline, 70–110 μm high. *Paraphyses* simple, septate, 2–3 μm wide, tips up to 5 μm . *Hypothecium* hyaline, not interspersed, paraplectenchymatous. *Asci* 8-spored, clavate, *Teloschistes*-type, 58–65 \times 13–17 μm . *Ascospores* polarilocular, oblong to broadly ellipsoid, (12.0–)13.5–15.7–18.3(–20.0) \times (5.0–)6.2–7.3–9.6(–11.0) μm , septa (1.5–)1.9–2.2–2.7(–3.0) μm wide.

Pycnidia not seen.

Spot test. Thallus and apothecia: K+ reddish brown, C–, KC+ dark red.

Chemistry. Thallus and apothecia with a large proportion of parietin and smaller proportions of teloschistin, fallacinal, parietinic acid and emodin (chemosynndrome A *sensu* Söchting (1997)).

Etymology. The specific epithet '*chikarensis*' (Latin) refers to the type locality Chikar, where the specimen was collected.

Habitat and distribution. The new species was collected in the Himalayas of Pakistan, on siliceous rocks in moist temperate forests, dominated mainly by *Pinus roxburghii*, *Pyrus pashia*, *Quercus oblongata* and *Q. glauca*. The maximum daily temperature of the region varies from 20–30 °C during the summer and averages 4 °C during the winter, with moderate rainfall.

Additional specimens examined. **Pakistan:** Khyber Pakhtunkhwa: Kaghan Valley, moist temperate forest, 34°30'N, 73°18'E, 2500–3000 m alt., on rock, 2021, N. S. Afshan & F. Iftikhar (SG-20) (LAH37548). **Azad Jammu and Kashmir:** Chikar, moist temperate forest, 36°23'N, 75°47'E, 3850 m alt., on rock, 2021, N. S. Afshan, I. Fayyaz & F. Iftikhar (CKT-39) (LAH37547).

Squamulea flakusii (Wilk) Arup, Söchting & Bungartz

In Bungartz, Söchting & Arup, *Plant and Fungal Systematics* **65**, 564 (2020).

(Fig. 3)




Fig. 3. *Squamulea flakusii* (LAH37549). A, squamulose thallus in natural habitat. B, apothecia. Scales: A = 1 cm; B = 1 mm. In colour online.

Thallus squamulose or strongly reduced and almost invisible, 2–3 cm diam., orbicular to irregular in outline, pale yellow to dark yellow, pruinose, squamules flat to convex. *Prothallus* absent. *Cortex* thin to thick (in well-developed thallus), 27–50 μm thick, paraplectenchymatous. *Algal layer* continuous. *Photobiont cells* trebouxoid, more or less globose, 9–14 μm diam.

Apothecia abundant, somewhat crowded and aggregated, rounded, angular, old apothecia distinctly undulate, adnate, zeorine, (0.5–)0.8–1.0–1.2(–1.4) mm wide. *Disc* plane to strongly convex, yellow-orange to pale orange, pruinose, often cracked (especially in old apothecia). *Proper margin* thin, slightly prominent in young apothecia, then level with disc, slightly paler than disc. *Thalline margin* 65–90 μm thick, conspicuous, much reduced, algal cells abundant, forming a continuous layer. *Epihymenium* light brown, 25–38 μm thick. *Hymenium* 60–78 μm thick. *Paraphyses* simple to slightly branched, 2–3 μm broad at base, with upper cells slightly thickened, up to 4 μm . *Hypothecium* 80–140 μm thick, paraplectenchymatous, hyaline. *Asci* 8-spored, clavate, 50–63 \times 12–22 μm . *Ascospores* polarilocular, usually broadly ellipsoid, 12–18 \times 7–11 μm , septum (1.0–)1.3–2.0–2.6(–3.0) μm wide.

Pycnidia not seen.

Spot test. Thallus and apothecia: K+ purple, C–, KC+ purple.

Table 2. A comparison of morphological and anatomical features of *Squamulea chikarensis* sp. nov. with selected *Squamulea* species.

| Characters | <i>S. chikarensis</i> | <i>S. flakusii</i> (Pakistan) | <i>S. flakusii</i> (Peru) | <i>S. flakusii</i> (Galapagos) | <i>S. kiamae</i> | <i>S. loekoesiana</i> | <i>S. osseophila</i> | <i>S. parviloba</i> | <i>S. phyllidizans</i> | <i>S. squamosa</i> | <i>S. subsoluta</i> |
|------------------------------------|------------------------------|----------------------------------|------------------------------|-----------------------------------|-------------------------------|---------------------------------------|---------------------------|---|---|---|---|
| Thallus morphology | areolate to subsquamulose | squamulose or strongly reduced | squamulose | more conspicuously squamulose | rosettes of radiating areoles | areolate | barely subsquamulose | areolate to subsquamulose; areoles with small lobules | subsquamulose to minutely squamulose | squamulose | areolate to subsquamulose |
| Thallus colour | pale green to yellow | pale yellow to dark yellow | orange | deep orange | brownish orange | dull yellow-brownish, yellow-greenish | deep yellow | yellow-orange to orange | yellowish orange | yellow-orange to orange or reddish orange | yellow-orange to orange or reddish orange |
| Apothecium type | zeorine | zeorine | zeorine | - | lecanorine/zeorine | zeorine | lecanorine | zeorine | lecanorine | zeorine | usually biatorine/zeorine |
| Apothecium diam. (mm) | 0.8–1.8 | 0.5–1.4 | 0.2–1.0 | - | 0.3–0.7 | 0.2–0.5 | - | 0.2–0.5 | 0.8 | 0.5–2.0 | 0.1–1.1 |
| Apothecial disc | pale yellow to yellow-orange | yellow-orange to pale orange | reddish | - | reddish to reddish brown | dull yellow to bright yellow | deep orange | darker orange than thallus | deep orange | darker orange than thallus | darker orange than thallus |
| Hymenium (μm) | 70–110 | 60–78 | 70–85 | - | 75–80 | 50–60 | - | 40–70 | - | 65–80 | 60–80 |
| Ascospore size (μm) | 12–20 \times 5–11 | 12–18 \times 7–11 | 10–15 \times 5.0–9.5 | - | 10–16 \times 6–8 | 8.5–12 \times 4.5–6 | 7.6–17.3 \times 4.4–7.4 | 10–14 \times 5.5–7.0 | 8.3–12.8 \times 3.5–8.2 | 9.5–14.0 \times 5.5–7.0 | 9.5–14.0 \times 5.5–7.0 |
| Ascospore septum (μm) | 1.5–3 | 1–3 | 2–4 | - | 1–4 | 4–7 | 1.7–4.1 | 3–4 | 1.7–5.9 | 3–4 | 3–4 |
| Distribution | South Asia (Pakistan) | South Asia (Pakistan) | South America (Peru) | South America (Galapagos) | Australia | Eastern Asia (South Korea) | South America (Galapagos) | south-western North America | south-western North America and South America | south-western North America | North America and Europe |
| References | This paper | This paper | Wilk 2020 | Bungartz et al. 2020 | Kondratyuk et al. 2007 | Kondratyuk et al. 2017b | Bungartz et al. 2020 | Wetmore 2003 | Bungartz et al. 2020 | Wetmore 2003 | Wetmore 2003 |

Specimens examined. **Pakistan:** *Khyber Pakhtunkhwa:* Kaghan Valley, moist temperate forest, 34°30'N, 73°18'E, 2500–3000 m alt., on rock, 2021, N. S. Afshan & F. Iftikhar (SG-45) (LAH37549). *Azad Jammu and Kashmir:* Garhi Dupatta, moist temperate forest, 34°36'N, 73°35'E, 817 m alt., on rock, 2021, N. S. Afshan, I. Fayyaz & F. Iftikhar (CKR-07) (LAH37550).


Discussion

The species *Squamulea chikarensis* is characterized by minute squamules, usually scattered, large apothecia and large ascospores with rather thin septa. The comparison of the new species with its close relatives is presented in Table 2. Morphologically the new species resembles *S. subsoluta*, a widespread species also occurring in Pakistan (under the name *Caloplaca irrubescens*; Aptroot & Iqbal 2012). However, *S. subsoluta* differs from the new species in having a yellow-orange to orange or reddish orange thallus, a blackish prothallus, a darker orange apothecial disc and smaller ascospores, 9.5–14.0 × 5.5–7.0 μm (Wetmore 2003). *Squamulea flakusii*, newly recognized in the lichen biota of Pakistan, differs from *S. chikarensis* by having a pale yellow to dark yellow thallus, apothecia that are 0.5–1.4 mm wide, a hymenium 60–78 μm high and smaller ascospores, 12–18 × 7–11 μm.

In the present study, *Squamulea flakusii* is morpho-anatomically and molecularly characterized, and confirmed as a new record for Pakistan. The morphological features of the Pakistani collections agree with the published description of *S. flakusii* (from South America, Peru) (Wilk 2020), with the exception of the following (features in brackets refer to the Peru collections): less reduced thallus (vs more reduced thallus), pale yellow to dark yellow thallus (vs orange thallus), less crowded and aggregated apothecia (vs more crowded and aggregated apothecia), larger apothecia up to 0.5–1.4 mm wide (vs 0.2–1.0 mm), yellow-orange to pale orange apothecial discs (vs reddish apothecial discs), larger ascospores, 12–18 × 7–11 μm (vs 10–15 × 5.0–9.5 μm) and a narrower ascospore septum, 1–3 μm (vs 2–4 μm). The morphology of Pakistani specimens of *S. flakusii* is, however, quite different to specimens from the Galapagos. Pakistani specimens have a pale yellow to dark yellow thallus (vs deep orange thallus) and are collected at an elevation of 817–3000 m alt. (vs above 1000 m alt. for Galapagos specimens). For a comparison of *S. flakusii* with *S. chikarensis* and other taxa, see Table 2 and the discussion under *S. chikarensis*.

Although *Squamulea flakusii* has been described from South America (Peru) and was then recorded only from the Galapagos Islands (Bungartz *et al.* 2020; Wilk 2020), its distribution in the Northern Hemisphere is possible. This could indicate that the species may be a widespread taxon, as in the case of *S. subsoluta* known also from Austria. To date, due to the uncertainty regarding phenotypic characters, phylogenetic analysis of DNA sequences is almost the only reliable way to distinguish species and to study the geographical distribution of *Squamulea* species (Bungartz *et al.* 2020). Therefore, our knowledge of those species should increase in the future with sampling and analysis of further *Squamulea* species. In Pakistan, *S. flakusii* was collected in the high mountain area, at 817–3000 m alt., in humid forest habitat, on siliceous rock, which more or less corresponds with the previous reports from South America. One Pakistani specimen sequenced (OP341346), however, was collected at c. 817 m alt., in moist temperate forest. The genus is probably more abundantly represented in Pakistan's lichen biota than is currently known.

Acknowledgements. We are very grateful to Dr Ulf Arup (Biological Museum, Lund University, Sweden) for his guidance and to Dr Karina Wilk (IB PAS, Krakow) for her useful comments, both helping to improve the manuscript considerably. We also thank Jason Hollinger (The Edgewood Institute, USA) for reviewing the English and helping us overcome grammatical mistakes. We are indebted to anonymous reviewers whose comments greatly improved the manuscript.

Author ORCIDs.  Najam-ul-Sehar Afshan, 0000-0003-4538-3626; Iram Fayyaz, 0000-0001-6193-1069; Fatima Iftikhar, 0000-0003-4440-3787; Maria Jabben, 0000-0001-9045-9293; Abdul Nasir Khalid, 0000-0002-5635-8031.

References

- Aptroot A and Iqbal SH (2012) Annotated checklist of the lichens of Pakistan, with reports of new records. *Herzogia* 25, 211–229.
- Arup U, Søchting U and Frödén P (2013) A new taxonomy of the family Teloschistaceae. *Nordic Journal of Botany* 31, 016–083.
- Bungartz F, Søchting U and Arup U (2020) Teloschistaceae (lichenized Ascomycota) from the Galapagos Islands: a phylogenetic revision based on morphological, anatomical, chemical, and molecular data. *Plant and Fungal Systematics* 65, 515–576.
- Gardes M and Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2, 113–118.
- Gaya E, Högnabba F, Holguin Á, Molnár K, Fernández-Brime S, Stenroos S, Arup U, Søchting U, van den Boom P, Lücking R, *et al.* (2012) Implementing a cumulative supermatrix approach for a comprehensive phylogenetic study of the Teloschistales (Pezizomycotina, Ascomycota). *Molecular Phylogenetics and Evolution* 63, 374–387.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acid Symposium Series* 41, 95–98.
- Khan M, Khalid AN and Lumbsch HT (2018) A new species of *Lecidea* (Lecanorales, Ascomycota) from Pakistan. *MycKeys* 38, 25–34.
- Kondratyuk SY, Kärnefelt I, Elix JA and Thell A (2007) New species of the genus *Caloplaca* in Australia. *Bibliotheca Lichenologica* 95, 341–386.
- Kondratyuk S, Jeong M-H, Yu N-H, Kärnefelt I, Thell A, Elix JA, Kim J, Kondratyuk AS and Hur J-S (2013) Four new genera of teloschistoid lichens (Teloschistaceae, Ascomycota) based on molecular phylogeny. *Acta Botanica Hungarica* 55, 251–274.
- Kondratyuk SY, Jeong M-H, Yu N-N, Kärnefelt I, Thell A, Elix JA, Kim J, Kondratyuk AS and Hur J-S (2014) A revised taxonomy for the subfamily Caloplacoideae (Teloschistaceae, Ascomycota) based on molecular phylogeny. *Acta Botanica Hungarica* 56, 93–123.
- Kondratyuk SY, Kärnefelt I, Thell A, Elix JA, Kim J, Kondratyuk AS and Hur J-S (2015a) *Tassiloo*, a new genus in the Teloschistaceae (lichenized Ascomycetes). *Graphis Scripta* 27, 22–26.
- Kondratyuk SY, Lökös L, Farkas E, Oh S-O and Hur J-S (2015b) New and noteworthy lichen-forming and lichenicolous fungi 2. *Acta Botanica Hungarica* 57, 77–141.
- Kondratyuk SY, Lökös L, Kim JA, Kondratyuk AS, Jeong M-H, Jang SH, Oh S-O and Hur J-S (2015c) Three new monotypic genera of the caloplacoid lichens (Teloschistaceae, lichen-forming Ascomycetes). *Mycobiology* 43, 195–202.
- Kondratyuk SY, Kärnefelt I, Thell A, Elix JA, Kim J, Kondratyuk AS and Hur J-S (2015d) *Brownlielloideae*, a new subfamily in the Teloschistaceae (Lecanoromycetes, Ascomycota). *Acta Botanica Hungarica* 57, 321–341.
- Kondratyuk SY, Lökös L, Halda JP, Upreti DK, Mishra GK, Haji Moniri M, Farkas E, Park JS, Lee BG, Liu D, *et al.* (2016) New and noteworthy lichen-forming and lichenicolous fungi 5. *Acta Botanica Hungarica* 58, 319–396.
- Kondratyuk SY, Lökös L, Halda JP, Roux C, Upreti DK, Schumm F, Mishra GK, Nayaka S, Farkas E, Park JS, *et al.* (2017a) New and noteworthy lichen-forming and lichenicolous fungi 6. *Acta Botanica Hungarica* 59, 137–260.
- Kondratyuk SY, Lökös L, Upreti DK, Nayaka S, Mishra GK, Ravera S, Jeong M-H, Jang S-H, Park JS and Hur J-S (2017b) New monophyletic branches

- of the *Teloschistaceae* (lichen-forming *Ascomycota*) proved by three gene phylogeny. *Acta Botanica Hungarica* **59**, 71–136.
- Löytynoja A and Goldman N** (2010) webPRANK: a phylogeny-aware multiple sequence aligner with interactive alignment browser. *BMC Bioinformatics* **11**, 1–7.
- Miller MA, Pfeiffer W and Schwartz T** (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 November 2010, New Orleans, Louisiana, pp. 1–8.
- Orange A, James PW and White FJ** (2010) *Microchemical Methods for the Identification of Lichens*. London: British Lichen Society.
- Posada D** (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**, 1253–1256.
- Rambaut A, Suchard MA, Xie D and Drummond AJ** (2014) *FigTree 1.4.2* Institute of Evolutionary Biology, University of Edinburgh. [WWW resource] URL <http://tree.bio.ed.ac.uk/software/figtree/>.
- Sochting U** (1997) Two major anthraquinone chemosyndromes in *Teloschistaceae*. *Bibliotheca Lichenologica* **86**, 135–144.
- Sochting U, Sogaard MZ, Elix JA, Arup U, Elvebakk A and Sancho LG** (2014a) *Catenarina* (*Teloschistaceae*, *Ascomycota*), a new Southern Hemisphere genus with 7-chlorocatenarin. *Lichenologist* **46**, 175–187.
- Sochting U, Garrido-Benavent I, Seppelt R, Castello M, Pérez-Ortega S, De Los Ríos Murillo A, Sancho LG, Frödén P and Arup U** (2014b) *Charcotiana* and *Amundsenia*, two new genera in *Teloschistaceae* (lichenized *Ascomycota*, subfamily *Xanthorioideae*) hosting two new species from continental Antarctica, and *Austroplaca frigida*, a new name for a continental Antarctic species. *Lichenologist* **46**, 763–782.
- Stamatakis A** (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Wetmore CM** (2003) The *Caloplaca squamosa* group in North and Central America. *Bryologist* **106**, 147–156.
- White TJ, Bruns TD, Lee SB and Taylor JW** (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In Innis MA, Gelfand DH, Sninsky JJ and White TJ (eds), *PCR Protocols: a Guide to Methods and Applications*. San Diego: Academic Press, pp. 315–322.
- Wilk K** (2020) *Huriella flakusii* (*Teloschistaceae*, lichenized *Ascomycota*), a new species from the Colca Canyon region in Peru. *Lichenologist* **52**, 37–47.
- Wilk K, Pabijan M, Saluga M, Gaya E and Lücking R** (2021) Phylogenetic revision of South American *Teloschistaceae* (lichenized *Ascomycota*, *Teloschistales*) reveals three new genera and species. *Mycologia* **113**, 278–299.