

EPP0689

Body satisfaction and sexuality in pregnant and postpartum women

S. Bader^{1*}, M. aloulou¹, Z. zran¹, A. Abdelmoula², A. Bouaziz¹ and W. abbes¹

¹Psychiatry and ²obstetric gynecology, University Hospital of Gabes, Gabes, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.772

Introduction: Pregnancy and postpartum is an important life event associated with profound physiognomic and psychosocial changes affecting the female body in all its physiological, psychic and affective reality. It might influence the sexual function in expectant mothers.

Objectives: To investigate the relationship between the body satisfaction and perception and the sexual function among pregnant and postpartum women.

Methods: It was a cross-sectional study established over a period of 3 months from the June 1st, 2023 to August 31, 2023. This study focused on a population of pregnant and postpartum women recruited from outpatient consultations and inpatient of the obstetric gynecology department at the university hospital of Gabes. We used a pre-established sheet exploring socio-demographic data, medical and gynecologic history, the body mass index (BMI) and informations concerning the marital relationship and the woman's sexual activity. We administered the validated Arabic version of the Arizona Sexual Experiences Scale (ASEX) to assess sexual functioning and we used the body satisfaction and global self-perception questionnaire (QSCPGS) to explore the body satisfaction and perception.

Results: Fifty-eight women were included. The average age was 35.6±5.5 years; they were from an urban origin in 75%. They were pregnant in the first, second and third trimester in (15.6%, 15.6% and 25% respectively). They were in postpartum in 43.8% of cases with a cesarean delivery in 73.3% and breastfeeding in 56%. All women reported being on good terms with their spouses and satisfied with their sexuality. The usual frequency of sexual relations was (1/day: 22.6%, 1/week: 74.2%, 1/month: 3.2%) and 25% reported wanting to reduce the frequency. The mean ASEX score was 13 ± 4.3 and 47% of the sample had sexual dysfunction. For the total score of the QSCPGS, we observe a mean value of 33 ± 28.3, which means that our sample has a good level of positive body satisfaction and self-perception. The mean value of the "body satisfaction" factor is higher (23.7 ± 10.4) than the mean value of the "self-perception" factor (11.4 ± 14.3). The mean value of BMI was 28.74 ± 4.4 which means an overweight. We found a significant association between the "body satisfaction" factor and the sexual dysfunction (p=0.03), insufficient lubrication (p=0.01) and difficulty reaching orgasm (p=0.001).

Conclusions: We found that body and physical changes among pregnant and postpartum women can negatively affect their body perception and it might deteriorate its global sexual function. Further researches are recommended to study other potential factors affecting sexual function during this period.

Disclosure of Interest: None Declared

E-mental Health

EPP0688

Development of a novel screening questionnaire for brain fog

Y. Jheng^{1*}, W.-C. Hsieh² and Y.-H. Chou^{1,2,3}

¹Center for Quality Management; ²Department of Psychiatry, Taipei Veterans General Hospital and ³Department of Psychiatry, School of Medicine, National Yang-Ming Chiao-Tung University, Taipei, Taiwan

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.773

Introduction: Amidst the widespread proliferation of the COVID-19 virus, brain fog has become one of the most critical issues of public health. Brain fog may lead to sub-health conditions, such as forgetfulness, difficulty thinking, and other related symptoms. Although they are not immediately life-threatening, these sub-health conditions could gradually erode the quality of life. Currently, there is no relevant screening tool for brain fog.

Objectives: The aim of this study was to develop a reliable screening tool.

Methods: A web-based brain fog screening questionnaire was developed in the study. It was based on previous studies, which summarized five parts of the most common clinical symptoms after COVID-19: forgetfulness, difficulty thinking, difficulty concentrating, feeling confused, and difficulty finding words or phrases to speak. Unfortunately, these items were used only in a way of yes or no answers in previous studies. Each of these items was expanded to five anchors to evaluate their severity in the study. Cronbach's alpha coefficient was used to assess internal consistency. K-means clustering was used as a second method to validate the cutoff points. Furthermore, the receiver operating characteristic (ROC) curve was applied to validate the appropriateness of the cutoff point.

Results: There were 534 participants who completely finished the questionnaire. It includes 183 males and 351 females, and all of them aged between 19 and 81 years. The Cronbach's Alpha value was 0.821. The cutoff point was at a total score of 6 in terms of K-means. Based on the result, the ROC curve revealed that an area under the curve (AUC) was 0.816 with a confidence interval of 0.784 to 0.849.

Conclusions: The study demonstrated the feasibility and reliability of the web-based screening test for brain fog.

Disclosure of Interest: None Declared

EPP0689

Efficacy of Digital Interventions for Anxiety Disorders: A Systematic Review and Meta-Analysis

H.-G. Ji^{1*}, S. Kang¹, H. Bae¹, G. Kim¹ and J.-W. Hur¹

¹Department of Psychology, Korea University, Seoul, Korea, Republic Of

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.774

Introduction: Anxiety disorders are one of the most common mental disorders, yet only less than 20% of people with anxiety disorders receive adequate treatment. Digital interventions for anxiety disorders can potentially increase access to evidence-based treatment. However, there is no comprehensive meta-analysis study that covers all modalities of digital interventions and all anxiety disorders.

Objectives: A preliminary meta-analysis was conducted to examine the treatment efficacy of digital interventions [e.g., virtual reality (VR)-, mobile application-, internet-based interventions] for anxiety disorders and to identify potential moderators that may lead to better treatment outcomes.

Methods: We searched Embase, PubMed, PsycINFO, Web of Science, and the Cochrane Library for randomized controlled trials examining the therapeutic efficacy of digital interventions for individuals with anxiety disorders from database inception to April 18, 2023. Search keywords were developed by combining the PICOS framework and MeSH terms. Data screening and extraction adhered to PRISMA guidelines. We used a random-effects model with effect sizes expressed as Hedge's *g*. The quality of the studies was assessed using the Revised Cochrane risk-of-bias tool for randomized trials (RoB 2). The study protocol was registered in PROSPERO on April 22, 2023 (CRD42023412139).

Results: A systematic literature search identified 19 studies with randomized controlled trials (21 comparisons; 1936 participants) with high overall heterogeneity ($Q = 104.49$; $P < .001$; $I^2 = 80.9\%$). Digital interventions reduced anxiety symptoms with medium to large effect sizes ($g = 0.78$; 95% CI: 0.55-1.02; $P < .001$), with interventions for specific phobia showing the largest effect size ($n = 6$; $g = 1.22$; 95% CI: 0.51-1.93; $P < .001$). VR-based interventions had a larger effect size ($n = 6$; $g = 0.98$; 95% CI: 0.39-1.57; $P < .001$) than mobile- or internet-based interventions, which had medium effect sizes. Meta-regression results exhibited that effect sizes of digital interventions were associated with the mean age of participants ($\beta = 0.04$; 95% CI: 0.02-0.06; $P < .001$).

Conclusions: The results of this study provide evidence for the efficacy of digital interventions for anxiety disorders. However, this also suggests that the degrees of effectiveness in reducing anxiety symptoms can be moderated by the specific diagnosis, the modalities of digital technologies, and mean age, implying that the application of digital interventions for anxiety disorders should be accompanied by personalized guidance.

Disclosure of Interest: None Declared

Sexual Medicine and Mental Health

EPP0690

Risk assessment and treatment - Evaluation of a group therapy for people with pedophilia

P. Heindl*, S. Schobel, K. Fischer, T. Nenov-Matt, A. Chrobok, M. Wertz and K. Schiltz

Forensic Department, LMU Klinikum, Munich, Germany

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.775

Introduction: Deviant sexual interest for children (pedophilia, hebephilia) is associated with a higher risk of sexual offending against children (CSA) and consuming child sexual abuse images

(CSAI). There is a general shortage of therapeutic programs for individuals who feel sexually attracted to juvenile bodies and are concerned about their sexual behaviour. Efforts to establish regional centres throughout Germany offering preventive support led to the prevention network "Don't become an offender" ("Kein Täter werden").

Objectives: To identify dynamic risk factors (DRFs) and evaluate a treatment programme aiming to reduce CSA and CSAI among potential or existing pedosexual offenders (who have not been legally charged). In addition, changes in the course of therapy are examined to provide information about the accessibility and motivation of the target group and its therapeutic responsiveness.

Methods: Participants undergo standardized diagnostic and treatment procedures. Therapy comprises an outpatient psychotherapy program (group therapy) over the course of approx. 48 weekly sessions, optional individual and partner/relative including sessions, as well as additional pharmaceutical treatment. Assessments are carried out through self- and other-reported psychometric test batteries pre-, during and post-treatment up to a 3.5 year follow-up. The test battery includes clinical questionnaires (WHO-5, CTQ-SF), personality questionnaires (ISK-K, NEO-FFI), sexuality questionnaires (EKK-R, KV-M, MSI, HBI-19) and risk assessment procedures (VRAG-R, STATIC-99, VRS:SO). Main outcome measures are self- and externally-reported DRF changes well as offending behaviour characteristics.

Results: By September 20, 2023, N=12 individuals were enrolled in the treatment program. All individuals had a deviant sexual preference (exclusive/non-exclusive pedo-/hebephilia). Nine individuals reported past and/or current use of CSAI. Of these, two individuals reported at least one CSA in the past. Three had no previous use of CSAI or CSA history.

In the first treatment group (N=6), preliminary results show reduction in dynamic risk factors (e.g., Cognitive Bias, Sexual Compulsivity, Impulsivity) after the first 12 weeks of treatment. The evaluation of additional clinical data is pending.

Conclusions: To date, therapy for individuals with pedophilia or hebephilia has been insufficient – particularly when not offending. Ongoing evaluation of the therapy program should provide further insight into responsiveness and therapeutic motivation of this target group. In particular, the impact of therapy on changing dynamic risk factors for CSA and CSAI remains to be examined.

Disclosure of Interest: None Declared

EPP0691

Psychological Background of Sexual Dysfunctions – a Comparative Study on Hungarian and Spanish Samples

Z. G. Pintér-Eszenyei^{1*}, S. Kató¹, Á. Hőgye-Nagy¹, A. K. Csinády¹, M. A. Rando Hurtado² and A. Szemán-Nagy¹

¹Faculty of Humanities, Institute of Psychology, University of Debrecen, Debrecen, Hungary and ²Faculty of Psychology and Speech Therapy, University of Málaga, Málaga, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.776

Introduction: Sexual dysfunctions are prevalent issues affecting individuals' sexual well-being and relationships. These conditions encompass a range of difficulties in sexual functioning, from desire