

FINITENESS OF CANONICAL QUOTIENTS OF DEHN QUANDLES OF SURFACES

NEERAJ K. DHANWANI and MAHENDER SINGH

(Received 9 October 2023; accepted 28 January 2024)

Communicated by Ben Martin

Abstract

The Dehn quandle of a closed orientable surface is the set of isotopy classes of nonseparating simple closed curves with a natural quandle structure arising from Dehn twists. In this paper, we consider the finiteness of some canonical quotients of these quandles. For a surface of positive genus, we give a precise description of the 2-quandle of its Dehn quandle. Further, with some exceptions for genus more than 2, we determine all values of n for which the n -quandle of its Dehn quandle is finite. The result can be thought of as the Dehn quandle analogue of a similar result of Hoste and Shanahan for link quandles [‘Links with finite n -quandles’, *Algebr. Geom. Topol.* **17**(5) (2017), 2807–2823]. We also compute the size of the smallest nontrivial quandle quotient of the Dehn quandle of a surface. Along the way, we prove that the involutory quotient of an Artin quandle is precisely the corresponding Coxeter quandle, and also determine the smallest nontrivial quotient of a braid quandle.

2020 *Mathematics subject classification*: primary 57K10, 57K20; secondary 57K12.

Keywords and phrases: Artin group, Coxeter quandle, Dehn quandle, involutory quandle, orientable surface, mapping class group.

1. Introduction

The Dehn quandle of a group with respect to a subset is the union of conjugacy classes of elements of the set equipped with the operation of conjugation. It turns out that these are precisely the quandles that embed into their enveloping groups, and many well-known classes of quandles satisfy this property [9]. The first is the class of free quandles, where a free quandle on a set is simply the union of conjugacy classes of the free generating set in the corresponding free group. The second interesting class of examples, which is also the focus of this paper, is given by surfaces. Let \mathcal{M}_g be the mapping class group of a closed orientable surface S_g of genus $g \geq 1$, and \mathcal{D}_g^{ns} the

N.K.D. acknowledges support from the NBHM via grant number 0204/1/2023/R &D-II/606. M.S. is supported by the Swarna Jayanti Fellowship grants DST/SJF/MSA-02/2018-19 and SB/SJF/2019-20/04.

© The Author(s), 2024. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

set of isotopy classes of nonseparating simple closed curves on S_g . It is well known that \mathcal{M}_g is generated by Dehn twists along finitely many simple closed curves from \mathcal{D}_g^{ns} [12, Theorem 4.1]. The binary operation $\alpha * \beta = \tau_\beta(\alpha)$, where $\alpha, \beta \in \mathcal{D}_g^{ns}$ and τ_β is the Dehn twist along β , turns the set \mathcal{D}_g^{ns} into a quandle called the Dehn quandle of the surface S_g . In fact, \mathcal{D}_g^{ns} can be viewed as a subquandle of the conjugation quandle of \mathcal{M}_g , by identifying the isotopy class of a nonseparating simple closed curve with the isotopy class of its corresponding Dehn twist. These quandles originally appeared in the work of Zablow [29, 30]. It turns out that two surfaces of genus more than two are homeomorphic if and only if their Dehn quandles are isomorphic [9, Proposition 6.5]. A homology theory based on Dehn quandles of surfaces has been derived in [31] and it has been shown that isomorphism classes of Lefschetz fibrations over a disk correspond to quandle homology classes in dimension two. In a related direction, the papers [19, 27, 28] considered a quandle structure on the set of isotopy classes of simple closed arcs on an orientable surface with at least two punctures, referred as the quandle of cords. In the case of a disk with n punctures, this quandle is simply the Dehn quandle of the braid group B_n on n strands with respect to its standard set of generators. It turns out that Dehn quandles of groups with respect to their subsets include many well-known constructions of quandles from groups including conjugation quandles of groups, free quandles, Coxeter quandles, Dehn quandles of closed orientable surfaces, quandles of cords of orientable surfaces, knot quandles of prime knots, core quandles of groups and generalised Alexander quandles of groups with respect to fixed-point free automorphisms (see [9] for more details). A presentation for the quandle of cords of the plane and the 2-sphere has been given in [19]. Beyond these cases, not much was known about presentations of Dehn quandles until the work [10], wherein two approaches to writing efficient presentations for Dehn quandles using presentations of their underlying groups have been given. Several computations of presentations have been given, including Dehn quandles of spherical Artin groups, surface groups and mapping class groups of orientable surfaces for small genera.

Since link quandles are always infinite except for the unknot and the Hopf link, it is natural to explore finiteness of n -quandles of link quandles. It had been conjectured by Przytycki that the n -quandle of an oriented link L in the 3-sphere is finite if and only if the fundamental group of the n -fold cyclic branched cover of the 3-sphere, branched over L , is finite. The conjecture has been proved by Hoste and Shanahan in [14], wherein they use Dunbar's classification [11] of spherical 3-orbifolds to determine all links with a finite n -quandle for some n . In this paper, we consider the analogous problem for Dehn quandles of surfaces. The two families of quandles have a rather curious intersection. Thanks to the work of Niebrzydowski and Przytycki [23], where they proved that the knot quandle of the trefoil is isomorphic to the Dehn quandle of the torus, the problem can be thought of as a quandle analogue of [5, Problem 28], which asks whether the normal closure of squares of Dehn twists in the mapping class group of a closed orientable surface is of finite index. See also [13, 16] for further work on the problem.

The paper is organised as follows. In Section 3, we establish some general results on finiteness of n -quandles of Dehn quandles. We prove that if the n -quandle of the Dehn quandle of a group is finite, then a certain canonical quotient of the group is finite (Theorem 3.4). We consider Dehn quandles of Artin groups in Section 4 and prove that the involutory quandle of an Artin quandle is the corresponding Coxeter quandle (Theorem 4.3). We also determine the size of the smallest nontrivial quandle quotient of the Dehn quandle of the braid group with respect to its standard generating set (Proposition 4.5). In Section 5, we discuss finiteness of n -quandles of Dehn quandles of surfaces. For a closed orientable surface of positive genus, we give a precise description of its 2-quandle (Theorem 5.4). Further, with some exceptions for genus more than two, we determine all values of n for which the n -quandle of the Dehn quandle is finite (Theorem 5.6). As a final result, we also determine the size of the smallest nontrivial quandle quotient of the Dehn quandle of a surface (Proposition 5.8).

2. Preliminaries

Throughout the paper, we consider right-distributive quandles. Recall that, a *quandle* is a nonempty set Q equipped with a binary operation $*$ satisfying the following axioms:

- (i) $x * x = x$ for all $x \in Q$;
- (ii) for each $x, y \in Q$, there exists a unique $z \in Q$ such that $x = z * y$;
- (iii) $(x * y) * z = (x * z) * (y * z)$ for all $x, y, z \in Q$.

The second quandle axiom implies that there exists a dual binary operation $*^{-1}$ on Q such that

$$(x * y) *^{-1} y = x = (x *^{-1} y) * y$$

for all $x, y \in Q$. Topologically, the three quandle axioms correspond to the three Reidemeister moves of planar diagrams of links in the 3-space. Though quandles are ubiquitous, groups and knots are prominent sources of quandles. For instance, if G is a group, then the binary operation $x * y = yxy^{-1}$ turns G into the quandle called the *conjugation quandle* of G . Concerning knot theory, every link can be assigned a quandle called the *link quandle*, which is a complete invariant of nonsplit links up to weak equivalence. This fundamental result appeared independently in the works of Joyce [17, 18] and Matveev [21], and has led to much of the recent work on the subject.

Morphisms and automorphisms of quandles are defined in the obvious way. Note that the quandle axioms are equivalent to saying that for each $y \in Q$, the map $S_y : Q \rightarrow Q$ given by $S_y(x) = x * y$ is an automorphism of Q fixing y . The group $\text{Inn}(Q)$ generated by such automorphisms is called the group of inner automorphisms of Q . The group $\text{Inn}(Q)$ acts on the quandle Q and the corresponding orbits are referred as connected components. For example, knot quandles are connected, whereas link quandles of split links are not connected.

The *enveloping group* $\text{Env}(Q)$ of a quandle Q is the group with the set of generators as $\{e_x \mid x \in Q\}$ and the defining relations as

$$e_{x*y} = e_y e_x e_y^{-1}$$

for all $x, y \in Q$. For example, the enveloping group of the link quandle of a link in the 3-space is the corresponding link group [17, 18]. The natural map

$$\eta : Q \rightarrow \text{Env}(Q)$$

given by $\eta(x) = e_x$ is a quandle homomorphism with $\text{Env}(Q)$ viewed as the conjugation quandle. The map η is not injective in general. In fact, Dehn quandles are precisely those for which this map is injective [9, Proposition 3.9]. The functor from the category of quandles to that of groups assigning the enveloping group to a quandle is left adjoint to the functor from the category of groups to that of quandles assigning the conjugation quandle to a group. Thus, enveloping groups play a crucial role in the understanding of quandles themselves.

Using defining axioms [26, Lemma 4.4.7], any element of a quandle Q can be written in a left-associated product of the form

$$((\dots((a_0 *^{\epsilon_1} a_1) *^{\epsilon_2} a_2) *^{\epsilon_3} \dots) *^{\epsilon_{n-1}} a_{n-1}) *^{\epsilon_n} a_n,$$

where $a_i \in Q$ and $\epsilon_i \in \{1, -1\}$. For simplicity, we write the preceding expression as

$$a_0 *^{\epsilon_1} a_1 *^{\epsilon_2} \dots *^{\epsilon_n} a_n.$$

Let $n \geq 2$ be an integer. A quandle Q is called an *n-quandle* if

$$x * \underbrace{y * y * \dots * y}_{n \text{ times}} = x$$

for all $x, y \in Q$. Equivalently, a quandle is an *n-quandle* if and only if each basic inner automorphism S_x has order dividing n . A 2-quandle is also called *involutory*. For example, the core quandle of any group is involutory.

3. Finiteness of n-quandles of Dehn quandles

Let G be a group, A a nonempty subset of G , and A^G the set of all conjugates of elements of A in G . The *Dehn quandle* $\mathcal{D}(A^G)$ of G with respect to A is defined as the set A^G equipped with the binary operation of conjugation, that is,

$$x * y = yxy^{-1}$$

for all $x, y \in \mathcal{D}(A^G)$. The class of Dehn quandles contains many well-known constructions of quandles from groups. Note that $\mathcal{D}(G^G)$ is the conjugation quandle of G . If $F(S)$ is the free group generated by S , then $\mathcal{D}(S^{F(S)})$ is the free quandle on S . If (W, S) is a Coxeter system, then $\mathcal{D}(S^W)$ is the so called Coxeter quandle [2, 24]. Let S_g be a closed orientable surface of genus g and \mathcal{M}_g its mapping class group. If S is the set

of Dehn twists about nonseparating simple closed curves, then $\mathcal{D}(S^M_g)$ is the Dehn quandle of the surface [19, 29, 30].

Given a quandle Q and an integer $n \geq 2$, the n -quandle $(Q)_n$ of Q is defined as the quotient of Q by the relations

$$x *^n y := x * \underbrace{y * y * \cdots * y}_{n \text{ times}} = x$$

for all $x, y \in Q$. Note that there is a natural epimorphism $\pi : Q \rightarrow (Q)_n$. Further, the construction satisfies the universal property that for any quandle homomorphism $f : Q \rightarrow Y$, where Y is an n -quandle, there exists a unique quandle homomorphism $\bar{f} : (Q)_n \rightarrow Y$ such that $\bar{f} \pi = f$. The construction is of particular interest from the point of view of knot theory. Since link quandles are always infinite except for the unknot and the Hopf link, it is interesting to explore finiteness of n -quandles of link quandles. Using Thurston’s geometrisation theorem and Dunbar’s classification of spherical 3-orbifolds, a complete classification of links whose n -quandles are finite for some n has been given in [14].

We begin with the following general observation.

PROPOSITION 3.1. *Let Q be a quandle with a presentation $\langle S \mid R \rangle$. Then, for each $n \geq 2$, $(Q)_n$ has a presentation $\langle S \mid R \cup T \rangle$, where $T = \{x *^n y = x \mid x, y \in S\}$.*

PROOF. Let $n \geq 2$ and X be the quandle with presentation $\langle S \mid R \cup T \rangle$. First, we claim that X is an n -quandle. Consider elements $x \in X$ and $y \in S$. We write $x = a_1 *^{\epsilon_1} a_2 *^{\epsilon_2} \cdots *^{\epsilon_{l-1}} a_l$, where $a_i \in S$ and $\epsilon_i \in \{1, -1\}$. By right distributivity, we obtain

$$x * y = (a_1 * y) *^{\epsilon_1} (a_2 * y) *^{\epsilon_2} \cdots *^{\epsilon_{l-1}} (a_l * y)$$

and a repeated application of the property further gives

$$x *^n y = (a_1 *^n y) *^{\epsilon_1} (a_2 *^n y) *^{\epsilon_2} \cdots *^{\epsilon_{l-1}} (a_l *^n y) = a_1 *^{\epsilon_1} a_2 *^{\epsilon_2} \cdots *^{\epsilon_{l-1}} a_l = x,$$

which is desired.

Now take $x, y \in X$, and use induction on the word length of y . When y is of word length one, we are in the preceding case. We assume that $y = a_1 *^{\epsilon_1} a_2 *^{\epsilon_2} \cdots *^{\epsilon_{k-1}} a_k$, where $k > 1$. Then we write $y = y_1 *^{\epsilon_{k-1}} a_k$, where $y_1 := a_1 *^{\epsilon_1} a_2 *^{\epsilon_2} \cdots *^{\epsilon_{k-2}} a_{k-1}$ is of word length less than k . By the induction hypothesis, the assertion holds for all words of length less than k . Then, using [26, Lemma 4.4.7],

$$\begin{aligned} x *^n y &= x *^n (y_1 *^{\epsilon_{k-1}} a_k) \\ &= (x *^{-\epsilon_{k-1}} a_k * y_1 *^{\epsilon_{k-1}} a_k) *^{n-1} (y_1 *^{\epsilon_{k-1}} a_k) \\ &= (x *^{-\epsilon_{k-1}} a_k *^2 y_1 *^{\epsilon_{k-1}} a_k) *^{n-2} (y_1 *^{\epsilon_{k-1}} a_k) \\ &= ((x *^{-\epsilon_{k-1}} a_k) *^n y_1) *^{\epsilon_{k-1}} a_k \\ &= (x *^{-\epsilon_{k-1}} a_k) *^{\epsilon_{k-1}} a_k, \quad \text{by induction hypothesis} \\ &= x. \end{aligned}$$

Thus, X is an n -quandle, proving our claim. Finally, it follows from the universal property that $(Q)_n \cong X$. □

COROLLARY 3.2. *If the n -quandle of a quandle is finite for some integer $n \geq 2$, then its d -quandle is also finite for each divisor d of n .*

The presentation of the enveloping group of a quandle can be reduced substantially [26, Theorem 5.1.7].

COROLLARY 3.3. *Let $Q = \langle S \mid R \rangle$ be a quandle presentation. Then*

$$\text{Env}((Q)_n) = \langle \tilde{S} \mid \tilde{R} \cup \tilde{T} \rangle \cong \text{Env}(Q) / \langle\langle \tilde{T} \rangle\rangle,$$

where $\tilde{S} = \{e_x \mid x \in S\}$, $\tilde{T} = \{e_x^n e_y e_x^{-n} e_y^{-1} \mid x, y \in S\}$ and \tilde{R} consists of relations of R with each $x * y$ replaced by $e_y e_x e_y^{-1}$.

Given a quandle Q , define a map $\nu : Q \rightarrow \mathbb{N} \cup \{0\}$ by setting

$$\nu(x) = \begin{cases} \text{order of } S_x & \text{if order of } S_x \text{ is finite,} \\ 0 & \text{if order of } S_x \text{ is infinite.} \end{cases}$$

Consider the subgroup $Z(Q) = \langle e_x^{\nu(x)} \mid x \in Q \rangle$ of $\text{Env}(Q)$. It follows from the construction of $Z(Q)$ that it is a central subgroup of $\text{Env}(Q)$ [3, Lemma 2.1]. Let $F(Q) = \text{Env}(Q)/Z(Q)$ be the corresponding quotient group. It is known that if Q is a finite quandle, then $F(Q)$ is a finite group [3, Proposition 3.1].

THEOREM 3.4. *Let G be a group generated by a set S and $G_n = G / \langle\langle s^n \mid s \in S \rangle\rangle$. Then there exists a surjective group homomorphism $F((\mathcal{D}(S^G))_n) \rightarrow G_n$. In particular, if $(\mathcal{D}(S^G))_n$ is finite, then G_n is finite.*

PROOF. It follows from [9, Theorem 3.6] that the map $\Phi : \text{Env}(\mathcal{D}(S^G)) \rightarrow G$, given by $\Phi(e_x) = x$, is a surjective homomorphism. By Corollary 3.3, we have a surjective homomorphism $\tilde{\Phi} : \text{Env}((\mathcal{D}(S^G))_n) \rightarrow H_n$, where $H_n = G / \langle\langle x^n y x^{-n} y^{-1} \mid x, y \in S \rangle\rangle$. This, in turn, gives a surjective homomorphism

$$\bar{\Phi} : F((\mathcal{D}(S^G))_n) \rightarrow H_n / \langle\langle x^n \mid x \in S \rangle\rangle.$$

The assertion now follows from the fact that $H_n / \langle\langle x^n \mid x \in S \rangle\rangle \cong G_n$. □

4. Involutory quotients of Artin quandles

Recall that an Artin group \mathbf{A} is a group with a presentation

$$\mathbf{A} = \{s_1, s_2, \dots, s_n \mid (s_i s_j)_{m_{ij}} = (s_j s_i)_{m_{ij}}, \text{ where } m_{ij} \in \{2, 3, \dots\} \cup \{\infty\}\},$$

where $(s_i s_j)_{m_{ij}}$ is the word $s_i s_j s_i s_j s_i \cdots$ of length m_{ij} if $m_{ij} < \infty$ and there is no relation of type $(s_i s_j)_{m_{ij}} = (s_j s_i)_{m_{ij}}$ if $m_{ij} = \infty$. We set $\mathbf{S} = \{s_1, s_2, \dots, s_n\}$. The corresponding Coxeter group W is the quotient of \mathbf{A} by imposing additional relations $s_i^2 = 1$ for all $s_i \in \mathbf{S}$.

To distinguish the presentation of \mathbf{A} from that of its corresponding Coxeter group W , we write the presentation of W without using bold letters. More precisely,

$$W = \{s_1, s_2, \dots, s_n \mid s_i^2 = 1, (s_i s_j)_{m_{ij}} = (s_j s_i)_{m_{ij}}, \text{ where } m_{ij} \in \{2, 3, \dots\} \cup \{\infty\}\}.$$

Setting $S = \{s_1, s_2, \dots, s_n\}$, the pair (W, S) is referred to as a Coxeter system and elements of W that are conjugates of elements of S are called reflections. Following [9], Dehn quandles of Artin and Coxeter groups with respect to their standard generating sets are referred to as Artin and Coxeter quandles, respectively. In this section, we prove that the involutory quandle of an Artin quandle is the corresponding Coxeter quandle.

To prove the result, we must first understand centralisers of Coxeter generators, for which we follow [4]. Let (W, S) be a Coxeter system and Δ the labelled graph with vertex set S such that there is an edge between s and s' with label $m_{ss'}$ whenever $m_{ss'} < \infty$. Let Δ^{odd} be the subgraph of Δ consisting of only odd labelled edges. It is easy to see that connected components of Δ^{odd} correspond to conjugacy classes of reflections in W . For each $s \in S$, let Δ_s^{odd} denote the connected component of Δ^{odd} containing s .

Let (W, S) be a Coxeter system and $\gamma = (t_0, t_1, \dots, t_n)$ an edge-path in Δ^{odd} with the edge joining t_{i-1} and t_i labelled $2l_i + 1$. We set

$$p_\gamma := (t_1 t_0)^{l_1} (t_2 t_1)^{l_2} \cdots (t_n t_{n-1})^{l_n}$$

if $n > 0$ and $p_\gamma = 1$ if $n = 0$. If u is a vertex of Δ such that there is an edge joining u and t_n with even label 2λ , then we define

$$r_{\gamma,u} := p_\gamma (u t_n)^{\lambda-1} u p_\gamma^{-1}.$$

Henceforth, the notation $r_{\gamma,u}$ means that the vertex u is joined to the end point of γ by an even labelled edge. We need the following results [4, Corollaries 6 and 8]. See also [6, 7, 15].

THEOREM 4.1. *Let (W, S) be a Coxeter system, $s \in S$ and $C_W(s)$ the centraliser of s in W .*

- (1) *Let W_Ω be the subgroup of $C_W(s)$ generated by all the reflections it contains except s . Let Γ_Ω be the subgroup of W generated by elements p_γ , where γ is an edge-loop in Δ^{odd} based at s . Then $C_W(s) = \langle s \rangle \times (W_\Omega \rtimes \Gamma_\Omega)$.*
- (2) *Let Z be a set of edge-loops in Δ_s^{odd} generating the fundamental group $\pi_1(\Delta^{\text{odd}}, s)$. Then $\{p_\gamma \mid \gamma \in Z\}$ generates Γ_Ω .*
- (3) *Let Y be the set consisting of one edge-path δ_t in Δ_s^{odd} from s to t for each vertex $t \in \Delta_s^{\text{odd}}$, and let X be the set of vertices u of Δ such that there is an even labelled edge joining u to the end point of some δ_t . Then $\{p_\gamma \mid \gamma \in Z\}$ and $\{r_{\delta_t,u} \mid \delta_t \in Y, u \in X\}$ together generate $W_\Omega \rtimes \Gamma_\Omega$.*

Let $\nu : \mathbf{A} \rightarrow W$ be the natural surjection given by $\nu(\mathbf{s}) = s$. Clearly, $\ker(\nu)$ is normally generated by $\{\mathbf{s}^2 \mid \mathbf{s} \in \mathbf{S}\}$.

LEMMA 4.2. *The restriction map $C_{\mathbf{A}}(\mathbf{s}) \rightarrow C_W(s)$ is surjective for each $\mathbf{s} \in \mathbf{S}$.*

PROOF. Let $\mathbf{s} \in \mathbf{S}$ be fixed such that $\nu(\mathbf{s}) = s$. Let $\gamma = (t_0, t_1, \dots, t_n)$ be an edge-path in Δ^{odd} with the edge joining t_{i-1} and t_i labelled $2i + 1$, and let $p_\gamma = (t_1 t_0)^{l_1} (t_2 t_1)^{l_2} \cdots (t_n t_{n-1})^{l_n}$ be the corresponding element. Let $\mathbf{p}_\gamma = (\mathbf{t}_1 \mathbf{t}_0)^{l_1} (\mathbf{t}_2 \mathbf{t}_1)^{l_2} \cdots (\mathbf{t}_n \mathbf{t}_{n-1})^{l_n}$ denote the lift of p_γ in \mathbf{A} . If we set $\bar{\mathbf{p}}_\gamma = (\mathbf{t}_{n-1} \mathbf{t}_n)^{l_n} (\mathbf{t}_{n-2} \mathbf{t}_{n-1})^{l_{n-1}} \cdots (\mathbf{t}_0 \mathbf{t}_1)^{l_1}$, then it follows that $\nu(\bar{\mathbf{p}}_\gamma) = p_\gamma^{-1}$. It is easy to see that the identities

$$\mathbf{p}_\gamma \mathbf{t}_n = \mathbf{t}_0 \mathbf{p}_\gamma \quad \text{and} \quad \bar{\mathbf{p}}_\gamma \mathbf{t}_0 = \mathbf{t}_n \bar{\mathbf{p}}_\gamma$$

hold in \mathbf{A} . Thus, if γ is an edge loop based at s , then both \mathbf{p}_γ and $\bar{\mathbf{p}}_\gamma$ commute with \mathbf{s} . Similarly, let δ_t be a edge-path in Δ_s^{odd} from s to t and u a vertex of Δ that is joined to the end point of δ_t with an edge labelled 2λ . If we set $\mathbf{r}_{\delta_t, u} = \mathbf{p}_{\delta_t} (\mathbf{u} \mathbf{t})^{\lambda-1} \mathbf{u} \bar{\mathbf{p}}_{\delta_t}$, then we can see that

$$\begin{aligned} \mathbf{r}_{\delta_t, u} \mathbf{s} &= \mathbf{p}_{\delta_t} (\mathbf{u} \mathbf{t})^{\lambda-1} \mathbf{u} \bar{\mathbf{p}}_{\delta_t} \mathbf{s} \\ &= \mathbf{p}_{\delta_t} (\mathbf{u} \mathbf{t})^{\lambda-1} \mathbf{u} \mathbf{t} \bar{\mathbf{p}}_{\delta_t} \\ &= \mathbf{p}_{\delta_t} \mathbf{t} (\mathbf{u} \mathbf{t})^{\lambda-1} \mathbf{u} \bar{\mathbf{p}}_{\delta_t} \\ &= \mathbf{s} \mathbf{p}_{\delta_t} (\mathbf{u} \mathbf{t})^{\lambda-1} \mathbf{u} \bar{\mathbf{p}}_{\delta_t} \\ &= \mathbf{s} \mathbf{r}_{\delta_t, u}, \end{aligned}$$

and hence $\mathbf{r}_{\delta_t, u}$ commutes with \mathbf{s} . By Theorem 4.1, the group $C_W(s)$ is generated by $\{s, p_\gamma, r_{\delta_t, u}\}$. Since each generator has a pre-image under ν that lies in $C_A(\mathbf{s})$, the assertion follows. □

We now prove the main result of this section.

THEOREM 4.3. *The involutory quandle of an Artin quandle is the corresponding Coxeter quandle, that is, $(\mathcal{D}(\mathbf{S}^A))_2 \cong \mathcal{D}(S^W)$.*

PROOF. The surjection $\nu : \mathbf{A} \rightarrow W$ induces a surjective quandle homomorphism $\tilde{\nu} : \mathcal{D}(\mathbf{S}^A) \rightarrow \mathcal{D}(S^W)$. Since $(\mathcal{D}(\mathbf{S}^A))_2$ is involutory, $\tilde{\nu}$ descends to a surjective quandle homomorphism

$$\bar{\nu} : (\mathcal{D}(\mathbf{S}^A))_2 \rightarrow \mathcal{D}(S^W).$$

If $\pi : \mathcal{D}(\mathbf{S}^A) \rightarrow (\mathcal{D}(\mathbf{S}^A))_2$ is the natural surjection, then $\bar{\nu} \pi = \tilde{\nu}$. We claim that $\bar{\nu}$ is an isomorphism. Suppose that $\tilde{\nu}(\mathbf{x}) = \tilde{\nu}(\mathbf{y})$ for some $\mathbf{x}, \mathbf{y} \in \mathcal{D}(\mathbf{S}^A)$. Since $\tilde{\nu}$ preserves orbits (under action of the inner automorphism group), it follows that \mathbf{x} and \mathbf{y} must be in the same orbit. Thus, we can write $\mathbf{x} = \mathbf{s}_{i_k}^{\epsilon_k} \cdots \mathbf{s}_{i_1}^{\epsilon_1} \mathbf{s} \mathbf{s}_{i_1}^{-\epsilon_1} \cdots \mathbf{s}_{i_k}^{-\epsilon_k}$ and $\mathbf{y} = \mathbf{s}_{j_r}^{\delta_r} \cdots \mathbf{s}_{j_1}^{\delta_1} \mathbf{s} \mathbf{s}_{j_1}^{-\delta_1} \cdots \mathbf{s}_{j_r}^{-\delta_r}$ for some $\mathbf{s}, \mathbf{s}_{i_t}, \mathbf{s}_{j_t} \in \mathbf{S}$ and $\epsilon_t, \delta_t \in \{1, -1\}$. Now $\tilde{\nu}(\mathbf{x}) = \tilde{\nu}(\mathbf{y})$ implies that $\nu(\mathbf{s}_{j_1}^{-\delta_1} \cdots \mathbf{s}_{j_r}^{-\delta_r} \mathbf{s}_{i_k}^{\epsilon_k} \cdots \mathbf{s}_{i_1}^{\epsilon_1})$ commutes with $\nu(\mathbf{s}) = s$. In view of Lemma 4.2, the short exact sequence $1 \rightarrow \ker(\nu) \rightarrow \mathbf{A} \xrightarrow{\nu} W \rightarrow 1$ induces a short exact sequence

$$1 \rightarrow \ker(\nu) \cap C_A(\mathbf{s}) \rightarrow C_A(\mathbf{s}) \rightarrow C_W(s) \rightarrow 1.$$




FIGURE 1. Arcs $c_{i,j}$ on the disc for $n = 6$.

Thus, we can write $\mathbf{s}_{j_1}^{-\delta_1} \cdots \mathbf{s}_{j_r}^{-\delta_r} \mathbf{s}_{i_k}^{\epsilon_k} \cdots \mathbf{s}_{i_1}^{\epsilon_1} = \mathbf{u}\mathbf{v}$ for some $\mathbf{u} \in \ker(\nu) = \langle\langle \mathbf{s}^2 \mid \mathbf{s} \in \mathbf{S} \rangle\rangle$ and some $\mathbf{v} \in C_A(\mathbf{s})$. Thus,

$$\begin{aligned} & \pi(\mathbf{s}_{j_1}^{-\delta_1} \cdots \mathbf{s}_{j_r}^{-\delta_r} \mathbf{s}_{i_k}^{\epsilon_k} \cdots \mathbf{s}_{i_1}^{\epsilon_1} \mathbf{s} \mathbf{s}_1^{-\epsilon_1} \cdots \mathbf{s}_{i_k}^{-\epsilon_k} \mathbf{s}_{j_r}^{\delta_r} \cdots \mathbf{s}_{j_1}^{\delta_1}) \\ &= \pi(\mathbf{u}\mathbf{v}\mathbf{s}\mathbf{v}^{-1}\mathbf{u}^{-1}) \\ &= \pi(\mathbf{u}\mathbf{s}\mathbf{u}^{-1}), \quad \text{since } \mathbf{v} \in C_A(\mathbf{s}) \\ &= \pi(\mathbf{s}), \quad \text{since } \mathbf{u} \in \ker(\nu), \end{aligned}$$

which shows that $\pi(\mathbf{x}) = \pi(\mathbf{y})$ and hence $\bar{\nu}$ is an isomorphism. □

As a consequence of Propositions 3.1 and 4.3, we obtain a presentation of the enveloping group of the Coxeter quandle, which extends [2, Proposition 3.3].

COROLLARY 4.4. *Let (W, S) be a Coxeter system. Then $\text{Env}(\mathcal{D}(S^W)) \cong \mathbf{A}/\mathbf{N}$, where \mathbf{A} is the corresponding Artin group and $\mathbf{N} = \langle\langle \mathbf{s}_i^2 \mathbf{s}_j \mathbf{s}_i^{-2} \mathbf{s}_j^{-1} \mid \mathbf{s}_i, \mathbf{s}_j \in \mathbf{S} \rangle\rangle$.*

Our next result gives the size of the smallest nontrivial quandle quotient of the Dehn quandle of the braid group with respect to its standard generating set. The proof is essentially rephrasing [20, Lemma 8] in the language of quandles.

PROPOSITION 4.5. *Let $n \geq 5$ and $f : \mathcal{D}(S^{B_n}) \rightarrow Q$ be a surjective quandle homomorphism onto a quandle with at least two elements. Then, $|Q| \geq n(n - 1)/2$ and the bound is sharp.*

PROOF. We view the braid group B_n as the mapping class group of the disc \mathbb{D}^2 with n marked points. Consider the set $\{c_{i,j} \mid 1 \leq i < j \leq n\}$ of $n(n - 1)/2$ arcs on \mathbb{D}^2 as shown in Figure 1 for $n = 6$. Let $\sigma_{i,j}$ denote the isotopy class of the (anti-clockwise) half Dehn twist along the arc $c_{i,j}$ and let $X = \{\sigma_{i,j} \mid 1 \leq i < j \leq n\}$. Then B_n is generated by the subset $S = \{\sigma_{1,2}, \sigma_{2,3}, \dots, \sigma_{n-1,n}\}$ of X . Note that all the elements of X are conjugate to each

other in B_n and hence lie in $\mathcal{D}(S^{B_n})$. Let $f : \mathcal{D}(S^{B_n}) \rightarrow Q$ be a surjective quandle homomorphism onto a quandle with at least two elements. Then we have the following cases:

Case (1): Suppose that f is not injective on S , that is, $f(\sigma_{i,i+1}) = f(\sigma_{j,j+1})$ for some $i \neq j$. Since $n \geq 5$, by construction of X , we can choose an arc $c_{k,k+1}$ such that it has exactly one end point in common with precisely one of $c_{i,i+1}$ or $c_{j,j+1}$ and no end point in common with the other. Without loss of generality, we can assume that $c_{k,k+1}$ and $c_{j,j+1}$ have one end point in common, while $c_{k,k+1}$ and $c_{i,i+1}$ are disjoint. Then, the half Dehn twists satisfy the relations

$$\sigma_{k,k+1}\sigma_{j,j+1}\sigma_{k,k+1} = \sigma_{j,j+1}\sigma_{k,k+1}\sigma_{j,j+1} \quad \text{and} \quad \sigma_{k,k+1}\sigma_{i,i+1} = \sigma_{i,i+1}\sigma_{k,k+1}.$$

Applying f to the preceding relations and using the equality $f(\sigma_{i,i+1}) = f(\sigma_{j,j+1})$, we obtain $f(\sigma_{k,k+1}) = f(\sigma_{i,i+1})$. Proceeding in this manner, we show that f is constant on S . Since S also generates $\mathcal{D}(S^{B_n})$ as a quandle [9, Proposition 3.2], it follows that f is constant on $\mathcal{D}(S^{B_n})$, which is a contradiction. Hence, f is injective on S .

Case (2): Suppose that f is not injective on X . There are two possibilities here.

Subcase (2a): Suppose that $f(\sigma_{i,i+1}) = f(\sigma_{k,l})$ where $\{i, i + 1\} \neq \{k, l\}$ as sets. Since $n \geq 5$, we can choose an arc $c_{r,r+1}$ such that it has exactly one end point in common with precisely one of $c_{i,i+1}$ or $c_{k,l}$, and no end point in common with the other. If $c_{r,r+1}$ and $c_{i,i+1}$ have one end point in common, then as in Case (1), we obtain $f(\sigma_{i,i+1}) = f(\sigma_{r,r+1})$. Similarly, if $c_{r,r+1}$ and $c_{k,l}$ have one end point in common, then as in Case (1), we get $f(\sigma_{i,i+1}) = f(\sigma_{r,r+1})$. Again, proceeding as in Case (1) leads to a contradiction.

Subcase (2b): Suppose that $f(\sigma_{i,j}) = f(\sigma_{k,l})$ where $\{i, j\} \neq \{k, l\}$. Since $n \geq 5$, we can choose an arc $c_{r,r+1}$ such that it has exactly one end point in common with precisely one of $c_{i,j}$ or $c_{k,l}$, and no end point in common with the other. Proceeding as before, we deduce that $f(\sigma_{i,j}) = f(\sigma_{k,l}) = f(\sigma_{r,r+1})$, which is Subcase (2a).

Hence, it follows that the map f must be injective on the set X , and therefore $|Q| \geq n(n - 1)/2$. For sharpness of the bound, consider the surjective group homomorphism $B_n \rightarrow \Sigma_n$, where Σ_n is the symmetric group on n symbols. The group homomorphism induces a surjective quandle homomorphism $\mathcal{D}(S^{B_n}) \rightarrow \mathcal{D}(T^{\Sigma_n})$, where $T = \{(i, i + 1) \mid 1 \leq i \leq n - 1\}$ is the image of S . Since $|\mathcal{D}(T^{\Sigma_n})| = n(n - 1)/2$, the assertion follows. □

REMARK 4.6. It follows from Theorem 4.3 that $\mathcal{D}(T^{\Sigma_n})$ is precisely the 2-quandle of $\mathcal{D}(S^{B_n})$. It is intriguing to know the smallest quandle quotients of general Artin quandles.

5. Finiteness of n -quandles of Dehn quandles of surfaces

In this main section, we consider finiteness of n -quandles of Dehn quandles of closed orientable surfaces. For each genus, with some exceptions, we determine all values of n for which the n -quandle of the Dehn quandle of the surface S_g is finite. We also determine the explicit structures of their 2-quandles.




FIGURE 2. Surface of genus g with curves a_i, b_i and c_i .

5.1. 2- and 3-quandles of Dehn quandles of surfaces. Let S_g be a closed oriented surface of genus $g \geq 1$ and

$$\Psi : \mathcal{M}_g \rightarrow \text{Sp}(2g, \mathbb{Z})$$

the symplectic representation of the mapping class group \mathcal{M}_g of S_g . For an integer $n \geq 2$, let $\psi_n : \text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}_n)$ be the mod n reduction homomorphism and

$$\Psi_n := \psi_n \Psi : \mathcal{M}_g \rightarrow \text{Sp}(2g, \mathbb{Z}_n)$$

the mod n reduction of the symplectic representation. Given a simple closed curve a in S_g , let τ_a denote the (right-hand) Dehn twists about a . Consider the surface S_g with curves a_i, b_i, c_i as in Figure 2.

LEMMA 5.1. *Let p be a prime and $g \geq 1$. Then, an element $A \in \text{Sp}(2g, \mathbb{Z}_p)$ lies in the centraliser of $\Psi_p(\tau_{a_1})$ if and only if A has first column $(a, 0, 0, \dots, 0)$ and second row $(0, a, 0, 0, \dots, 0)$, where $a \equiv \pm 1 \pmod{p}$.*

PROOF. Let $E_{i,j}$ denote the elementary matrix with (i, j) -entry as 1 and each other entry as 0. Observe that $\Psi_p(\tau_{a_1}) = I_{2g} - E_{1,2}$. If $A \in \text{Sp}(2g, \mathbb{Z}_p)$ commutes with $\Psi_p(\tau_{a_1})$, then it follows from elementary linear algebra that A has first column $(a, 0, 0, \dots, 0)$ and second row $(0, a, 0, 0, \dots, 0)$ for some $a \in \mathbb{Z}_p^\times$. Since $A^t J A = J$, where J represents the symplectic form, we obtain $a^2 \equiv 1 \pmod{p}$. Since $a^2 \equiv 1 \pmod{p}$ if and only if $a \equiv \pm 1 \pmod{p}$, the assertion follows. Conversely, any symplectic matrix of this form commutes with $\Psi_p(\tau_{a_1})$. \square

LEMMA 5.2. *Let p be a prime and $g \geq 1$. Then the centraliser of $\Psi_p(\tau_{a_1})$ in $\text{Sp}(2g, \mathbb{Z}_p)$ is generated by $\Psi_p(S) \cup \{-I_{2g}\}$, where*

$$S = \{\tau_{a_1}, \tau_{a_2}, \dots, \tau_{a_g}, \tau_{b_2}, \dots, \tau_{b_g}, \tau_{c_1}, \tau_{c_2}, \dots, \tau_{c_{g-1}}\}.$$

PROOF. For $g = 1$, a direct computation shows that the centraliser of $\Psi_p(\tau_{a_1})$ in $\text{Sp}(2, \mathbb{Z}_p)$ is generated by $\Psi_p(\tau_{a_1}) \cup \{-I_{2g}\}$. We assume that $g \geq 2$. Clearly, $\Psi_p(S) \cup \{-I_{2g}\}$ is a subset of the centraliser of $\Psi_p(\tau_{a_1})$ in $\text{Sp}(2g, \mathbb{Z}_p)$. If $A = (A_{i,j})$ lies in the centraliser of $\Psi_p(\tau_{a_1})$ in $\text{Sp}(2g, \mathbb{Z}_p)$, then Lemma 5.1 implies that $A_{1,1} = \pm 1$. Multiplying A by $-I_{2g}$, we can assume that $A_{1,1} = 1$. It remains to prove such an A can be written as a product of elements from $\Psi_p(S)$. Consider the curves d_1, d_2, \dots, d_{g-1} as




FIGURE 3. Surface of genus g with curves d_i .

shown in Figure 3. For each $1 \leq i \leq g - 1$, let $M_i := \Psi_p(\tau_{d_i} \tau_{a_1}^{-1} \tau_{a_{i+1}}^{-1})$. Then, we can see that

$$M_i = I_{2g} + E_{2i+1,2} + E_{1,2i+2}.$$

Let $N_i := \Psi_p(\tau_{a_{i+1}} \tau_{b_{i+1}} \tau_{a_{i+1}})^{-1} M_i \Psi_p(\tau_{a_{i+1}} \tau_{b_{i+1}} \tau_{a_{i+1}})$. Then, a direct computation shows that

$$N_i = I_{2g} - E_{2i+2,2} + E_{1,2i+1}.$$

Now, consider the matrix

$$B := \left(\prod_{i=1}^{g-1} N_{g-i}^{A_{2(g-i)+2,2}} M_{g-i}^{-A_{2(g-i)+1,2}} \right) A,$$

where a matrix raised to the power by the entry $A_{i,j}$ represents the same matrix raised to the power by a representative of $A_{i,j}$ in the subset $\{0, 1, 2, \dots, p - 1\}$ of integers. By the choice of matrices M_i and N_i , it follows that the second column of B is $(B_{1,2}, 1, 0, 0, \dots, 0)$. Next, take

$$C := \Psi_p(\tau_{a_1})^{B_{1,2}} B.$$

It is easy to see that the first two columns of C are $(1, 0, 0, \dots, 0)$ and $(0, 1, 0, \dots, 0)$, respectively. Since C is symplectic, it has the form

$$C = \begin{bmatrix} I_2 & O_{2,2g-2} \\ O_{2g-2,2} & D \end{bmatrix},$$

where $O_{m,n}$ is the $m \times n$ null matrix and $D \in \text{Sp}(2g - 2, \mathbb{Z}_p)$. Note that $\text{Sp}(2g - 2, \mathbb{Z}_p)$ is generated by $\Psi_2(\{\tau_{a_2}, \dots, \tau_{a_g}, \tau_{b_2}, \dots, \tau_{b_g}, \tau_{c_2}, \dots, \tau_{c_{g-1}}\})$. We claim that each τ_{d_i} can be written as a product of elements from S , which would complete the proof. Since $d_1 = c_1$, the claim follows when $g = 2$.

Let $\mathcal{M}_g(\vec{\alpha})$ be the subgroup of \mathcal{M}_g consisting of all elements that preserve a nonseparating simple closed curve α with orientation. Let $\mathcal{M}_{g,\alpha}$ be the mapping class group of the surface $S_{g,\alpha}$ (which is homeomorphic to a surface of genus $g - 1$ and two boundary components) obtained by cutting the surface S_g along the curve α and then taking its closure. Let $\{\delta_1, \delta_2\}$ be the two boundary components of $S_{g,\alpha}$. Then, by [25, Lemma 1.20], there is a short exact sequence of groups

$$1 \rightarrow \langle \tau_{\delta_1} \tau_{\delta_2}^{-1} \rangle \rightarrow \mathcal{M}_{g,\alpha} \rightarrow \mathcal{M}_g(\vec{\alpha}) \rightarrow 1,$$

where $\langle \tau_{\delta_1}, \tau_{\delta_2}^{-1} \rangle \cong \mathbb{Z}$. Applying this result to the oriented curve a_1 and using the generating set for \mathcal{M}_{g,a_1} (see [12, Corollary 4.16 and Figure 4.10]),

$$\mathcal{M}_g(\vec{a_1}) = \langle \tau_{a_1}, \tau_{a_2}, \dots, \tau_{a_g}, \tau_{b_2}, \dots, \tau_{b_g}, \tau_{c_1}, \tau_{c_2}, \dots, \tau_{c_{g-1}} \rangle.$$

Since each τ_{d_i} preserves the curve a_1 with orientation, the claim follows. □

Using Lemma 5.2 and choosing the hyperelliptic involution as a pre-image of $-I_{2g}$, the short exact sequence

$$1 \rightarrow \ker(\Psi_p) \rightarrow \mathcal{M}_g \xrightarrow{\Psi_p} \mathrm{Sp}(2g, \mathbb{Z}_p) \rightarrow 1$$

induces the following short exact sequence at the level of centralisers.

COROLLARY 5.3. *For each prime p and each $g \geq 1$, there is a short exact sequence of groups*

$$1 \rightarrow \ker(\Psi_p) \cap C_{\mathcal{M}_g}(\tau_{a_1}) \rightarrow C_{\mathcal{M}_g}(\tau_{a_1}) \rightarrow C_{\mathrm{Sp}(2g, \mathbb{Z}_p)}(\Psi_p(\tau_{a_1})) \rightarrow 1. \tag{5-1}$$

We briefly recall from [9, Section 7] the construction of the projective primitive homological quandle $\mathcal{P}_{g,n}$ of the surface S_g , where $g \geq 1$ and $n \geq 2$. Let \mathcal{P}'_g denote the set of all primitive elements in $H_1(S_g, \mathbb{Z})$ and $\mathcal{P}'_{g,n}$ denote the set of all primitive elements in $H_1(S_g, \mathbb{Z}_n)$. The algebraic intersection number $\hat{i}(-, -)$ gives a skew-symmetric (in fact, symplectic) bilinear form on the \mathbb{Z} -module $H_1(S_g, \mathbb{Z})$. For $x, y \in \mathcal{P}'_g$, the binary operation

$$x * y := x + \hat{i}(x, y)y$$

gives a quandle structure on \mathcal{P}'_g . Similarly, reduction modulo n defines a quandle structure on $\mathcal{P}'_{g,n}$.

Let $\mathcal{P}_g := \mathcal{P}'_g / \mathbb{Z}_2$ and $\mathcal{P}_{g,n} := \mathcal{P}'_{g,n} / \mathbb{Z}_2$ be quotients under the natural involutory action of $\mathbb{Z}_2 = \{1, -1\}$. It is clear that quandle structures descend to quandle structures on \mathcal{P}_g and $\mathcal{P}_{g,n}$. Further, reduction modulo n gives a surjective quandle homomorphism $\mathcal{P}'_g \rightarrow \mathcal{P}'_{g,n}$, which further induces a surjective quandle homomorphism $\mathcal{P}_g \rightarrow \mathcal{P}_{g,n}$. The quandles \mathcal{P}_g and $\mathcal{P}_{g,n}$ are called *projective primitive homological quandles* of S_g . Note that $\mathcal{P}_{g,2}$ is an involutory quandle of order $2^{2g} - 1$.

Recall from [12, Proposition 6.2] that a nonzero element of $H_1(S_g, \mathbb{Z})$ is primitive if and only if it is represented by an oriented simple closed curve. For the isotopy class a of an oriented simple closed curve in S_g , we denote by $[a] \in H_1(S_g, \mathbb{Z})$ its homology class. Since \mathcal{D}_g^{ns} consists of isotopy classes of unoriented simple closed curves, there are two choices for the homology class $[a] \in \mathcal{P}'_g$ for each $a \in \mathcal{D}_g^{ns}$. We choose $[a]$ such that the entry in its first nonzero coordinate (from left) is positive. This gives a surjection $\mathcal{D}_g^{ns} \rightarrow \mathcal{P}'_g$ given by $a \mapsto [a]$. Composing the surjections $\mathcal{D}_g^{ns} \rightarrow \mathcal{P}'_g$, $\mathcal{P}'_g \rightarrow \mathcal{P}_g$ and $\mathcal{P}_g \rightarrow \mathcal{P}_{g,n}$ gives a surjection

$$\phi : \mathcal{D}_g^{ns} \rightarrow \mathcal{P}_{g,n},$$

which is a quandle homomorphism due to [9, Theorem 7.1]. Let $(\mathcal{D}_g^{ns})_n$ denote the n -quandle of the Dehn quandle \mathcal{D}_g^{ns} and

$$\pi : \mathcal{D}_g^{ns} \rightarrow (\mathcal{D}_g^{ns})_n$$

the natural quotient map.

THEOREM 5.4. *Let \mathcal{D}_g^{ns} be the Dehn quandle of the closed orientable surface S_g of genus $g \geq 1$. Then the following assertions hold:*

- (1) $(\mathcal{D}_g^{ns})_2 \cong \mathcal{P}_{g,2}$ for each $g \geq 1$;
- (2) $(\mathcal{D}_2^{ns})_3 \cong \mathcal{P}_{2,3}$.

PROOF. Let $\phi : \mathcal{D}_g^{ns} \rightarrow \mathcal{P}_{g,2}$ be the surjective quandle homomorphism as discussed above. Since $(\mathcal{D}_g^{ns})_2$ is involutory, ϕ induces a surjective quandle homomorphism

$$\bar{\phi} : (\mathcal{D}_g^{ns})_2 \rightarrow \mathcal{P}_{g,2}$$

such that $\bar{\phi} \pi = \phi$. We claim that $\bar{\phi}$ is an isomorphism. Suppose that $\phi(x) = \phi(y)$ for $x, y \in \mathcal{D}_g^{ns}$. It suffices to prove that $\pi(x) = \pi(y)$. We identify elements of \mathcal{D}_g^{ns} with corresponding Dehn twists in \mathcal{M}_g with the quandle operation as conjugation. Since \mathcal{D}_g^{ns} is a connected quandle, we have $x = \tau_{a_1} *^{\epsilon_1} \tau_{z_1} *^{\epsilon_2} \tau_{z_2} * \dots *^{\epsilon_k} \tau_{z_k}$ and $y = \tau_{a_1} *^{\delta_1} \tau_{w_1} *^{\delta_2} \tau_{w_2} * \dots *^{\delta_r} \tau_{w_r}$ for some $z_i, w_j \in \mathcal{D}_g^{ns}$ and $\epsilon_i, \delta_j \in \{1, -1\}$, where a_1 denotes the usual curve (see Figure 2). Since $\phi(x) = \phi(y)$, we have

$$\phi(\tau_{z_k}^{\epsilon_k} \dots \tau_{z_2}^{\epsilon_2} \tau_{z_1}^{\epsilon_1} \tau_{a_1}^{-\epsilon_1} \tau_{z_2}^{-\epsilon_2} \dots \tau_{z_k}^{-\epsilon_k}) = \phi(\tau_{w_r}^{\delta_r} \dots \tau_{w_2}^{\delta_2} \tau_{w_1}^{\delta_1} \tau_{a_1}^{-\delta_1} \tau_{w_2}^{-\delta_2} \dots \tau_{w_r}^{-\delta_r}).$$

It follows from [9, Proposition 7.6(ii)] that ϕ is simply the restriction of $\Psi_2 : \mathcal{M}_g \rightarrow \text{Sp}(2g, \mathbb{Z}_2)$. Thus, the preceding equation can be written as

$$\Psi_2(\tau_{z_k}^{\epsilon_k} \dots \tau_{z_2}^{\epsilon_2} \tau_{z_1}^{\epsilon_1} \tau_{a_1}^{-\epsilon_1} \tau_{z_2}^{-\epsilon_2} \dots \tau_{z_k}^{-\epsilon_k}) = \Psi_2(\tau_{w_r}^{\delta_r} \dots \tau_{w_2}^{\delta_2} \tau_{w_1}^{\delta_1} \tau_{a_1}^{-\delta_1} \tau_{w_2}^{-\delta_2} \dots \tau_{w_r}^{-\delta_r}),$$

which implies that $\Psi_2(\tau_{w_1}^{-\delta_1} \tau_{w_2}^{-\delta_2} \dots \tau_{w_r}^{-\delta_r} \tau_{z_k}^{\epsilon_k} \dots \tau_{z_2}^{\epsilon_2} \tau_{z_1}^{\epsilon_1})$ commutes with $\Psi_2(\tau_{a_1})$. Using Corollary 5.3, we can write $\tau_{w_1}^{-\delta_1} \tau_{w_2}^{-\delta_2} \dots \tau_{w_r}^{-\delta_r} \tau_{z_k}^{\epsilon_k} \dots \tau_{z_2}^{\epsilon_2} \tau_{z_1}^{\epsilon_1} = uv$ such that $\Psi_2(u) = I_{2g}$ and v commutes with τ_{a_1} (by choosing an appropriate section to (5-1)). By [16, Proposition 2.1], we know that $\ker(\Psi_2)$ is generated by squares of Dehn twists along nonseparating curves. Thus,

$$\begin{aligned} &\pi(\tau_{a_1} *^{\epsilon_1} \tau_{z_1} *^{\epsilon_2} \tau_{z_2} * \dots *^{\epsilon_k} \tau_{z_k} *^{-1} *^{-\delta_r} \tau_{w_r} * \dots *^{-\delta_2} \tau_{w_2} *^{-\delta_1} \tau_{w_1}) \\ &= \pi(\tau_{a_1} * v * u) \\ &= \pi(\tau_{a_1} * u), \quad \text{since } v \text{ commutes with } \tau_{a_1} \\ &= \pi(\tau_{a_1}) * \pi(u) \\ &= \pi(\tau_{a_1}), \end{aligned}$$

which implies that $\pi(x) = \pi(y)$. This completes the proof of assertion (1).

For assertion (2), we claim that a kernel of $\Psi_3 : \mathcal{M}_2 \rightarrow \text{Sp}(4, \mathbb{Z}_3)$ is generated by cubes of Dehn twists along nonseparating curves. Let $N = \langle\langle \tau_{a_1}^3 \rangle\rangle$ be the normal closure

in \mathcal{M}_2 of the set of cubes of Dehn twists along all nonseparating simple closed curves on S_2 . By [16, Proposition 2.1], $\ker(\Psi) \leq N$. Let $M = \langle\langle I_4 + 3E_{1,2} \rangle\rangle$ be the normal closure of $I_4 + 3E_{1,2}$ in $\text{Sp}(4, \mathbb{Z})$ and $\psi_3 : \text{Sp}(4, \mathbb{Z}) \rightarrow \text{Sp}(4, \mathbb{Z}_3)$ be the mod 3 reduction homomorphism. By [22, Hilfssatz 9.2], $\ker(\psi_3) = M$. Hence, it follows that

$$N/\ker(\Psi) \cong \Psi(N) = M = \ker(\psi_3).$$

This gives

$$\mathcal{M}_2/N \cong (\mathcal{M}_2/\ker(\Psi))/(N/\ker(\Psi)) \cong \text{Sp}(4, \mathbb{Z})/\ker(\psi_3) \cong \text{Sp}(4, \mathbb{Z}_3).$$

Since $\Psi_3 = \psi_3\Psi$, the preceding isomorphism is, in fact, induced by Ψ_3 . Hence, it follows that $\ker(\Psi_3) = N = \langle\langle \tau_{a_1}^3 \rangle\rangle$. The proof of assertion (2) now follows analogously to that of assertion (1). □

REMARK 5.5. Lemma 5.2 is inspired by [1, 8], wherein similar arguments have been used to understand the stabiliser of a homological vector in the mapping class group \mathcal{M}_g , which also preserves the fibre over that homological vector under the homomorphism $\phi : \mathcal{D}_g^{ns} \rightarrow \mathcal{P}_{g,n}$.

5.2. n -quandles of Dehn quandles of surfaces. We now present our main result, which can be thought of as the Dehn quandle analogue of a similar result of Hoste and Shanahan [14] for link quandles.

THEOREM 5.6. *Let \mathcal{D}_g^{ns} be the Dehn quandle of the closed orientable surface of genus $g \geq 1$. Then the following assertions hold:*

- (1) $(\mathcal{D}_1^{ns})_n$ is finite if and only if $n = 2, 3, 4, 5$;
- (2) $(\mathcal{D}_2^{ns})_n$ is finite if and only if $n = 2, 3$;
- (3) if $g \geq 3$, then $(\mathcal{D}_g^{ns})_n$ is finite for $n = 2$ and infinite for $n \neq 2, 3, 4, 6, 8, 12$.

PROOF. It follows from [23] that $\mathcal{D}_1^{ns} \cong Q(3_1)$, the knot quandle of the trefoil knot 3_1 . Further, it follows from [14, Section 5] that $(Q(3_1))_n$ is finite if and only if $n = 2, 3, 4, 5$, which proves assertion (1).

By Theorem 5.4, we obtain that $(\mathcal{D}_2^{ns})_n$ is finite for $n = 2, 3$. Let \mathcal{M}_g^b be the mapping class group of the surface S_g^b of genus g with $b \geq 0$ boundary components and let $(\mathcal{M}_g^b)_n$ be the quotient of \mathcal{M}_g^b by the normal closure of n th powers of Dehn twists along nonseparating simple closed curves. We avoid writing b in the notation whenever $b = 0$. It is proved in [16, Theorem 4] that $(\mathcal{M}_2^b)_n$ is infinite for $b \geq 0$ and $n > 3$. Taking $b = 0$, it follows from Theorem 3.4 that $(\mathcal{D}_2^{ns})_n$ is infinite for $n > 3$, which proves assertion (2).

Assume that $g \geq 3$. By Theorem 5.4, $(\mathcal{D}_g^{ns})_2$ is finite. However, [13, Corollary 1.2] gives that $(\mathcal{M}_g)_n$ is infinite for $n \neq 2, 3, 4, 6, 8, 12$. Thus, Theorem 3.4 implies that $(\mathcal{D}_g^{ns})_n$ is infinite for $n \neq 2, 3, 4, 6, 8, 12$. □

REMARK 5.7. We do not know whether $(\mathcal{D}_g^{ns})_n$ is finite for $g \geq 3$ and $n = 3, 4, 6, 8, 12$.

We conclude with a bound on the size of the smallest nontrivial quandle quotient of the Dehn quandle of a surface. The proof is based on [20, Lemma 11].

PROPOSITION 5.8. *Let $g \geq 1$ and $f : \mathcal{D}_g^{ns} \rightarrow Q$ be a surjective quandle homomorphism onto a quandle with at least two elements. Then, $|Q| \geq 2^{2g} - 1$ and the bound is sharp.*

PROOF. By [9, Theorem 7.1], there is a surjective quandle homomorphism from \mathcal{D}_g^{ns} onto the projective primitive homological quandle $\mathcal{P}_{g,2}$ of order $2^{2g} - 1$. It follows from [12, Proposition 6.2] that a nonzero element of $H_1(S_g, \mathbb{Z}_2)$ is primitive if and only if it is represented by an oriented simple closed curve on S_g . Following [20, Section 5], we suitably choose simple closed curves on S_g such that they represent all primitive vectors in $H_1(S_g, \mathbb{Z}_2)$. Let X be the set of isotopy classes of (right) Dehn twists about these suitably chosen curves. Viewing \mathcal{D}_g^{ns} as a subquandle of the conjugation quandle of \mathcal{M}_g , let $f : \mathcal{D}_g^{ns} \rightarrow Q$ be a surjective quandle homomorphism. Suppose that $f(\tau_\alpha) = f(\tau_\beta)$ for some $\tau_\alpha, \tau_\beta \in X$ with $\tau_\alpha \neq \tau_\beta$. Then, by the construction of X , one can find a simple closed curve γ such that τ_γ commutes with precisely one of τ_α or τ_β and satisfies the braid relation with the other. Without loss of generality, suppose that

$$\tau_\gamma \tau_\alpha \tau_\gamma = \tau_\alpha \tau_\gamma \tau_\alpha \quad \text{and} \quad \tau_\gamma \tau_\beta = \tau_\beta \tau_\gamma.$$

Applying f to the preceding relations and using the equality $f(\tau_\alpha) = f(\tau_\beta)$, we obtain $f(\tau_\alpha) = f(\tau_\gamma)$. Proceeding in this manner, we show that f is constant on X . Since X also generates \mathcal{D}_g^{ns} as a quandle [9, Proposition 3.2], it follows that f is constant on \mathcal{D}_g^{ns} , which is a contradiction. Hence, f must be injective on X , and therefore $|Q| \geq 2^{2g} - 1$. \square

Acknowledgements

The authors thank Prof. Louis Funar for pointing out an oversight in our earlier proof of assertion (3) of Theorem 5.6. Some of the ideas in this paper were developed by the authors during the preparation of [9, 10]. They thank Dr. Hitesh Raundal for his interest in the ideas.

References

- [1] N. Agarwal, S. Dey, N. K. Dhanwani and K. Rajeevsarathy, ‘Liftable mapping class groups of regular cyclic covers’, *Houston J. Math.* **47**(1) (2021), 223–243.
- [2] T. Akita, ‘The adjoint group of a Coxeter quandle’, *Kyoto J. Math.* **60** (2020), 1245–1260.
- [3] T. Akita, A. Hasegawa and M. Tanno, ‘Structure of the associated groups of quandles’, *Kodai Math. J.* **45** (2022), 270–281.
- [4] D. Allcock, ‘Reflection centralizers in Coxeter groups’, *Transform. Groups* **18**(3) (2013), 599–613.
- [5] J. S. Birman, *Braids, Links, and Mapping Class Groups*, Annals of Mathematics Studies, 82 (Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1974).

- [6] B. Brink, 'On centralizers of reflections in Coxeter groups', *Bull. Lond. Math. Soc.* **28**(5) (1996), 465–470.
- [7] B. Brink and R. B. Howlett, 'Normalizers of parabolic subgroups in Coxeter groups', *Invent. Math.* **136**(2) (1999), 323–351.
- [8] S. Dey, N. K. Dhanwani, H. Patil and K. Rajeevsarathy, 'Generating the liftable mapping class groups of regular cyclic covers', Preprint, 2021, [arXiv:2111.01626v1](https://arxiv.org/abs/2111.01626v1).
- [9] N. K. Dhanwani, H. Raundal and M. Singh, 'Dehn quandles of groups and orientable surfaces', *Fund. Math.* **263** (2023), 167–201.
- [10] N. K. Dhanwani, H. Raundal and M. Singh, 'Presentations of Dehn quandles', *J. Algebra* **636** (2023), 207–247.
- [11] W. D. Dunbar, 'Geometric orbifolds', *Rev. Mat. Univ. Complut. Madrid* **1**(1–3) (1988), 67–99.
- [12] B. Farb and D. Margalit, *A Primer on Mapping Class Groups*, Princeton Mathematical Series, 49 (Princeton University Press, Princeton, NJ, 2012).
- [13] L. Funar, 'On the TQFT representations of the mapping class groups', *Pacific J. Math.* **188**(2) (1999), 251–274.
- [14] J. Hoste and P. D. Shanahan, 'Links with finite n -quandles', *Algebr. Geom. Topol.* **17**(5) (2017), 2807–2823.
- [15] R. B. Howlett, 'Normalizers of parabolic subgroups of reflection groups', *J. Lond. Math. Soc.* (2) **21**(1) (1980), 62–80.
- [16] S. Humphries, 'Normal closures of powers of Dehn twists in mapping class groups', *Glasg. Math. J.* **34**(3) (1992), 314–317.
- [17] D. Joyce, 'An algebraic approach to symmetry with applications to knot theory', PhD Thesis, University of Pennsylvania, 1979.
- [18] D. Joyce, 'A classifying invariant of knots, the knot quandle', *J. Pure Appl. Algebra* **23** (1982), 37–65.
- [19] S. Kamada and Y. Matsumoto, 'Certain racks associated with the braid groups', in: *Knots in Hellas '98 (Delphi)*, Series on Knots and Everything, 24 (eds C. McA. Gordon, V. F. R. Jones, L. Kauffman, S. Lambropoulou and J. H. Przytycki) (World Scientific Publishing, River Edge, NJ, 2000), 118–130.
- [20] S. Kolay, 'Smallest non-cyclic quotients of braid and mapping class groups', *Geom. Topol.* **27** (2023), 2479–2496.
- [21] S. V. Matveev, 'Distributive groupoids in knot theory', *Russian Mat. Sb. (N.S.)* **119**(1) (1982), 78–88; translated in *Math. USSR Sb.* **47**(1) (1984), 73–83.
- [22] J. Mennicke, 'Zur theorie der Siegelschen Modulgruppe', *Math. Ann.* **159** (1965), 115–129.
- [23] M. Niebrzydowski and J. H. Przytycki, 'The quandle of the trefoil knot as the Dehn quandle of the torus', *Osaka J. Math.* **46** (2009), 645–659.
- [24] T. Nosaka, 'Central extensions of groups and adjoint groups of quandles', in: *Geometry and Analysis of Discrete Groups and Hyperbolic Spaces*, RIMS Kokyuroku Bessatsu, B66 (eds M. Fujii, N. Kawazumi and K. Ohshika) (Research Institute for Mathematical Sciences (RIMS), Kyoto, 2017), 167–184.
- [25] A. Putman, 'Lectures on the Torelli group'. www3.nd.edu/~andyp/teaching/2014SpringMath541/TorelliBook.pdf.
- [26] S. K. Winker, 'Quandles, knots invariants and the n -fold branched cover', PhD Thesis, University of Illinois at Chicago, 1984.
- [27] D. N. Yetter, 'Quandles and Lefschetz fibrations', Preprint, 2002, [arXiv:math/0201270](https://arxiv.org/abs/math/0201270).
- [28] D. N. Yetter, 'Quandles and monodromy', *J. Knot Theory Ramifications* **12**(4) (2003), 523–541.
- [29] J. Zablow, 'Loops, waves, and an "algebra" for Heegaard splittings', PhD Thesis, City University of New York, 1999.
- [30] J. Zablow, 'Loops and disks in surfaces and handlebodies', *J. Knot Theory Ramifications* **12**(2) (2003), 203–223.
- [31] J. Zablow, 'On relations and homology of the Dehn quandle', *Algebr. Geom. Topol.* **8**(1) (2008), 19–51.

NEERAJ K. DHANWANI, Department of Mathematical Sciences,
Indian Institute of Science Education and Research (IISER) Mohali,
Sector 81, S. A. S. Nagar, P. O. Manauli, Punjab 140306, India
e-mail: neerajk.dhanwani@gmail.com

MAHENDER SINGH, Department of Mathematical Sciences,
Indian Institute of Science Education and Research (IISER) Mohali,
Sector 81, S. A. S. Nagar, P. O. Manauli, Punjab 140306, India
e-mail: mahender@iisermohali.ac.in