

Individual and group behavioural reactions of small delphinids to remote biopsy sampling

JJ Kiszka^{*†‡}, B Simon-Bouhet[†], F Charlier[§], C Pusineri[§] and V Ridoux[†]

[†] LIENSs (LIttoral, ENvironnement et Sociétés), UMR 6250, CNRS-Université de La Rochelle, 2, rue Olympe de Gouges, F-17000, La Rochelle, France

[‡] Direction de l'Environnement et du Développement Durable, Collectivité Départementale de Mayotte, BP 101 F-97600 Mamoudzou, Mayotte

[§] Office National de la Chasse et de la Faune Sauvage, Délégation Régionale Outre-Mer, Coconi, Mayotte

* Contact for correspondence and requests for reprints: jeremy.kiszka@wanadoo.fr/jeremy.kiszka@univ-lr.fr

Abstract

Biopsy sampling is an effective technique for collecting cetacean skin and blubber samples for various biological studies. However, determining the impact of this research practice is important, as it may vary between sites, species and equipment used. We examined the short-term behavioural reactions of four small (160–278 cm in length) delphinid species (*Stenella longirostris*, *S. attenuata*, *Tursiops aduncus* and *Peponocephala electra*) to remote biopsy sampling in the vicinity of the island of Mayotte (12°50'S, 45°10'E, SW Indian Ocean). Two scales of behavioural reactions were considered: i) the behavioural reaction of the individual, and ii) the reaction of the focal group to which the targeted individual belonged. Three main categories of behavioural responses were defined on the basis of character and duration: low, moderate and strong. This study underlines that biopsy sampling induces moderate reactions in individuals. No inter-specific variations of responses, at the scale of the individual or focal group, were observed. In other words, smaller delphinids were not more reactive than larger ones. No effect of group size was observed on the strength of behavioural reactions. However, it was clear that biopsy success during sampling sessions was higher in species with large group size. Finally, in the spinner dolphin (*S. longirostris*), we investigated whether initial behavioural state affected the level of reaction. Resting and socialising groups showed a stronger response than milling and travelling groups. This study confirms the limited impact of remote biopsy sampling in small delphinids, especially in the spinner dolphin. However, as a precautionary approach, wherever possible, biopsy sampling of milling and travelling dolphins may be preferable.

Keywords: animal welfare, behavioural impact, dolphins, group reactions, individual reactions, remote biopsy sampling

Introduction

In wildlife studies, a number of invasive techniques may be used to collect biological samples in order to answer a variety of questions which may be of particular relevance in terms of management and/or conservation. It is crucial that the impact of such practices is quantitatively assessed and managed, as the process of sample collection may have a negative impact at both an individual or population level, over a range of scales (eg injuries, individual stress, individual/group displacement, behavioural change, etc).

The use of skin and blubber biopsy samples from free-ranging cetaceans is a widespread and highly effective technique for answering many questions, including those relating to population genetics (stock identity, social organisation, population size, phylogeny, genetic connectivity: Amos & Hoelzel 1990; Bérubé *et al* 1998), feeding ecology and trophic relationships using stable isotope and

fatty acid analyses (Herman *et al* 2005; de Stephanis *et al* 2008; Gross *et al* 2009), and pollutant analysis (Godard *et al* 2004). In order to collect samples, modified crossbows, rifles and hand-held biopsy poles have been used, both for large and small cetaceans, including delphinids (Weinrich *et al* 1991; Krützen *et al* 2002; Bilgmann *et al* 2007). The behavioural effect of biopsy sampling has been investigated in large whales, such as right whales (*Eubalaena glacialis* and *E. australis*: Brown *et al* 1991; Best *et al* 2005), humpback whales (*Megaptera novaeangliae*: Weinrich *et al* 1991; Clapham & Mattila 1993), other large balaenopterid whales (Gauthier & Sears 1999), and delphinids such as short-beaked common dolphin (*Delphinus delphis*: Bearzi 2000), bottlenose dolphin (*Tursiops* spp: Krützen *et al* 2002; Bilgmann *et al* 2007; Gorgone *et al* 2008) and Indo-Pacific humpback dolphin (*Sousa chinensis*: Jefferson & Hung 2008).

The International Whaling Commission considers biopsy sampling to be acceptable, since no long-term effects (change of behaviour) have been observed for individuals and populations (International Whaling Commission 1991). Levels of short-term reactions to biopsy sampling could potentially vary among species, populations and individuals. However, for both small and large cetaceans, the behavioural impact of biopsy sampling is generally considered to be low. Animal responses can be characterised as reactions to a brief noxious stimulus of low-to-moderate amplitude (Weinrich *et al* 1992; Best *et al* 2005; Bilgmann *et al* 2007; Jefferson & Hung 2008). In small cetaceans, an incidence of death has been reported in a short-beaked common dolphin, underlining that remote biopsy sampling is not without risk (Bearzi 2000). Consequently, the use of less-invasive sampling techniques may be preferred. Other methods include skin swabbing and faecal sampling (Harlin *et al* 1999; Parsons *et al* 1999). However, these techniques provide a limited amount of material, and DNA may not be of sufficient quality to undertake multiple markers analyses and other analyses (such as pollutant analyses, for example). Biopsy sampling is generally preferred for molecular genetic studies (Parsons *et al* 2003) and other analyses, such as those involving stable isotopes (Gross *et al* 2009). In addition, the use of remote techniques, using a gun or a crossbow, is more effective than a pole system for studies of population structure and parentage because animals can be sampled even if they do not bowride. Remote sampling also allows the individual identification of targeted dolphins (Bilgmann *et al* 2007). Proper identification of bowriding animals is generally not possible as it is dependent upon achieving an appropriate angle for photography.

The objective of this study was to characterise the short-term reactions of four small delphinid species to remote biopsy sampling: the spinner dolphin (*Stenella longirostris*, 160–208 cm), the pantropical spotted dolphin (*S. attenuata*, 160–260 cm), one of the smallest delphinids, the Indo-Pacific bottlenose dolphin (*Tursiops aduncus*, 230–270 cm) and the melon-headed whale (*Peponocephala electra*, 240–278 cm), one of the least known delphinids. This study provides, as far as we are aware, the first information on the effects of biopsy sampling in these species.

In order to collect skin and blubber samples for stable isotope, genetic and histopathological investigations, remote biopsy sampling was conducted from December 2004 to October 2008. Levels of behavioural reactions were recorded at two different levels: i) the behavioural reaction of the individual, sampled dolphin; and ii) the behavioural reaction of the focal group to which the targeted animal belonged. This latter component has not been investigated in previous studies of any other cetacean, as far as we are aware, and allows for an understanding of the impact of remote biopsy sampling on a broader scale, ie groups as opposed to merely targeted individuals.

Materials and methods

Study area

The island of Mayotte (45°10'E, 12°50'S), which is part of the Comoros archipelago, is located in the northern Mozambique Channel (SW Indian Ocean) between Madagascar and Southeast Africa (Figure 1). It covers an area of 376 km² and is composed of two main islands: the main inhabited island on the east and, on the barrier reef, a smaller inhabited island. The other islands are small islets dispersed throughout the lagoon. The island of Mayotte is characterised by a large diversity of marine mammals (22 species including 12 delphinids; Kiszka *et al* 2007). The most common species are the spinner dolphin, the pantropical spotted dolphin, the Indo-Pacific bottlenose dolphin and the melon-headed whale; all of which are resident throughout the whole year (Kiszka *et al* 2007).

Biopsy collection

From December 2004 to September 2008, small-boat-based cetacean surveys were conducted throughout the year in Mayotte waters in sea conditions not exceeding Beaufort 3. Observational efforts concentrated mainly on the lagoon and over the insular slope in adjacent waters of the barrier reef. Biopsy attempts were made opportunistically, when groups and individuals were easily approachable and when conditions were optimal (Beaufort < 2, dolphins closely approaching the boat). Optimal weather conditions allowed stability of the research boat and afforded a better chance of sampling the animals safely and successfully. Several types of boats were used: a 7-m catamaran equipped with two, four-stroke, 60-hp outboard engines; a 7-m mono hull boat equipped with two, two-stroke, 40-hp outboard engines; a 6.4-m cabin cruiser equipped with one, four-stroke, and 150-hp outboard engine; and a 10.8-m cabin cruiser equipped with two, four-stroke, 115-hp outboard engines. Biopsies were collected by using a crossbow (BARNETT Veloci-Speed® Class, 68-kg draw weight) with Finn Larsen (Ceta-Dart, Copenhagen, Denmark) bolts and tips (dart: 25 mm in length, 5 mm diameter). A conical plastic stopper caused the bolt to rebound after impact with the dolphin. When sufficiently close (3–10 m) to the research boat, the dolphins were hit below the dorsal fin. Focal groups/individuals were approached at speeds of 1–4 knots. Blubber and skin biopsy samples were preserved individually in 90% ethanol before shipping and subsequent analysis. Biopsy sampling was conducted under French scientific permit #78/DAF/2004 (September 10, 2004) and permit #032/DAF/SEF/2008 (May 16, 2008) after examination of the project by Conseil National de Protection de la Nature.

Behavioural observations

During biopsy-sampling sessions, an observer recorded the behavioural reactions of dolphins at two different scales: the targeted individual and the focal group with which the targeted individual was associated. The focal group was defined as a group of dolphins engaged in the same activity

and travelling in the same direction (Shane 1990). Three levels of behavioural reaction were defined for individuals and focal groups. These reactions followed Hooker *et al* (2000) and were adapted for the species investigated in this study: 1) No reaction — the individual and focal group continued to show the same behaviour as before the biopsy attempt; 2) Moderate reaction — the individual or the focal group modified its behaviour but gave no prolonged (> 5 min) evidence of behavioural disturbance: reactions included, eg acceleration, twitch and immediate dive and simple immediate dive (a dive was considered as a behavioural response to biopsy sampling when it lasted more than 5 min); and 3) Strong reaction — the individual or the focal group modified its behaviour in a succession of percussive behaviours (strong and short-lived reactions), including escape from the research boat of the individual and/or focal group (leaping, breaches, tail slaps).

Data analysis


We investigated the occurrence (events and their proportions) of reactions described above and factors responsible for the variability of reactions (group size, species, activity), at the scale of hit/targeted individuals as well as focal groups. Group size was defined prior to biopsy sampling as the number of animals at the surface within five body lengths of each other (Smolker *et al* 1992). The estimates of group size were more stochastic for spinner dolphins, pantropical spotted dolphins and melon-headed whales, as group size for these species was important (mostly > 50 individuals). Determining absolute group size was not possible for large groups of delphinids. The predominant behaviour was recorded as the activity displayed by the majority of the animals of the group during the first 10 min. These data were collected during scan sampling of the group (Mann 1999) using six different behavioural categories: travelling, milling, resting, feeding/foraging, playing and socialising (Shane 1990).

Analysis of individual behavioural reactions were differentiated when the animal was missed (the bolt did not reach the animal) or hit. An individual was considered as hit when the bolt reached the body. No differentiation was made between biopsy hit providing or not providing a sample. We tested how group size may affect individual behavioural reactions, especially for the most frequently sampled species, the spinner dolphin. For this species, we also investigated the effect of initial behavioural state on the levels of reaction and the long-term effect of biopsy sampling. In this later case, we hypothesised that avoidance behaviour would increase across the study period. Significance of this increase has been tested using a Pearson's correlation. For comparisons, Fisher exact tests, Kruskal-Wallis tests and contingency table analyses were performed using Rv2.10.0 (R Development Core Team 2009).

Results

Biopsy sampling was undertaken from December 2004 to September 2008 (n = 271 attempts, n = 193 samples). Four species (spinner dolphin, pantropical spotted dolphin, melon-

Figure 1




Location of study area.

headed whale and Indo-Pacific bottlenose dolphin) constituted 96% of the biopsies sampled (n = 259 attempts, n = 181 samples). Other species included the Fraser's dolphin (*Lagenodelphis hosei*: n = 7), the common bottlenose dolphin (*Tursiops truncatus*: n = 2), the Indo-Pacific humpback dolphin (n = 2) and the short-finned pilot whale (*Globicephala macrorhynchus*: n = 1), but these data have not been included in this study. Three types of vessel were used to undertake biopsy sampling, but no significant differences in individual behavioural reactions were found between boat types (all species combined, $\chi^2 = 3.7$, df = 6; $P = 0.391$).

Among the four species, no significant inter-specific differences in reactions were recorded, both at the scale of the individual (Fisher exact test; $P = 0.9$) and the group ($P = 0.643$). Sampling success (a hit) varied between species from 65 to 78% (Table 1). On 34 occasions overall, the hit was successful but no sample was retained in the biopsy tip. Individual behavioural reactions to remote biopsy sampling were recorded on 252 occasions (180 hits, 72 misses), while focal group behavioural reactions were recorded on 271 occasions (193 hits, 78 misses). There were no statistical differences between individual behavioural reactions between biopsy hits and misses (all species

Table 1 Number of attempts, biopsy samples collected and sampling success in delphinids sampled around the island of Mayotte from December 2004 to September 2008.

Species	Attempts (n)	Samples (n)	Success (%)	Biopsy sessions (n)	Mean group size	Average biopsies per session
<i>Stenella longirostris</i>	137	96	70	30	70.5	3.2
<i>Stenella attenuata</i>	77	50	65	20	78.5	2.5
<i>Peponocephala electra</i>	23	18	78	5	31.0	3.6
<i>Tursiops aduncus</i>	22	17	77	15	6.3	1.1

Figure 2Individual behavioural reactions of delphinids (*Stenella longirostris*, *S. attenuata*, *Tursiops aduncus* and *Peponocephala electra*) to a biopsy hit or miss.

combined, Fisher exact test, $P = 0.068$). Similarly, at the scale of focal groups, no significant differences between biopsy hits and misses were found (all species combined, KW test; $H = 0.702$; $df = 1$; $P = 0.402$).

At the individual scale, 94% of individual reactions were moderate, ie twitch and immediate dive, and simple immediate dive (Figure 2, Table 2). Strong reactions (tail slap, leaping, successive breaches and escape) only represented 2% of behavioural responses of individual dolphins. Escape and leaping was only observed in spinner and spotted dolphins. Increase in speed was observed once in a bottlenose dolphin group (Table 2).

Group behavioural reactions were frequent (54% of sampling sessions), with dive being the commonest moderate reaction (45%, Figure 3, Table 2). Strong reactions of focal groups were rare, representing only 4% of responses. These strong reactions consisted of increased swimming speed or escape (Table 2, Figures 2 and 3).

We did not find any correlation between group size and behavioural reactions (Fisher exact test; $P = 0.431$). However, there is a clear relationship between the mean


specific group size and the mean number of biopsies collected per biopsy session (Table 1). The average number of biopsies collected during each session was lowest for the Indo-Pacific bottlenose dolphin, which had the lowest mean group size (Table 1).

On six occasions, hit dolphins were observed bow-riding just after having been sampled (fresh wound of the biopsy hit observed below the dorsal fin or in adjacent areas). These cases were observed in the pantropical spotted dolphin ($n = 2$ events), spinner dolphin ($n = 2$) and Indo-Pacific bottlenose dolphin ($n = 2$). During sampling sessions, significant signs of avoidance of the research vessel by groups were observed on a number of occasions: in spinner dolphins ($n = 2$, after one and four biopsy attempts) and in melon-headed whales ($n = 2$, after one and six biopsy attempts).

We hypothesised that group reactions to biopsy sampling would differ according to activity (milling/travelling, resting, socialising, and playing). We tested this for spinner dolphins, as the dataset for that species was the largest. A significant difference was found between group reactions and initial

Table 2 Individual and focal group behavioural reactions of *Stenella longirostris*, *S attenuata*, *Peponocephala electra* and *Tursiops aduncus* to remote biopsy sampling (numbers represent events).

Reactions	<i>Stenella longirostris</i>	<i>Stenella attenuata</i>	<i>Peponocephala electra</i>	<i>Tursiops aduncus</i>
<i>Individual reactions</i>				
Twitch and dive	83	21	8	10
Successive breaches	1	1	0	0
Tail slap	2	0	0	0
Escape (leaping)	1	0	0	0
No reaction	1	0	0	0
<i>Group behavioural reactions</i>				
Dive	41	19	7	9
Increase swimming speed	2	0	0	0
Escape	1	1	0	1
No reaction	40	28	10	6

Figure 3Focal group behavioural reactions of delphinids (*Stenella longirostris*, *S attenuata*, *Tursiops aduncus* and *Peponocephala electra*) to a biopsy hit or miss.

behavioural states in which spinner dolphin groups were engaged (Fisher exact test; $P = 0.041$). Spinner dolphins predominantly showed a stronger response to biopsy sampling when resting and socialising. When milling and travelling, reactions were moderate. We did not observe changes in dolphin reactions (increased avoidance behaviour) to the research vessel prior to biopsy sampling over the study period (nearly four years, Pearson's correlation; $r = 0.324$; $P > 0.05$).

Discussion

In this study, we observed the behavioural reactions of four delphinid species to remote biopsy sampling. The biopsy success rate was 65–78%, which is consistent with previous studies. This was mostly due to the high accessibility of the

targeted species which generally came close to the research vessel, especially dolphins of the genus *Stenella*, often coming to ride waves created by the bow of the boat. No significant inter-species differences were found in reactions to remote biopsy sampling. Indeed, the smallest species (spinner and pantropical spotted dolphins) did not have a higher occurrence of moderate reactions than larger ones (Indo-Pacific bottlenose dolphin and melon-headed whale), as might have been expected. However, the strongest reactions, such as breaches and escaping, occurred (but were very rare). Such extreme reactions were only observed in the smaller species, especially spinner and pantropical spotted dolphins. However, due to the small sample size for Indo-Pacific bottlenose dolphins and melon-headed whales, we cannot rule out the

possibility that these species could also react strongly, in the same manner as spinner and spotted dolphins.

The mean number of biopsies per session was greater for species with a larger mean group size, ie melon-headed whales, spinner and pantropical spotted dolphins. In larger groups, animals are more accessible for biopsying, as there are more individuals to choose from. This is likely to not merely be a function of the behaviour or the group reaction, but also a result of the higher number of individuals.

Despite the fact that we used three different types of boats, no differences in reactions were found between vessels. These differences have been documented in other studies, with generally stronger reactions when smaller boats were used (Bilgmann *et al* 2007). However, in this later study, sampling was carried out on bowriding dolphins, and the boat types and length differed to a much larger extent than in this study.

Delphinids around Mayotte exhibited short-term behavioural reactions to biopsy attempts, characterised by acceleration, twitch and immediate dive and simple immediate dive. Strong reactions to biopsy sampling were previously recorded in common bottlenose dolphins (Parsons *et al* 2003). Conversely, reactions of common bottlenose dolphins appear to be minimal in other areas such as in the eastern US (Gorgone *et al* 2008). Dolphins of all species sampled react in a similar fashion to biopsy hits and misses. This has been previously documented for other species such as the Indo-Pacific humpback dolphin (Jefferson & Kung 2008), meaning that the hit of the bolt on the water has a significant effect on the reactions of dolphins at the immediate proximity of the impact. In this study, focal groups were frequently impacted by biopsy sampling, meaning that remote biopsy sampling does have a broader effect on small cetaceans, ie on adjacent individuals within the group. This effect was also greater in species consisting of small groups, ie Indo-Pacific bottlenose dolphins, as the biopsy success decreased during sampling sessions for such species. The group behavioural reactions of spinner and pantropical spotted dolphins were relatively low, apparently because they formed larger aggregations. However, results underlined that there is variability in reactions according to initial behavioural state. Indeed, for the spinner dolphin, we observed that the animals had stronger reactions to remote biopsy sampling when resting and socialising. When milling and travelling, reactions were more moderate. This suggests that remote biopsy sampling should be conducted preferably during travelling and milling activities.

Animal welfare implications

Overall, conducting remote biopsy sampling is effective on small delphinids and induces a limited short-term (less than 5 min) behavioural impact on hit and missed individuals, including those in the smallest delphinid species (especially dolphins of the genus *Stenella*). However, we also observed that biopsy sampling does not only impact hit individuals, but groups to which the targeted individual belongs. No long-term effect of biopsy sampling was observed, such as an increase in avoidance of research vessels. This confirms that the method has no long-term impact on the animals.

However, as a precautionary approach, our findings suggest that it may be preferable to conduct biopsy sampling when the animals are milling or travelling. It is critical however that we reconsider the practice of biopsy sampling to answer scientific questions.

Acknowledgements

This research has been funded by the Collectivité Départementale de Mayotte and the Ministère de l'Ecologie et du Développement Durable (French Ministry of Environment). We thank Robin Rolland, Alban Jamon, Wilfrid Fousse, Ismaël Oussené (DAF) and the personnel of Brigade Nature (CDM/ONCFS) for assistance in the field in Mayotte. Special thanks are addressed to Didier Fray (CDM), our main boat pilot, for his patience and perseverance in the field. We warmly thank Tom Jefferson and William Perrin (NOAA Fisheries) for their helpful comments on the early version of the manuscript, and the two anonymous reviewers for their constructive comments and corrections.

References

- Amos W and Hoelzel AR** 1990 DNA fingerprinting cetacean biopsy samples for individual identification. *Report of the International Whaling Commission 12*: 79-85
- Bearzi G** 2000 First report of a common dolphin (*Delphinus delphis*) death following penetration of a biopsy dart. *Journal of Cetacean Research and Management 2*: 217-221
- Bérubé M, Aguilar A, Dendanto D, Larsen F, Notarbartolo di Sciara G, Sears R, Sigurjónsson R, Urban-RJ and Palsbøl PJ** 1998 Population genetic structure of North Atlantic, Mediterranean and Sea of Cortez fin whales, *Balaenoptera physalus* (Linnaeus 1758): analysis of mitochondrial and nuclear loci. *Molecular Ecology 7*: 585-599
- Best PB, Reeb D, Rew MB, Palsbøl PJ, Schaeff C and Brandão A** 2005 Biopsying southern right whales: their reactions and effect on reproduction. *Journal of Wildlife Management 69*: 1171-1180
- Bilgmann K, Griffiths OJ, Allen SJ and Möller LM** 2007 A biopsy pole system for bow-riding dolphins: sampling success, behavioral responses, and test for sampling bias. *Marine Mammal Science 23*: 218-225
- Brown MW, Kraus SD and Gaskin DE** 1991 Reaction of north Atlantic right whales (*Eubalaena glacialis*) to skin biopsy sampling for genetic and pollutant analysis. In: Hoelzel AE and Donovan GP (eds) *Genetic Ecology of Whales and Dolphins* pp 81-89. The International Whaling Commission: Cambridge, UK
- Clapham PJ and Mattila DK** 1993 Reactions of humpback whales to skin biopsy sampling on a West Indies breeding ground. *Marine Mammal Science 9*: 382-391
- De Stephanis R, Garcia-Tiscar S, Verborgh P, Esteban-Pavo R, Pérez S, Minvielle-Sebastia L and Guinet C** 2008 Diet of social groups of long-finned pilot whales (*Globicephala melas*) in the Strait of Gibraltar. *Marine Biology 154*: 603-612
- Gauthier J and Sears R** 1999 Behavioral response of four species of balaenopterid whales to biopsy sampling. *Marine Mammal Science 15*: 85-101
- Godard CAG, Smolowitz RM, Wilson JY, Payne RS and Stegeman JJ** 2004 Induction of cetacean cytochrome P450IA1 by β -naphthoflavone exposure of biopsy slices. *Toxicological Sciences 80*: 268-275

- Gorgone AM, Haase PA, Griffith ES and Hohn A** 2008 Modelling response of target and nontarget dolphins to biopsy darting. *Journal of Wildlife Management* 72: 926-932
- Gross A, Kiszka J, Van Canneyt O, Richard P and Ridoux V** 2009 A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, southwest Indian Ocean. *Estuarine, Coastal and Shelf Science* 84: 367-374
- Harlin AD, Würsig B, Baker CS and Markowitz TM** 1999 Skin swabbing for genetic analysis: application to Dusky dolphins (*Lagenorhynchus obscurus*). *Marine Mammal Science* 15: 409-425
- Herman DP, Burrows DG, Wade PR, Durban JW, Matkin CO, LeDuc RG, Barret-Lennard LG and Krahn MM** 2005 Feeding ecology of eastern North Pacific killer whales *Orcinus orca* from fatty acid, stable isotope, and organochlorine analyses from blubber biopsies. *Marine Ecology Progress Series* 302: 275-291
- Hooker SK, Baird RW, Al-Omari S, Gowans S and Whitehead H** 2000 Behavioral reactions of northern bottlenose whales (*Hyperoodon ampullatus*) to biopsy darting and tag attachment procedures. *Fishery Bulletin* 99: 303-308
- International Whaling Commission** 1991 Report of the *ad hoc* working group on the effect of biopsy sampling on individual cetaceans. In: Hoelzel AE and Donovan GP (eds) *Genetic Ecology of Whales and Dolphins* pp 23-27. The International Whaling Commission: Cambridge, UK
- Jefferson TA and Hung SK** 2008 Effects of biopsy sampling on Indo-Pacific humpback dolphins (*Sousa chinensis*) in a polluted coastal environment. *Aquatic Mammals* 34: 310-316
- Kiszka J, Ersts PJ and Ridoux V** 2007 Cetacean diversity around the Mozambique Channel island of Mayotte (Comoros archipelago). *Journal of Cetacean Research and Management* 9: 105-109
- Krützen M, Barre LM, Möller LM, Heithaus MR, Simms C and Sherwin WB** 2002 A biopsy system for small cetaceans: darting success and wound healing in *Tursiops* spp. *Marine Mammal Science* 18: 863-878
- Mann J** 1999 Behavioral sampling methods for cetaceans: a review and critique. *Marine Mammal Science* 15: 102-122
- Parsons KM, Dallas JF, Claridge DE, Durban JW, Balcomb KC, Thompson PM and Noble LR** 1999 Amplifying dolphin mitochondrial DNA from faecal plumes. *Molecular Ecology* 8: 1766-1768
- Parsons KM, Durban JW and Claridge DE** 2003 Comparing two alternative methods for genetic sampling of small cetaceans. *Marine Mammal Science* 19: 224-231
- R Development Core Team** 2009 R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing: Vienna, Austria
- Shane SH** 1990 Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. In: Leatherwood S and Reeves RR (eds) *The Bottlenose Dolphin* pp 245-265. Academic Press: San Diego, USA
- Smolker RA, Richards AF, Connor RC and Pepper JW** 1992 Sex differences in patterns of association among Indian Ocean bottlenose dolphins. *Behaviour* 123: 38-69
- Weinrich MT, Lambertsen RH, Baker CS, Schilling MR and Belt CR** 1991 Behavioral responses of humpback whales *Megaptera novaeangliae* in the southern Gulf of Maine to biopsy sampling. In: Hoelzel AR and Donovan GP (eds) *Genetic Ecology of Whales and Dolphins* pp 91-98. International Whaling Commission: Cambridge, UK
- Weinrich MT, Lambertsen RH, Belt CR, Shilling MR, Iken JH and Syrjala SE** 1992 Behavioral responses of humpback whales *Megaptera novaeangliae* to biopsy procedures. *Fishery Bulletin* 90: 588-598