
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSIDE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

Contents

	Pages
Effect of the organophosphorous insecticide, chlorpyrifos (Dursban), on growth, fecundity and mortality of <i>Biomphalaria alexandrina</i> and on the production of <i>Schistosoma mansoni</i> cercariae in the snail. W. L. F. IBRAHIM, P. FURU, A. M. IBRAHIM and N. Ø. CHRISTENSEN	79-88
Seasonal patterns in the transmission of <i>Schistosoma haematobium</i> in Attaouia, Morocco. K. KHALLAAYOUNE and H. LAAMRANI	89-95
Sites of encystment by the metacercariae of <i>Echinoparyhium recurvatum</i> in <i>Lymnaea peregra</i> . M. E. ADAM and J. W. LEWIS	96-99
Growth, development and allometry of <i>Philophthalmus nocturnus</i> in the eyes of domestic chicks. V. G. M. SWARNAKUMARI and R. MADHAVI	100-107
Effect of concurrent infection with <i>Muellerius capillaris</i> on the development of redial generations of <i>Fasciola hepatica</i> in <i>Lymnaea truncatula</i> . P. HOURDIN, D. RONDELAUD and J. CABARET	108-117
Clinical manifestations of taeniasis in Taiwan aborigines. P. C. FAN, W. C. CHUNG, C. Y. LIN and C. H. CHAN	118-123
Comparative cytotoxicity of secondary hydatid cysts, protoscoleces and <i>in vitro</i> developed microcysts of <i>Echinococcus granulosus</i> . D. JANSSEN, A. OSUNA, J. LAZUEN and P. H. DE RYCKE	124-131
Differential establishment and survival of <i>Hymenolepis diminuta</i> in syngeneic and outbred rat strains. A. ISHIIH, M. NISHIMURA and M. SANO	132-136
<i>In vivo</i> passage through calves of nematophagous fungi selected for biocontrol of parasitic nematodes. M. LARSEN, J. WOLSTRUP, S. A. HENRIKSEN, J. GRØNVOLD and P. NANSEN	137-141
Isozymic pattern of lactate dehydrogenase in cases of bancroftian filariasis. S. MISRA, T. M. MAHOPATRA and S. RATHAUR	142-146
Induction of protective immunity to <i>Brugia pahangi</i> in jirds by drug-abbreviated infection. Y. HORII, H. NAKANISHI, A. MORI, M. UEDA, K. KUROKAWA, M. ZAITSU, T. ODA and K. FUJITA	147-154
Protease resistant interleukin-3 stimulating components in excretory and secretory products from adult worms of <i>Strongyloides ratti</i> . T. ABE, Y. NAWA and K. YOSHIMURA	155-158
International Commission on Zoological Nomenclature	117