

RESEARCH ARTICLE

Child undernutrition and associated factors among children 6–23 months old in largely food insecure areas of rural Ethiopia

Zelalem Tafese* , Fekadu Reta, Biruk Mulugeta and Anchamo Anato 

School of Nutrition, Food Science and Technology, Hawassa University, P. O. Box 05, Hawassa, Ethiopia

(Received 15 November 2021 – Final revision received 1 June 2022 – Accepted 12 July 2022)

Journal of Nutritional Science (2022), vol. 11, e63, page 1 of 9

doi:10.1017/jns.2022.61

Abstract

Child malnutrition is the leading public health problem in Sub-Saharan Africa, resulting in poor health and economic losses. Ethiopia has one of the highest child undernutrition rates in the world that occurs to multifaceted factors, including food insecurity. Thus, we performed a cross-sectional study to assess the prevalence and risk factors for child undernutrition in largely food insecure areas of Ethiopia. Data were collected from 354 mother–child pairs from the Siraro district. Both bivariate and multivariate logistic regression was used for statistical analysis. Variables with a *P*-value of <0.05 in multivariate analysis were used to detect statistical significance at a 95 % confidence level. About 67 % of households are food insecure. The prevalence of stunting wasting and underweight were 42.7, 9.9 and 27.7 %, respectively. Female gender, higher age of the child (12–23 months *v.* 6–11 months), living in a household with five or more siblings, not getting therapeutic zinc supplement at least once, inadequate diet diversity, lack of growth monitoring service, and maternal own income increases the likelihood of child undernutrition. It can be concluded that child undernutrition is a severe public health problem in the study area. Improving primary healthcare services related to zinc supplementation, growth monitoring and promotion, and improving infant and child feeding practices can be considered as a strategy to address the problem.

Key words: Diet diversity: Food insecurity: Mothers income: Stunting: Underweight: Zinc supplement

Background

Widespread child undernutrition is one important problem in Africa⁽¹⁾ and Ethiopia^(2,3). According to the World Health Organization (WHO), wasting, stunting and being underweight are the main forms of undernutrition defined as *z*-scores less than -2 standard deviations of weight for height, height for age and weight for age, respectively⁽⁴⁾. Globally, 144 million children under 5 were stunted in 2019⁽⁵⁾. Among the three forms of the above-mentioned undernutrition, stunting is a devastating result of poor nutrition during fetal development and in early childhood⁽⁶⁾. At the global level, more than one in four children under the age of 5 years are stunted and Sub-Saharan Africa and

South Asia suffer the heaviest burden, with 75 % of the world's stunted children⁽⁷⁾.

Undernutrition in children occurs from multifaceted factors, including food insecurity^(8–10). One of the underlying causes of undernutrition in the conceptual framework of UNICEF is mainly linked to poor dietary intake⁽¹¹⁾. This indicates a high vulnerability to undernutrition among children from food-insecure households. Although food insecurity affects the nutritional status of the general population, its effect is more serious on the venerable segment of the population mothers and children⁽¹²⁾.

Ethiopia is one of the countries reported with the worst food crises in 2018⁽¹³⁾. Food insecurity in the Ethiopian context is a serious problem and the majority of the country's population

* Corresponding author: Zelalem Tafese, email wudasiez@gmail.com

Abbreviations: COVID-19, coronavirus disease 2019; DDS, diet diversity; HFIAS, household food insecurity access scale; IYCF, infant and young child feeding; MAD, minimum acceptable diet; WHO, World Health Organization



lives in rural areas^(14,15). Moreover, the number of food-insecure people in the country is increasing from year to year. For comparison 2.9 million in 2014 and 4.5 million in 2015 were estimated to be food insecure, and by the end of the same year, this figure had increased by more than twofold (10.2 million)⁽¹⁶⁾. According to the report of FAO, despite the ongoing assistance in Ethiopia, an estimated 8 million people were severely food insecure and the situation worsens between July and September, due to erratic rains, conflict and high food prices⁽¹⁷⁾.

Food insecurity and malnutrition were one of the public health problems in Ethiopia and throughout Sub-Saharan Africa⁽¹⁸⁾. In these regions, a high number of children were reported to be suffering from undernutrition⁽¹⁹⁾, and an increased risk of food shortage related to a variety of factors⁽²⁰⁾. The country is facing multiple, underlying vulnerabilities, for child undernutrition including food insecurity^(21,22). In addition to several factors^(10–14), the emerging global problem, the COVID-19 is likely to exacerbate the existing food security problems in these regions⁽²³⁾. The situation may be exacerbated in many ways including being an obstacle to imports and transportation problems, related to a combination of lockdowns and travel restrictions. And hence it is logical that this may exacerbate the already considerable burden of malnutrition and food insecurity, and the worst effect is expected among the poor communities⁽²⁴⁾. Because of these and related conditions, the country is at risk of secondary impacts of COVID-19 such as increased cases of acute and chronic malnutrition⁽²⁵⁾; although data are reported inadequate on the socio-economic impacts of COVID-19 in settings vulnerable to food insecurity⁽²⁶⁾.

During our data collection, the Ethiopian government began a contact tracing and isolating those who tested positive for the virus, closed schools, banned all public gatherings and sporting activities, and recommended social distancing a few days after the report of the first case of COVID-19 in March 2020. Although the travel restrictions can be effective in minimising the spread of the virus, they may play a negative role in the economy of the country⁽²⁷⁾. Therefore, the present study aimed to assess the food insecurity situation, nutritional status and risk factors among children 6–23 months during the emergence of the pandemic COVID-19 enabling us to anticipate its effect in this poor rural setting.

Methods

Study area and design

A community-based cross-sectional study was conducted in the Siraro district, East Shoa Zone of the Oromia region from March to May 2020. Siraro district is located 322 km southwest of the capital Addis Ababa. The population of the district is estimated at 213 741 of which, 106 870 are females. The total households of the district were reported at 44 531. Administratively Siraro District is divided into 32 kebeles (smallest administrative unit in Ethiopia) (28 rural and 4 towns). The mean annual temperature of the district is found between 13 and 25°C. However, there is a slight variation in temperature from month to month. The district is among the most impacted by climate variability-induced hazards in the region,

and one of the largely food-insecure districts targeted for social security programmes by the government of Ethiopia⁽²⁸⁾.

Sample size and sampling procedure

Children 6–23 months of age with their mothers living in the Siraro district were the study population. The sample size was calculated using the single population proportion formula with assumptions of 32 % stunting prevalence in Ethiopia⁽²⁹⁾, 5 % marginal error, 95 % confidence level and 5 % nonresponse rate. A total of 371 mother–child pairs were included in the study. The numbers of subjects were allocated proportionally from the kebeles based on the total number of households with 6–23 months children in each kebele. Study subjects were then selected by simple random sampling using a list of households in each kebele.

Data collection tools and techniques

We used the survey research method to collect the necessary information from sample respondents by a semi-structured questionnaire administered in a local language (Afan Oromo) as a major type of data collection method. These data were collected focusing on socio-demographic, economic, food insecurity status, maternal characteristics (antenatal and postnatal care), dietary, anthropometric and child morbidity characteristics (respiratory illness, diarrhoea and ear infection). The demographic data included the age and gender of the children, and household size (number of residents in a household). Child mothers or caretakers were asked to provide information on the child's age which was confirmed using child immunisation cards where available. Where cards were unavailable, the mothers or caretakers were asked to recall or use references to calendar events. The nutritional status of the child was assessed by anthropometric measurements undertaken in all eligible respondents in the selected households including height/length and weight for children standard categories of nutritional status are reported according to the WHO classification of anthropometric measurements cut-offs⁽³⁰⁾. After removing shoes and extra clothing, the child weight was measured to the nearest 0.1 kg using a calibrated SECA electronic balance with a measuring range of 25 kg. Instrument calibration was checked before weighing each child and the weighing scale was tested daily against a standard weight for accuracy. Height and length were measured to the nearest 0.1 cm using the UNICEF wooden height and length boards while weight was assessed to the nearest 0.1 kg using a calibrated SECA electronic balance.

The dietary diversity score (DDS) was developed from a single-pass 24 h recall by asking mothers about all foods the child had consumed for meals and snacks in the 24 h before the survey. The data collector wrote a list of the foods consumed; the numbers of meals and snacks were summarised and the foods consumed were subdivided into the seven standardised food groups after completing the interview. The consumption of any amount of food from a food group was sufficient for it to be included. The seven food groups were (1) cereals, roots and tubers; (2) legumes and



nuts; (3) dairy products; (4) flesh foods (any meat, fish or poultry product); (5) eggs; (6) vitamin A-rich vegetables and fruits and (7) other fruits and vegetables. Consuming ≥ 4 of the seven standardised food groups was labelled as adequate diversity and < 4 groups were inadequate. The dietary diversity score was computed using Mean \pm SD and children who scored less than and more than four food groups were also reported⁽³¹⁾. The meal frequency of the child was determined by asking the mother how many times the child took solid, semisolid or soft foods in the 24 h preceding the survey. Accordingly, ≥ 2 times for breastfed infants aged 6–8 months, ≥ 3 times for breastfed children aged 9–23 months and 4 times for non-breastfed children aged 6–23 months were considered to mean the children received the minimum meal frequency⁽³²⁾.

The Household Food Insecurity Access Scale (HFIAS) is a continuous measure of the degree of food insecurity (access) in the household in the past 4 weeks (30 days). The total HFIAS can range from 0 to 27, indicating the degree of insecure food access. For the present study, it was assessed by classifying it as food secure if it had not experienced any food insecurity conditions or had rarely worried about not having enough food, whereas food-insecure households were categorised as mild, moderate and severe following the guidelines. The HFIAS was used to measure the status of food insecurity. This scale categorises households into four levels of household food insecurity: food secure, mild, moderate and severely food insecure. This was proposed by the Food and Nutrition Technical Assistance (FANTA)⁽³³⁾. This instrument is a simple and valid tool to measure the access component of household food insecurity⁽³⁴⁾. The research instrument was pre-tested with 5 % of the total sample size out of the study area. This instrument was assessed for clarity, time to complete, understandability and completeness. Some questions were re-formed and re-ordered to carry out the objectives of the study and interview respondents smoothly. Adequate training was given to data collectors and supervisors on data collection techniques by the lead author. The data collectors administered the questionnaire privately to ensure confidentiality

Statistical analysis

The analysis was done using STATA 14 (Stata/SE 14) statistical package. Frequency, percentage, mean and standard deviation were computed from continuous variables. Variance inflation factor (VIF) was used to check collinearity and non-collinear variables were included in the independent binary logistic regression model. Variables with a $P < 0.05$ in the multivariable logistic regression analysis were used to declare statistical significance with a 95 % confidence interval.

Results

Socio-demographic characteristics

Three hundred fifty-four study participants were involved in this study of 371 participants planned to be included, with a response rate of 95 %. Nearly half of the participant children were females (Table 1). The mean (\pm SD) age of the children was 14.5 (± 4.6)

mo and the mean family size was 4.9 (± 1.8) persons. More than 14 % of households had three or more under-five children. Nearly 67 % of the households were food insecure, out of which 30 % were severely food insecure (Table 4).

Maternal characteristics

The majority (92 %) of mothers reported the pregnancy for the first child within the age range of 15–26 years (Table 2). Nearly 26 % of mothers gave five or more live births and 21 % of child mothers had more than three antenatal care (ANC) visits for their most recent pregnancy. Four percent of mothers gave their most recent birth at a health facility. More than 50 % of participant mothers did not ever have postnatal care (PNC) for their most recent delivery. Furthermore, more than 95 % of mothers did not get postpartum vitamin A supplements after this delivery as recommended by the World Health Organization⁽³⁵⁾.

Child feeding practice

Fifty-three percent of children started complementary food at 6 months of age (Table 2). Nearly 20 % of children had adequate diet diversity reported from the 24 h dietary

Table 1. Socio-demographic and anthropometric measures of participants from Siraro district, Ethiopia, 2020 (n 354)

Variable name	Percentage (%) or Mean (sd)
Age of the child in months	
6–11	27.8
12–23	72.2
Sex of the child	
Male	50.9
Female	49.2
Marital status of the mother	
Not married	6.78
Married/living together	86.2
Divorced/Widowed/Separated	7.08
Educational status of mother	
Secondary school and above	64
Primary school (1–8)	30.5
No formal education	5.4
Occupation of mother	
Housewife	91.5
Farmer	3.4
Other employment	0.85
Merchant	0.85
Daily labourer	3.3
Mother had own income	32
Income of mother/month (1 US dollar = 32 Eth.birr)	
≥ 500 Eth.birr	15
< 500 Eth.birr	85
Family size	4.97 \pm 1.77
Number of < 5 years children	1.69 \pm 0.94
Stunted	42.7
Wasted	9.9
Underweight	27.7
LAZ/HAZ	–1.69 \pm 1.75
WAZ	–1.24 \pm 1.23
WLZ	–0.50 \pm 1.49

LAZ, length for age z-score; sd, standard deviation; WAZ, weight for age z-score; WLZ, weight for length z-score.



Table 2. Child feeding practices, and health characteristics of participants from Siraro district, Ethiopia, 2020 (*n* 354)

Variable name	Percentage (%) or Mean (sd)
Age complementary food started in months	
<4 months	3.4
4–6 months	4.2
At 6 months	53.4
Beyond 6 months	38.9
Child met minimum acceptable diet	13
Child had respiratory illness in the last 2 weeks	43.8
Child had diarrhoea in the last 2 weeks	38.1
Child had zinc at least for at least one prior diarrhoeal episode	15.3
Child had ear infection in the last 2 weeks	20.1
Child immunised for age	89.9
Child received vitamin A supplement in last 6 months	69.8
Used iodised salt for complementary food preparation	52.8
Age of the mother at first pregnancy in years	18.36 ± 3.30
Total number of births	3.35 ± 1.93
Number of ANC visits	2.41 ± 1.31
Number of PNC visits	1.12 ± 1.32
Delivery place of the recent birth	
Health facility	4.5
Home with healthcare workers	47.2
Home with family	48.3
Mother had vitamin A supplement during the recent birth	3.11
Mother had iron/folate supplement during the recent pregnancy	58.8

Table 3. Dietary diversity score of infants and young children (6–23 months) from Siraro district, Ethiopia, 2020 (*n* 354)

Food groups recalled by mother for previous day and night	<i>N</i>	%
Starchy staples	336	94.9
Legumes and nuts foods	215	60.7
Dairy products (milk, yoghurt, cheese)	91	25.7
Flesh foods (meat, fish, poultry, liver/organ meats)	53	14.9
Eggs	88	24.9
Vitamin A-rich (yellow, green and red coloured) vegetables and fruits	66	18.6
Other fruits and vegetables	38	10.6
Dietary diversity score, overall		
Mean ± sd	2.51 ± 1.32	
Min and Max, respectively	1–7	
Scored less than four food groups	280	79.1
Scored four and more food groups	74	20.9

information (Table 3). The majority of children (94.9 %) are fed cereal-based foods. Only 15 % had flesh foods (meat, fish, poultry, liver/organ meats). The egg was consumed nearly by 25 % and vitamin A-rich (yellow, green and red coloured) vegetables and fruits by 18.6 %. Nearly 13 % of participant children met the minimum acceptable diet criteria, 53.1 % of participant children ate animal source food within a week interval and 15.3 % of children had zinc at least for one prior diarrhoeal episode (Table 2). Participant mothers who used iodised salt for complementary food were 52.8 %.

Child health characteristics

In the 2 weeks preceding the survey, 44 % of mothers reported their children had respiratory infections, 20 % reported an ear infection, and 38 % reported their children experienced diarrhoea (Table 2).

Prevalence of child undernutrition

Totally, 42.7 % (95 % CI 37.5, 47.8) were stunted, 9.9 % (95 % CI 7.12, 13.4) wasted and 27.7 % (95 % CI 23.2, 32.6) underweight (Table 1). Stunting and underweight were common among children in the 12–23 months age group compared to the 6–11 months age (Table 4).

Factors associated with child undernutrition

The multivariate logistic regression model that adjusted for covariates (Table 5), identified significant associations with stunting at $P < 0.01$ for age, gender and zinc supplements as well as associations at $P < 0.05$ for parity, iodised salt intake and child diet diversity. Among these variables, more than fourfold higher odds of stunting were found among children in the 12–23 months group (AOR 4.02; 95 % CI 2.27, 7.12), compared to 6–11 months infants. Girls have more than 1.9 times higher odds of being stunted (AOR 1.84; 95 % CI 1.21, 3.08), compared to boys. Similarly, children who never had therapeutic zinc supplements for diarrhoea showed more than four folds higher odds of being stunted (AOR 4.93; 95 % CI 2.12, 10.92), compared to those who had at least once in life. Those children whose mothers had five or more times birth also showed nearly double fold higher odds of being stunted (AOR 1.95; 95 % CI 1.15, 3.30), compared to those whose mothers had less than five births. The adjusted odds of stunting among children who consumed less than four food groups in the 24 h before the survey was 1.92 (95 % CI 1.05, 3.49), compared to those who had four or more food groups. The adjusted model predicting wasting (Table 6) identified family size, growth monitoring and maternal income to be significantly associated ($P < 0.05$).

Being in a family size of five or more was significantly associated with wasting (AOR 2.40; 95 % CI 1.10, 5.23), compared to those children from a family size of less than five. Children who do not receive growth monitoring services have relatively higher odds to be wasted (AOR 0.43; 95 % CI 0.20, 0.92), compared to those who received. Likewise, children of mothers who have their income showed more than threefold higher odds of being wasted (AOR 3.57; 95 % CI 1.21, 10.53).

Regarding underweight, the multivariate logistic model (Table 7) detected an association of zinc supplements, and total number of birth ($P < 0.01$), and diet diversity ($P < 0.05$). Children from mothers with five or more number births showed more than two folds higher odds of being underweight (AOR 2.00; 95 % CI 1.19, 3.37) than children from mothers with less than five births. Likewise, children who had never received zinc supplements were >4 times more likely to be underweight (AOR 4.35, 95 % CI 1.66, 11.40),

**Table 4.** Distribution of household food insecurity access prevalence (HFIAP) Siraro district, Ethiopia, 2020 (n 354)

Questions	Frequency		
	Rarely	Sometimes	Often
1a	88	54	15
2a	92	66	17
3a	104	57	12
4a	57	54	10
5a	82	67	12
6a	82	63	13
7a	49	30	8
8a	37	25	5
9a	26	18	4
Clue to the severity conditions			
		Food secure /access/ = 32.8%	
		Mildly food insecure /access/ = 10.5%	
		Moderately food insecure /access/ = 26.3%	
		Severely food insecure /access/ = 30.5%	

compared to those who had at least once in life. Finally, more than two folds higher odds of being underweight were found among children who do not meet the minimum diet diversity (AOR 2.09, 95 % CI 1.06, 4.14), compared to those who had met.

Discussion

The prevalence of stunting is consistent with prior studies reported from Ethiopia^(36,37) but much higher than stunting rates reported in previous studies in 2014, 23.3 %⁽³⁸⁾, and in 2015 17.1 %⁽³⁹⁾. Although an association between household food insecurity and child undernutrition has been reported in previous studies from Bangladesh⁽⁴⁰⁾ and Pakistan⁽⁴¹⁾, our study did not show significant associations of this variable

with either stunting wasting or underweight. Quite likely a lack of statistical symmetry on the distribution of households in the present study contributed because the study specifically was conducted in a largely food-insecure area. A similar finding was also noted in our previous published study⁽⁴²⁾.

Our study revealed the risk of stunting increased with the age of the child. A similar finding was reported from prior studies in 2017 and 2018 from Ethiopia^(43,44). The plausible reason may be as children grow older they have greater energy needs. Besides, stunting reflects chronic malnutrition that can be manifested after long-term nutritional deficiency. The other factor associated with stunting in the present study was inadequate diet diversity, which was consistent with the finding from previous studies^(45,46). It is a known fact that adequate complementary feeding is a challenge for children aged 6–23

**Table 5.** Factors associated with stunting among 6–23 months children from Siraro district, Ethiopia, 2020^a

Variables	COR [95 % CI]	P-value	AOR [95 % CI]	P-value
Age of child				
12–23 months	3.75 [2.18, 6.45]	<0.001**	4.02 [2.27, 7.12]	<0.001**
6–11 months ^b	1		1	
Gender of child				
Female	1.81 [1.18, 2.77]	0.006*	1.94 [1.21, 3.08]**	0.005*
Male ^b	1		1	
Total number of births				
≥5	1.84 [1.14, 2.97]	0.012*	1.95 [1.15, 3.30]**	0.011*
<5 ^b	1		1	
Child had zinc supplement for diarrhoea at least once				
No	4.49 [2.12, 9.51]	<0.001**	4.93 [2.12, 10.92]**	<0.001**
Yes ^b	1		1	
Child's foods had iodised salt				
No	1.90 [1.24, 2.91]	0.003*	1.56 [0.97, 2.50]	0.062
Yes ^b	1		1	
Diet diversity of child				
<4 food groups	1.72 [1.00, 2.96]	0.047*	1.92 [1.05, 3.49]	0.048*
≥4 food groups ^b	1		1	

AOR, adjusted odd ratio; COR, crude odd ratio.

^a n 354.^b Reference categories.* Statistically significant $P < 0.05$.** Statistically significant $P < 0.001$.

months in Ethiopia^(3,10,21). This is due to the number of food groups the child had has been considered as a proxy indicator of diet quality and nutrient adequacy^(47,48), and this may play a crucial role in the linear growth of children.

Contrary to previous studies^(41,49,50), the present study showed that girls were more stunted than boys. As reported by our previous study⁽⁴²⁾, in Ethiopia, where girls are discriminated against⁽⁵¹⁾, the first choice may have been given to the needs of male children, especially as the household experienced a greater food shortage.

Our data showed high odds of being stunted among children who never received zinc supplements for diarrhoea. Though, a significant proportion of zinc-deficient children in Ethiopia⁽⁵²⁾, poor implementation of micronutrient supplementation have also been reported⁽³⁾. These emphasised the need for improvement of the micronutrient supplement programme.

Table 6. Factors associated with wasting among 6–23 months children from Siraro district, Ethiopia, 2020^a

Variables	COR [95 % CI]	P-value	AOR [95 % CI]	P-value
Family size ^b				
≥5	2.27 [1.05, 4.89]	0.035*	2.40 [1.10, 5.23]	0.027*
<5 ^c	1		1	
Received growth monitoring at least once				
No	0.40 [0.19, 0.83]	0.015*	0.43 [0.20, 0.92]	0.031*
Yes ^c	1		1	
Mother who had own income				
No	4.02 [1.38, 11.68]	0.011*	3.57 [1.21, 10.53]	0.021*
Yes ^c	1		1	

AOR, adjusted odd ratio; COR, crude odd ratio.

^a n 354.^b Family size in number.^c Reference categories.* Statistically significant $P < 0.05$.

Having more siblings was associated with higher odds of being stunted and underweight, and this is a piece of supporting evidence for a previous study from Ethiopia which stated that children whose mothers gave birth to more than four children were more likely to be stunted compared to those children who born from mothers who had only one child⁽⁵³⁾. This may be related to families with more children may face difficulty in providing proper care for child growth and development.

The age of the child is one of the risk factors for child undernutrition. This finding suggests a need for initiatives focused on improving infant and young child feeding (IYCF) practices and diet diversity, particularly those associated with complementary feeding. Failure to receive zinc supplementation for the treatment of diarrhoea increased nutritional risk as did having more siblings which suggests targets for improved implementation in the healthcare system. Additional investigation is emphasised for higher risk for female children in food-insecure areas.

This is one of a few studies in food-insecure areas of Ethiopia to investigate the prevalence and factors associated with child undernutrition at the onset of the COVID-19 pandemic and that enables us to anticipate the negative impact of the pandemic on food insecurity. Food insecurity in the study area was already alarmingly high showing 67 % of the population food insecure, out of which nearly 31 % are severely food insecure. As reported by a prior study⁽⁵⁴⁾, in addition to other factors the key drivers of food insecurity regarding COVID-19 include an increase in food prices, exacerbated food shortages resulting from travel restrictions, reduced agricultural production and physical distancing measures. Because of these, the pandemic may likely have a worsening effect on the existing food insecurity situation. As a result, it is logical to predict the situation becomes very severe shortly, particularly in the already food-insecure areas.

**Table 7.** Factors associated with underweight among 6–23 months children from Siraro district, Ethiopia, 2020^a

Variables	COR [95 % CI]	P-value	AOR [95 % CI]	P-value
Total number of births				
≥5	2.02 [1.22, 3.36]	0.006**	2.00 [1.19, 3.37]	0.047**
<5 ^b	1		1	
Child had zinc supplement for diarrhoea at least once				
No	4.40 [1.69, 11.40]	0.002**	4.35 [1.66, 11.40]	0.003**
Yes ^b	1		1	
Diet diversity of child				
<4 food groups	2.29 [1.17, 4.46]	0.015*	2.09 [1.06, 4.14]	0.036*
≥4 food groups ^b	1		1	
Family size				
≥5	1.66 [1.03, 2.69]	0.036*	1.19 [0.67, 2.10]	0.492
<5 ^b	1		1	

AOR, adjusted odd ratio; COR, crude odd ratio.

^a n 354.

^b Reference categories.

* Statistically significant $P < 0.05$.

** Statistically significant $P < 0.001$.

Furthermore, the ongoing desert locust outbreak should also be emphasised because it may further deteriorate food security⁽⁸⁾. Food may become unavailable, inaccessible, and unaffordable and malnutrition may be increased in these areas. Hence, all nutrition implementers should consider all this and work on reducing child undernutrition and maybe food shortage. Similarly, the existing social protection programmes consider promoting nutritious, safe, affordable and sustainable diets that support adequate nutrition and prevent undernutrition among infants and young children in the study area and similar settings.

The limitation of the present study was the failure to collect information on variables like the seasonality of food availability, food taboos and the COVID-19 situation in the area. There may also be recall bias in reporting different food groups consumed over the previous day referring to dietary diversity score.

Conclusion

The main factors significantly associated with child undernutrition in the study area were having more siblings, lack of zinc supplement for diarrhoea, lack of child growth monitoring, inadequate diet diversity and poor income of mothers. These factors may be particularly important targets for intervention in the study population. Our finding suggests a need for initiatives focused on improving IYCF practices and diet diversity, particularly those associated with complementary feeding. Improving zinc supplementation for the treatment of diarrhoea should be taken as one intervention strategy to mitigate child undernutrition in the study area. Limiting the number of birth and improving the growth monitoring and counselling service suggests improvement in the health care system. Our result also showed the importance of involving mothers in income-generating activities for improved nutritional status of the child.

Based on the study results, child undernutrition and food insecurity situation are alarmingly high in the study area. The situation is calling the attention of the existing social security

programme. Furthermore, all actors of nutrition should be prepared to handle the worsening effects of the mentioned factors on food insecurity in the area by contributing different aspects including humanitarian actions, and advocacy focusing on educating governments and the public on the importance of nutrition to people's survival. The nutrition community also has a role in advising the government on approaches to target the most vulnerable populations for safety nets and food aid in the context of reduced food access.

Acknowledgements

The authors acknowledge study participants and data collectors. We also acknowledge Hawassa University SPIR-DFSA project for the financial support in accomplishing this paper. The authors would also like to extend their deep thanks to all individuals who contribute to this survey.

The financial support for this study came from the SPIR-DFSA learning agenda of Hawassa University funded by the United States Agency for International Development through World Vision Ethiopia. The contents are the responsibility of the authors and do not reflect the views of the funder.

Z. T. conceived of the study, carried out the analysis and interpretations, and drafted and edited the manuscript. A. A. helped to conceive the study and analysis and drafted and edited the manuscript. F. R. and B. M. helped with the analysis, drafting and critically edited all versions of the manuscript. All authors have read and approved this manuscript to be published.

The authors declare they have no competing interests in this work.

The data will be available from the corresponding author upon reasonable request.

The proposal gained ethical clearance from the Institutional Review Board (IRB) of the College of Health Sciences of Hawassa University. Participation in the study was conducted voluntarily and with oral consent, from participant mothers. Before administering the questionnaire the study participants



discussed the research objective and requested permission to participate. The participants were reassured about the confidentiality of the data.

References

1. Onyango AW, Jean-Baptiste J, Samburu B, *et al.* (2019) Regional overview on the double burden of malnutrition and examples of program and policy responses: African region. *Ann Nutr Metab* **75**, 127–130.
2. Girma A, Woldie H, Mekonnen FA, *et al.* (2019) Undernutrition and associated factors among urban children aged 24–59 months in Northwest Ethiopia: a community-based cross-sectional study. *BMC Pediatr* **19**, 214.
3. Central Statistical Agency/CSA/Ethiopia and ICF (2016) *Ethiopia Demographic and Health Survey*. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF. Available from: <http://www.ceigram.upm.es/wp-content/uploads/2020/03/HLPE>.
4. WHO/UNICEF (2009) *WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children*. ISBN: 9789241598163.
5. UNICEF, WHO & World Bank (2020) *Levels and Trends in Child Malnutrition: Key Findings of the 2020 Edition of the Joint Child Malnutrition Estimates*, vol. **24**. Geneva: UNICEF, WHO, World Bank, pp. 1–16.
6. de Onis M & Branca F (2016) Childhood stunting: a global perspective. *Matern Child Nutr* **12**, 12–26.
7. De Onis M, Dewey KG, Borghi E, *et al.* (2013) The World Health Organization's global target for reducing childhood stunting by 2025: rationale and proposed actions. *Matern Child Nutr* **9**, 6–26.
8. FAO, IFAD, UNICEF, WFP & WHO (2018) *Food Security and Nutrition in the World the State of Building Climate Resilience for Food Security and Nutrition*. FAO, IFAD, UNICEF, WFP & WHO. Available from: www.fao.org/publications
9. Vakili R, Moghadam ZE, Khademi G, *et al.* (2015) Child mortality at different world regions: a comparison review. *Int J Pediatr* **3**, 809–816.
10. Bantamen G, Belaynew W, Dube J, *et al.* (2014) Assessment of factors associated with malnutrition among under five years age children at Machakel Woreda, Northwest Ethiopia: a case control study. *J Nutr Food Sci* **4**, 1–7.
11. UNICEF (2019) *The State of the World's Children 2019. Children, Food and Nutrition: Growing Well in a Changing World*. New York: UNICEF.
12. UNICEF, WHO, World Bank Group & UN (2019) *Levels and Trends in Child Mortality Report 2019: Estimates Developed by the UN Inter-Agency Group for Child Mortality Estimation*. New York: UNICEF.
13. Lartey A (2008) Maternal and child nutrition in Sub-Saharan Africa: challenges and interventions. *Proc Nutr Soc* **67**, 105–108.
14. FSIN (2019) *Global Report on Food Crises. Joint Analysis for Better Decisions*. Rome: FSIN.
15. Bokora C (2015) The role of safety net in ensuring food security: the case of east Harerghe zone. *J Behav Econ Finance Entrepreneurship Account Transp* **3**, 76–82.
16. FAO, IFAD, UNICEF, WFP & WHO (2019) *The State of Food Security and Nutrition in the World 2019: Safeguarding Against Economic Slowdowns and Downturns*. Rome: FAO.
17. Berhanu A (2004) The food security role of agriculture in Ethiopia. *J Agric Dev Econ* **1**, 138–153.
18. FAO, UNICEF, IFAD, WFP & WHO (2019) *The State of Food Security and Nutrition in the World. Safeguarding Against Economic Slowdowns and Downturns*, pp. 1–219. Rome: FAO, UNICEF, IFAD, WFP & WHO.
19. Burchi F, Scarlato M & D'Agostino G (2018) Addressing food insecurity in Sub-Saharan Africa: the role of cash transfers. *Poverty Public Policy* **11**, 564–589.
20. UNICEF (2020) *Supporting Children's Nutrition during the COVID-19 Pandemic*. UNICEF.
21. Fufa DA & Laloto TD (2021) Factors associated with undernutrition among children aged between 6–36 months in Semien Bench district, Ethiopia. *Heliyon* **7**, e07072.
22. Yirga AA, Mwambi HG & Ayele DG (2019) Factors affecting child malnutrition in Ethiopia. *Afr Health Sci* **19**, 1897–1909.
23. Zidouemba PR, Kinda SR & Ouedraogo IM (2020) Could COVID-19 worsen food insecurity in Burkina Faso? *Eur J Dev Res* **32**, 1379–1401.
24. CFS HLPE (2020) *Impact of COVID-19 on Food Security and Nutrition (FSN)*, pp. 1–8. Available from: <http://www.ceigram.upm.es/wp-content/uploads/2020/03/HLPE>.
25. FAO (2020) *COVID-19 and Malnutrition: Situation Analysis and Options in Africa*. FAO, pp. 1–6.
26. Omar M, Elfagi S & Nouh F (2020) COVID-19 and nutrition: review of available evidence. *Sch J Appl Med Sci* **8**, 1158–1164.
27. Hirvonen K, Abate GT, De Brauw A (2020) *Food and Nutrition Security in Addis Ababa, Ethiopia during COVID-19 Pandemic*, pp. 1–28.
28. Sirraro district health office. Health profile report (2016).
29. Tasic H, Akseer N & Gebreyesus SH (2020) Drivers of stunting reduction in Ethiopia: a country case study. *Am J Clin Nutr* **112**, 875S–893S.
30. De Onis M (2006) WHO child growth standards based on length/height, weight and age. *Acta Paediatr Int J Paediatr* **95**, 76–85.
31. FAO (2018) *Dietary Assessment: A Resource Guide to Method Selection and Application in Low Resource Settings*. Rome: FAO.
32. WHO (2010) *Indicators for Assessing Infant and Young Child Feeding Practices: Part II: Measurement*. Geneva: WHO.
33. Coates J, Swindale A & Bilinsky P (2007) *Household Food Insecurity Access Scale (HFLAS) for Measurement of Food Access: Indicator Guide*. Washington, DC: Food and Nutrition Technical Project Version 3.
34. Knuettel D, Demment M & Kaiser L (2010) Validation of the household food insecurity access scale in rural Tanzania. *Public Health Nutr* **13**, 360–367.
35. WHO (2011) *Guideline: Vitamin A Supplementation in Postpartum Women*. Geneva: World Health Organization.
36. Amare ZY, Ahmed ME & Mehari AB (2019) Determinants of nutritional status among children under age 5 in Ethiopia: further analysis of the 2016 Ethiopia Demographic and Health Survey. *Global Health* **15**, 1–11.
37. Dake SK, Solomon FB, Bobe TM, *et al.* (2019) Predictors of stunting among children 6–59 months of age in Sodo Zuria District, south Ethiopia: a community based cross-sectional study. *BMC Nutr* **5**, 1–7.
38. Seedhol AE, Mohamed ES & Mahfouz EM (2014) Determinants of stunting among preschool children, Minia, Egypt. *Int Public Health Forum* **1**, 6–9.
39. Ubeysekara NH, Jayathissa R & Wijesinghe CJ (2015) Nutritional status and associated feeding practices among children aged 6–24 months in a selected community in Sri Lanka: a cross sectional study. *Eur J Prev Med* **3**, 15.
40. Hong R, Banta JE & Betancourt JA (2006) Relationship between household wealth inequality and chronic childhood under-nutrition in Bangladesh. *Int J Equity Health* **5**, 1–10.
41. Baig-Ansari N, Rahbar MH, Bhutta ZA, *et al.* (2006) Child's gender and household food insecurity are associated with stunting among young Pakistani children residing in urban squatter settlements. *Food Nutr Bull* **27**, 114–127.
42. Tafese Z, Reta Alemayehu F, Anato A, *et al.* (2020) Child feeding practice and primary health care as major correlates of stunting and underweight among 6- to 23-month-old infants and young children in food-insecure households in Ethiopia. *Curr Dev Nutr* **4**, 1–29.
43. Geberslassie SB, Abebe SM, Melsew YA, *et al.* (2018) Prevalence of stunting and its associated factors among children 6–59 months of age in Libo-Kemekem district, Northwest Ethiopia: a community based cross sectional study. *PLoS ONE* **13**, 1–11.
44. Derso T, Tariku A, Biks GA, *et al.* (2017) Stunting, wasting and associated factors among children aged 6–24 months in Dabat health and demographic surveillance system site: a community based cross-sectional study in Ethiopia. *BMC Pediatr* **17**, 96.
45. Arimond M & Ruel MT (2004) Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. *J Nutr* **134**, 2579–2585.



46. Mallard S, Houghton L, Filteau S, *et al.* (2016) Micronutrient adequacy and dietary diversity exert positive and distinct effects on linear growth in urban Zambian infants. *J Nutr* **146**, 2093–2101.
47. Bosch AM, Baqui AH & van Ginneken JK (2008) Early-life determinants of stunted adolescent girls and boys in Matlab, Bangladesh. *J Health Nutr* **26**, 189–199.
48. Islam MM, Sanin KI, Mahfuz M, *et al.* (2018) Risk factors of stunting among children living in an urban slum of Bangladesh: findings of a prospective cohort study. *BMC Public Health* **18**, 1–13.
49. Shrimpton R, Victora CG, De Onis M, *et al.* (2001) Worldwide timing of growth faltering: implications for nutritional interventions. *Pediatrics* **107**, 1–7.
50. Bork KA & Diallo A (2017) Boys are more stunted than girls from early infancy to 3 years of age in rural Senegal. *J Nutr* **147**, 940–947.
51. Belachew T, Hadley C & Lindstrom D (2011) Gender differences in food insecurity and morbidity among adolescents in southwest Ethiopia. *Pediatrics* **127**, e398–e405.
52. Dassoni F, Abebe Z, Ricceri F, *et al.* (2014) High frequency of symptomatic zinc deficiency in infants in Northern Ethiopia. *Dermatol Res Pract*, 1–5. doi: 10.1155/2014/719701.
53. Asfaw M, Wondaferash M & Taha M (2015) Prevalence of undernutrition and associated factors among children aged between six to fifty nine months in Bule Hora district, south Ethiopia. *BMC Public Health* **15**, 41.
54. WFP (2020) *Economic and Food Security Implications of the COVID-19 Outbreak*. WFP.