



# Endoscopic lifts from $PGL_3$ to $G_2$

Wee Teck Gan and Gordan Savin

## ABSTRACT

We determine essentially completely the theta correspondence arising from the dual pair  $PGL_3 \times G_2 \subset E_6$  over a  $p$ -adic field. Our first result determines the theta lift of any non-supercuspidal representation of  $PGL_3$  and shows that the lifting respects Langlands functoriality. Our second result shows that the theta lift  $\theta(\pi)$  of a (non-self-dual) supercuspidal representation  $\pi$  of  $PGL_3$  is an irreducible generic supercuspidal representation of  $G_2$ ; we also determine  $\theta(\pi)$  explicitly when  $\pi$  has depth zero.

## 1. Introduction

Let  $k$  be a non-archimedean local field of characteristic zero and residue characteristic  $p$ . Let  $\underline{H}$  be a split adjoint linear algebraic group of type  $E_6$  over  $k$ , and let  $H = \underline{H}(k)$  be its group of  $k$ -points. There is a dual reductive pair

$$PGL_3 \times G_2 \hookrightarrow H,$$

and one can thus consider the representation correspondence induced by the restriction of the minimal representation  $\Pi$  of  $H$  to this dual pair.

For an irreducible admissible representation  $\pi$  of  $PGL_3$ , we let  $\Theta(\pi)$  denote the set of irreducible admissible representations  $\pi'$  of  $G_2$ , counted with multiplicities, such that

$$\mathrm{Hom}_{PGL_3 \times G_2}(\Pi, \pi \otimes \pi') \neq 0.$$

In [GS97], Gross and Savin have given a precise conjecture regarding the determination of the set  $\Theta(\pi)$  in terms of  $\pi$ . For the convenience of the reader, we review the conjecture briefly.

The irreducible admissible representations of  $PGL_3$  are known to be parametrized by  $L$ -parameters, which are admissible homomorphisms

$$\varphi : W_k \times SL_2(\mathbb{C}) \rightarrow SL_3(\mathbb{C}),$$

where  $W_k$  denotes the Weil group of  $k$ . Let  $A_\varphi$  be the component group of the centralizer of the image of  $\varphi$ . Then

$$A_\varphi = \begin{cases} 1 & \text{if } \varphi \text{ is reducible,} \\ \mu_3 & \text{if } \varphi \text{ is irreducible.} \end{cases}$$

If  $\pi$  is supercuspidal, then its corresponding  $L$ -parameter  $\varphi$  is irreducible, so that  $A_\varphi = \mu_3$ .

On the other hand, the irreducible admissible representations of  $G_2$  are *conjecturally* parametrized by pairs  $(\varphi', \chi')$ , where

$$\varphi' : W_k \times SL_2(\mathbb{C}) \rightarrow G_2(\mathbb{C})$$

Received 8 April 2002, accepted in final form 14 August 2002.

*2000 Mathematics Subject Classification* 22E35, 22E50 (primary), 11F70 (secondary).

*Keywords:* theta correspondence, supercuspidal lifts.

The authors are partially supported by NSF grants DMS-0202989 and DMS-0138604 respectively. This journal is © [Foundation Compositio Mathematica](http://www.compositio-mathematica.org/) 2004.

is an  $L$ -parameter for  $G_2$ , and  $\chi'$  is an irreducible character of the finite component group  $A_{\varphi'}$ . This conjectured parametrization is known for non-supercuspidal representations but not for supercuspidal representations in general.

Now, up to conjugacy, there is a natural inclusion of dual groups

$$i : SL_3(\mathbb{C}) \hookrightarrow G_2(\mathbb{C}).$$

Composing  $\varphi$  with this natural inclusion, we obtain an  $L$ -parameter  $\varphi' = i \circ \varphi$  for the group  $G_2$ . The inclusion  $i$  also induces an injection  $i_* : A_\varphi \rightarrow A_{\varphi'}$  [GS97, Proposition 2.7], which identifies  $A_\varphi$  as a subgroup of index 1 or 2.

Consider the restriction of the minimal representation  $\Pi$  of  $H$  to the dual pair  $PGL_3 \times G_2$ . The maximal  $\pi$ -isotypic quotient

$$\Pi / \bigcap_{\phi \in \text{Hom}_{PGL_3}(\Pi, \pi)} \text{Ker}(\phi)$$

of  $\Pi$  can be expressed as  $\pi \otimes \theta(\pi)$ , for some smooth representation  $\theta(\pi)$  of  $G_2$  [MVW87, Lemme III.4] and, by definition, there is a  $PGL_3 \times G_2$ -equivariant surjection  $\Pi \rightarrow \pi \otimes \theta(\pi)$ . Then  $\Theta(\pi)$  is simply the set of irreducible quotients of  $\theta(\pi)$ , counted with multiplicities. The following is the conjecture from [GS97].

CONJECTURE. The correspondence is functorial with respect to the inclusion of dual groups. More precisely,  $\Theta(\pi)$  is the set of  $\pi'$  whose parameter  $(\varphi', \chi')$  satisfies

$$\begin{aligned} \varphi' &= i \circ \varphi_\pi, \\ \chi' \circ i_* &= 1. \end{aligned}$$

This conjecture was verified in [MS97] for  $\pi$  a tempered spherical representation, and in [GS97] for  $\pi$  a generalized principal series representation. The first main result of this paper is the complete determination of the set  $\Theta(\pi)$  when  $\pi$  is non-supercuspidal, given in Theorems 11, 14 and 15. As a result, we have the following.

THEOREM 1. *The conjecture is true for non-supercuspidal  $\pi$ .*

We highlight here an important special case, given by our Theorem 14. If  $\pi = St$  is the Steinberg representation of  $PGL_3$ , then its  $L$ -parameter  $\varphi$  is trivial on  $W_k$  and  $\varphi|_{SL_2(\mathbb{C})}$  is the homomorphism given by the adjoint representation of  $SL_2(\mathbb{C})$ . Thus,  $A_\varphi \cong \mu_3$  and  $A_{\varphi'} \cong S_3$ . The symmetric group  $S_3$  has three irreducible characters, 1,  $\rho$  and  $\epsilon$ , where  $\rho$  is a two-dimensional character and  $\epsilon$  is the sign character. We show that

$$\Theta(St) = \{\pi(1)', \pi_{sc}[1]\},$$

where  $\pi(1)'$  is a generic discrete series representation of  $G_2$  corresponding to the parameter  $(\varphi', 1)$  and  $\pi_{sc}[1]$  is a non-generic supercuspidal representation corresponding to the parameter  $(\varphi', \epsilon)$ .

Since the local Langlands conjecture for supercuspidal representations of  $G_2$  is not known, the conjecture of Gross–Savin cannot be verified literally for supercuspidal representations of  $PGL_3$ . However, if  $\pi$  is supercuspidal with  $L$ -parameter  $\varphi$ , then

$$A_{i \circ \varphi} = \begin{cases} \mu_3 & \text{if } \pi \text{ is not self-dual,} \\ S_3 & \text{if } \pi \text{ is self-dual.} \end{cases}$$

In particular, the conjecture predicts that

$$\#\Theta(\pi) = \begin{cases} 1 & \text{if } \pi \text{ is not self-dual,} \\ 2 & \text{if } \pi \text{ is self-dual.} \end{cases}$$

Our second main result verifies the above prediction when  $\pi$  is not self-dual.

**THEOREM 2.** *Let  $\pi$  be a supercuspidal representation of  $PGL_3$  which is not self-dual. Then  $\theta(\pi)$  is an irreducible generic supercuspidal representation of  $G_2$ .*

We remind the reader that if  $p \neq 2$ , there are no self-dual supercuspidal representations of  $PGL_3$ , so the above result holds for all supercuspidal representations.

The two theorems above essentially give complete information about the theta lifting from  $PGL_3$  to  $G_2$  (when  $p \neq 2$ ), except for the determination of the irreducible supercuspidal representation  $\theta(\pi)$  when  $\pi$  is supercuspidal. It may be possible to give a characterization of  $\theta(\pi)$  in terms of  $\pi$  without referring to the local Langlands conjecture. In the final section, we do this for  $\pi$  of depth zero, using the construction of supercuspidal representations by induction from open compact subgroups.

Note that  $PGL_3$  is an endoscopic group of  $G_2$  and the above results show that the theta correspondence for  $PGL_3 \times G_2$  can be used to construct elements of the corresponding endoscopic  $L$ -packets of  $G_2$ . To obtain every element of such a packet, we would need to consider the dual pair correspondence arising from  $PD^\times \times G_2$ , where  $D$  is a degree 3 division algebra. This dual pair is an inner twist of the one considered in this paper and was studied in [Sav99]. These local results are necessary preliminaries for the construction of global endoscopic  $L$ -packets using global theta liftings from  $PGL_3$  and  $PD^\times$ . We plan to pursue this global question in a future paper.

### 2. Representations of $l$ -groups

In this section, we establish some notations and discuss some basic facts on the representation theory of  $p$ -adic groups that are required later.

Let  $G$  be the  $k$ -points of a connected reductive algebraic group  $\underline{G}$  over  $k$ . Then recall that  $G$  is an  $l$ -group in the terminology of [BZ76]. Let  $\text{Alg}(G)$  be the category of smooth representations of  $G$ . For  $\pi \in \text{Alg}(G)$ , we let  $\pi^*$  denote the full linear dual of  $\pi$ ; it is a representation of  $G$  but is not usually smooth. Its subspace of smooth vectors is the contragredient representation  $\pi^\vee$ .

Recall that if  $P = MN$  is a parabolic subgroup of  $G$ , then we have an exact functor (normalized parabolic induction)

$$I_P : \text{Alg}(M) \longrightarrow \text{Alg}(G).$$

We also have the normalized Jacquet functor

$$R_P : \text{Alg}(G) \longrightarrow \text{Alg}(M).$$

Let  $\bar{P}$  be the opposite parabolic subgroup. Then we have the following adjointness properties:

$$\begin{aligned} \text{Hom}_G(\pi, I_P(\sigma)) &= \text{Hom}_M(R_P(\pi), \sigma), \\ \text{Hom}_G(I_P(\sigma), \pi) &= \text{Hom}_M(\sigma, R_{\bar{P}}(\pi)). \end{aligned}$$

A complete proof of the second property has been given in [Bus01].

Sometimes, we may wish to talk about unnormalized induction and unnormalized Jacquet functor. We use  $\text{Ind}$  and  $\text{ind}$  to denote unnormalized smooth and compact induction, respectively, and we write  $\pi \mapsto \pi_N$  (coinvariants) for unnormalized Jacquet functor. More generally, if  $\psi$  is a (unitary) character of  $N$ ,  $\pi_{N,\psi}$  denotes the maximal quotient of  $\pi$  on which  $N$  acts by  $\psi$ .

An irreducible representation  $\pi$  is said to belong to the discrete series or is square-integrable if  $\pi$  has unitary central character and its matrix coefficients are square-integrable modulo center. An example is the so-called Steinberg representation. A representation  $\pi \in \text{Alg}(G)$  is called supercuspidal if  $R_P(\pi) = 0$  for any proper parabolic subgroup  $P$ . If  $\pi$  is an irreducible unitarizable supercuspidal representation, then it is square-integrable. For any  $\pi \in \text{Alg}(G)$ , we have a decomposition (see [BZ76])

$$\pi = \pi_c \oplus \pi_i$$

where  $\pi_c$  is the supercuspidal part of  $\pi$  and  $\pi_i$  has no non-zero supercuspidal subquotient.

Let us also recall the notion of a ‘generic representation’ of  $G$ . For simplicity, assume that  $\underline{G}$  is split and has connected center. Let  $B = T \cdot U$  be a Borel subgroup with unipotent radical  $U$  and maximal split torus  $T$ . A character  $\psi$  of  $U$  is said to be in general position if its stabilizer in  $T$  is equal to the center of  $G$ . The group  $T$  acts transitively on the set of such characters of  $U$  and an irreducible representation  $\pi$  of  $G$  is said to be generic if  $\pi_{U,\psi} \neq 0$  for some (and hence any)  $\psi$  in general position. By Frobenius reciprocity, this is equivalent to saying that  $\pi$  admits a non-zero  $G$ -equivariant map into  $\text{Ind}_U^G \psi$ . In this case,  $\dim(\pi_{U,\psi}) = 1$ .

Finally, on several occasions, we use the following simple observation. Let

$$0 \longrightarrow V_1 \longrightarrow V_2 \longrightarrow V_3 \longrightarrow 0$$

be an exact sequence of  $G$ -modules. If an element of the Bernstein center of  $G$  acts on  $V_1$  and  $V_3$  by different eigenvalues, then  $V_1$  is a quotient of  $V_2$ , so that  $V_2 \cong V_1 \oplus V_3$ .

### 3. Structure of maximal parabolics

Let  $\tilde{Q}_1$  and  $\tilde{Q}_2$  be the two non-conjugate maximal parabolic subgroups of  $GL_3(k) = GL(k^3)$  stabilizing a one-dimensional subspace  $W_1$  and a two-dimensional subspace  $W_2$  of  $k^3$ , respectively. We fix  $W_1 \subset W_2$ . Their Levi factors are  $GL(W_1) \times GL(W_1^\perp)$  and  $GL(W_2) \times GL(W_2^\perp)$ , respectively, where  $W_1^\perp$  and  $W_2^\perp$  are annihilators of  $W_1$  and  $W_2$  in  $(k^3)^*$ . The corresponding maximal parabolic subgroups in  $PGL_3$  are denoted by  $Q_1 = L_1U_1$  and  $Q_2 = L_2U_2$ . We have isomorphisms

$$\begin{aligned} L_1 &\cong GL(W_1^\perp) \\ L_2 &\cong GL(W_2). \end{aligned}$$

Using these identifications, the modular characters of  $L_1$  and  $L_2$  are

$$\rho_1(g) = |\det g|^{1/2} \quad \text{and} \quad \rho_2(g) = |\det g|^{1/2}.$$

Further,  $Q_0 = Q_1 \cap Q_2$  is a Borel subgroup of  $PGL_3$ .

The maximal parabolic subgroups of  $G_2$  can be defined as the stabilizers of non-trivial nil subalgebras of the octonion  $k$ -algebra  $\mathbb{O}$ . A nil subalgebra of  $\mathbb{O}$  is a subspace consisting of trace zero elements with trivial multiplication (i.e. the product of any two elements is 0). The possible dimensions are 1 and 2. Fix a pair of nil-subalgebras  $V_1 \subset V_2$ . Then the stabilizers  $P_1 = M_1N_1$  and  $P_2 = M_2N_2$  of  $V_1$  and  $V_2$ , respectively, are two non-conjugate maximal parabolic subgroups of  $G_2$ , with  $P_0 = P_1 \cap P_2$  a Borel subgroup. Let

$$V_3 = \{x \in \mathbb{O} \mid \bar{x} = -x \text{ and } x \cdot V_1 = 0\}.$$

We have isomorphisms

$$\begin{aligned} M_1 &\cong GL(V_3/V_1) \\ M_2 &\cong GL(V_2). \end{aligned}$$

The action of the Levi factor of  $P_1$  on  $V_1$  is given by  $\det$ , and the modular characters are

$$\rho'_1(g) = |\det(g)|^{5/2} \quad \text{and} \quad \rho'_2(g) = |\det(g)|^{3/2}.$$

### 4. Representations of $PGL_3$

In this section, we describe the irreducible smooth representations of  $PGL_3$ . It will be convenient to describe the representations in terms of discrete series representations of Levi factors. First of all, we have the discrete series representations: the supercuspidal representations, the Steinberg representation  $St$  and its twists  $St_{\chi_E}$  and  $St_{\chi_E^2} \cong St_{\chi_E}^\vee$  by cubic characters  $\chi_E$ , each corresponding to a Galois cubic extension  $E$  of the base field  $k$ .

Next, let  $\delta$  denote a (unitarizable) discrete series representation of  $GL_2$ . Then  $\delta$  is either a supercuspidal representation or a twist  $st_\mu$  of the Steinberg representation  $st$  by a unitary character  $\mu$ . Let  $\delta_s$  be the twist of  $\delta$  by  $|\det|^s$  and write  $I_{Q_k}(\delta, s)$  for  $I_{Q_k}(\delta_s)$  ( $k = 1$  or  $2$ ). For  $s \geq 0$ , let

$$J_k(\delta, s) = \text{the unique irreducible quotient of } I_{Q_k}(\delta, s).$$

Here we have identified the Levi factors with  $GL_2$  as in the previous section. The following will be very useful for our calculations (here  $i \neq j$ ):

$$\begin{aligned} J_i(\delta, s) &\cong J_j(\delta, s)^\vee \\ J_i(\delta, s) &\subseteq I_{Q_j}(\delta^\vee, -s). \end{aligned}$$

Finally, we have the representations  $J(\chi_1, \chi_2, \chi_3)$  which are the unique irreducible quotient of representations obtained by (normalized) parabolic induction from the Borel subgroup  $Q_0 = Q_1 \cap Q_2$ . Here the  $\chi_i$  are three characters such that  $\chi_1 \chi_2 \chi_3 = 1$ . Furthermore, if we write each character  $\chi_i$  as  $\chi_i = \mu_i \cdot |\cdot|^{s_i}$ , where  $\mu_i$  is unitary and  $s_i \in \mathbb{R}$ , then

$$\begin{aligned} s_1 &\geq s_2 \geq s_3, \\ s_1 + s_2 + s_3 &= 0. \end{aligned}$$

In terms of intermediate induction, the representation  $J(\chi_1, \chi_2, \chi_3)$  can be realized as a submodule

$$J(\chi_1, \chi_2, \chi_3) \subset I_{Q_2}(\pi(\chi_3, \chi_2)).$$

Since  $J(\chi_1, \chi_2, \chi_3)^\vee \cong J(\chi_3^{-1}, \chi_2^{-1}, \chi_1^{-1})$ , we also have

$$J(\chi_1, \chi_2, \chi_3)^\vee \subset I_{Q_2}(\pi(\chi_1^{-1}, \chi_2^{-1})).$$

The following proposition is very useful for later purposes and can be read off from the above discussion.

**PROPOSITION 3.** *Let  $\pi$  be an irreducible smooth representation of  $PGL_3$  which is not square integrable. Then either  $\pi$  or its contragredient  $\pi^\vee$  is the unique irreducible submodule of an induced representation  $I_{Q_2}(\tau^\vee)$ , where  $\tau$  is an irreducible representation of  $L_2$  described as follows.*

- i) If  $\pi = J_i(\delta, s)$ , then  $\tau = \delta_s$ .
- ii) If  $\pi = J(\chi_1, \chi_2, \chi_3)$ , we may assume that  $\chi_2 = \mu_2 |\cdot|^{s_2}$  with  $s_2 \geq 0$  (by possibly replacing  $\pi$  by  $\pi^\vee$ ). There are now two cases as follows.
  - a) If  $\chi_1/\chi_2 \neq |\cdot|$ , then  $\tau = \pi(\chi_1, \chi_2)$ .
  - b) Otherwise, suppose that  $\chi_1 = \mu |\cdot|^{s+1}$  and  $\chi_2 = \mu |\cdot|^s$  for some unitary character  $\mu$ . Then  $\tau = \mu(\det)_{s+1/2}$ .

In each case, the central character  $\chi_\tau$  of  $\tau$  satisfies  $|\chi_\tau| = |\cdot|^t$  with  $t \geq 0$ . Further, whenever  $t = 0$ ,  $\tau$  is unitarizable and  $I_{Q_2}(\tau^\vee)$  is irreducible.

### 5. Minimal representation

There is a notion of a ‘minimal representation’ of a  $p$ -adic reductive group; we refer the reader to [Sav96] for the definition. For a split simply-laced group  $H$  of type  $A_{2n-1}$ ,  $D$  or  $E$ , it is known that a unitarizable minimal representation exists and is unique up to twisting by (one-dimensional) characters of  $H$ . For the purpose of this paper, we need to specify exactly what we mean by the minimal representation of  $H$ .

Since the algebraic group  $\underline{H}$  is split, we may assume that  $\underline{H}$  is defined over  $\mathbb{Z}$ . If  $A$  is the ring of integers of  $k$ , then  $K = \underline{H}(A)$  is a maximal compact subgroup of  $H$ . Recall that an irreducible representation of  $H$  is said to be unramified if it has a non-zero  $K$ -fixed vector. The set of isomorphism

classes of unramified representations is in natural bijection with semisimple conjugacy classes in the Langlands dual group  $H^\vee(\mathbb{C})$ . Given an unramified representation, the corresponding semisimple class in  $H^\vee(\mathbb{C})$  is called its Satake parameter.

The (unitarizable) minimal representation  $\Pi$  of  $H$  is an unramified representation and its Satake parameter is described as follows. Let  $\iota : SL_2(\mathbb{C}) \rightarrow H^\vee(\mathbb{C})$  be a homomorphism corresponding to the subregular unipotent orbit of  $H^\vee(\mathbb{C})$  by the Jacobson–Morozov theorem. Then the Satake parameter of  $\Pi$  is the conjugacy class of

$$s_\Pi = \iota \left( \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix} \right),$$

where  $q$  is the order of the residue field of  $k$ . The fact that this unramified representation is minimal was shown in [Sav94].

Henceforth,  $H$  denotes the split adjoint group of type  $E_6$ . We need various important properties of the minimal representation and these are reviewed in the following two sections. We conclude this section by observing that any automorphism of  $H$  fixes the isomorphism class of  $\Pi$  since it preserves the corresponding Satake parameter. Furthermore, there is an automorphism of  $H$  which stabilizes the subgroup  $PGL_3 \times G_2$ , acts trivially on  $G_2$  and maps to the non-trivial outer automorphism of  $PGL_3$ . As a consequence of this, we see that for any irreducible representation  $\pi$  of  $PGL_3$ ,  $\Theta(\pi) = \Theta(\pi^\vee)$ . Hence, for the purpose of determining  $\Theta(\pi)$ , we can restrict ourselves to representations  $\pi$  satisfying the conditions of Proposition 3.

### 6. Jacquet modules for $PGL_3$

In this section, we describe the  $L_2 \times G_2$ -module  $R_{Q_2}(\Pi)$ . The result is given by [MS97, Theorem 4.3]. To state it, we need some additional notations. There exists a maximal parabolic  $\Omega_2 = \mathfrak{L}_2 \mathfrak{U}_2$  in  $H$  whose Levi factor  $\mathfrak{L}_2$  is of type  $D_5$ , and such that

$$\begin{aligned} (PGL_3 \times G_2) \cap \mathfrak{L}_2 &= L_2 \times G_2 \\ PGL_3 \cap \mathfrak{U}_2 &= U_2. \end{aligned}$$

Let  $B$  be the Borel subgroup of  $GL(W_2)$  stabilizing the line  $W_1$ . The result is the following.

PROPOSITION 4. *The  $GL_2 \times G_2$ -module  $R_{Q_2}(\Pi)$  has a filtration*

$$0 = \Pi_0 \subset \Pi_1 \subset \Pi_2 \subset \Pi_3 = \Pi_{U_2}$$

such that

$$\begin{aligned} \Pi_1/\Pi_0 &\cong |\det|^{3/2} \otimes \text{Ind}_{GL_2 \times P_2}^{GL_2 \times G_2}(C_c^\infty(GL_2)) \\ \Pi_2/\Pi_1 &\cong |\det|^{3/2} \otimes \text{Ind}_{B \times P_1}^{GL_2 \times G_2}(C_c^\infty(GL_1)) \\ \Pi_3/\Pi_2 &= |\det|^{-1/2} \cdot \Pi_{U_2} \cong |\det|^{1/2} \otimes \Pi(\mathfrak{L}_2) + |\det|^{3/2} \otimes 1. \end{aligned}$$

Here  $\det$  is the determinant map on  $GL(W_2)$ . Moreover,  $C_c^\infty(GL_2)$  is the regular representation of  $GL(W_2) \times GL(V_2)$ , and  $C_c^\infty(GL_1)$  is the regular representation of  $GL(W_1) \times GL(V_1)$ . Finally,  $\Pi(\mathfrak{L}_2)$  is the minimal representation of  $\mathfrak{L}_2$ . The center of  $\mathfrak{L}_2$ , which coincides with the center of  $GL(W_2)$ , acts trivially on  $\Pi(\mathfrak{L}_2)$ .

Analogously, there is a maximal parabolic subgroup  $\Omega_1 = \mathfrak{L}_1 \mathfrak{U}_1$  of  $H$  whose Levi factor  $\mathfrak{L}_1$  has type  $D_5$  and such that

$$\begin{aligned} (PGL_3 \times G_2) \cap \mathfrak{L}_1 &= L_1 \times G_2 \\ PGL_3 \cap \mathfrak{U}_1 &= U_1. \end{aligned}$$

The two parabolic subgroups  $\mathfrak{Q}_1$  and  $\mathfrak{Q}_2$  are not conjugate in  $H$ , but are conjugate under a non-inner automorphism of  $H$ . We have an obvious analog of Proposition 4 for  $R_{Q_1}(\Pi)$ .

Now  $\mathfrak{Q}_0 = \mathfrak{Q}_1 \cap \mathfrak{Q}_2$  is a parabolic subgroup of  $H$  whose Levi factor  $\mathfrak{L}_0$  is of type  $D_4$ . Its intersection with  $PGL_3$  is equal to the Borel subgroup  $Q_0 = Q_1 \cap Q_2$  and the center of  $\mathfrak{L}_0$  is precisely the Levi factor  $L_0$  of  $Q_0$ . We have the following proposition.

PROPOSITION 5. *Let  $\Pi(D_4)$  be the minimal representation of  $\mathfrak{L}_1 \cap \mathfrak{L}_2$  with the center  $L_0$  acting trivially. Then the  $L_0 \times G_2$ -module  $R_{Q_0}(\Pi)$  has  $\rho_0 \otimes \Pi(D_4)$  as a quotient. Here,  $\rho_0$  is the modulus character of  $Q_0$ .*

*Proof.* The exponents of  $\Pi$  are known (see [Sav94]). In particular, it can be shown that  $\rho_0^2 \otimes \Pi(D_4)$  is a summand of  $\Pi_{\mathfrak{U}_0}$ , regarded as a  $L_0 \times G_2$ -module. Since  $\rho_0^{-1} \otimes \Pi_{\mathfrak{U}_0}$  is a quotient of  $R_{Q_0}(\Pi)$ , the proposition easily follows.  $\square$

### 7. Jacquet modules for $G_2$

In this section, we describe the structure of the  $PGL_3 \times M_2$ -module  $R_{P_2}(\Pi)$ . The result is given in [MS97, Theorem 7.6]. As in the previous section, there is a maximal parabolic subgroup  $\mathfrak{P}_2 = \mathfrak{M}_2\mathfrak{N}_2$  of  $H$  whose Levi factor  $\mathfrak{M}_2$  is of type  $A_5$  and such that

$$(PGL_3 \times G_2) \cap \mathfrak{M}_2 = PGL_3 \times M_2$$

$$G_2 \cap \mathfrak{N}_2 = N_2.$$

Let  $B'$  be the Borel subgroup of  $M_2 = GL(V_2)$  stabilizing the line  $V_1$ .

PROPOSITION 6. *The  $PGL_3 \times GL_2$ -module  $R_{P_2}(\Pi)$  has a filtration*

$$0 = \Pi_0 \subset \Pi_1 \subset \Pi_2 \subset \Pi_3 = R_{P_2}(\Pi)$$

such that

$$\Pi_1/\Pi_0 \cong |\det|^{1/2} \otimes \text{Ind}_{Q_1 \times GL_2}^{PGL_3 \times GL_2} C_c^\infty(GL_2) \bigoplus |\det|^{1/2} \otimes \text{Ind}_{Q_2 \times GL_2}^{PGL_3 \times GL_2} C_c^\infty(GL_2);$$

$$\Pi_2/\Pi_1 \cong |\det|^{1/2} \otimes \text{Ind}_{Q_0 \times B'}^{PGL_3 \times GL_2} C_c^\infty(GL_1),$$

$$\Pi_3/\Pi_2 \cong \Pi(\mathfrak{M}_2) \bigoplus 1 \otimes |\det|^{1/2}.$$

Here  $\det$  is the usual determinant on  $GL(V_2)$ . Moreover,  $C_c^\infty(GL_2)$  is the regular representation of

$$GL(W_1^\perp) \times GL(V_2) \quad \text{and} \quad GL(W_2) \times GL(V_2),$$

respectively, and  $C_c^\infty(GL_1)$  is the regular representation of

$$GL(W_1 \otimes W_2^\perp) \times GL(V_2/V_1).$$

Finally,  $\Pi(\mathfrak{M}_2)$  is the minimal representation of  $\mathfrak{M}_2$ . The center of  $\mathfrak{M}_2$ , which coincides with the center of  $GL(V_2)$ , acts trivially on  $\Pi(\mathfrak{M}_2) = \Pi(PGL_6)$ .

### 8. Non-square-integrable representations

In this section, we determine the set  $\Theta(\pi)$  for non-square-integrable  $\pi$  and verify the conjecture of Gross–Savin for such  $\pi$ . As we remarked before, there is no loss of generality in assuming that  $\pi$  satisfies the conditions of Proposition 3. In particular, we have an irreducible representation  $\tau$  of  $L_2$  according to that proposition and  $\pi \hookrightarrow I_{Q_2}(\tau^\vee)$ . We define the non-negative number  $t$  by  $|\chi_\tau| = |\cdot|^t$ .

PROPOSITION 7. *As representations of  $G_2$ ,*

$$\text{Hom}_{PGL_3}(\Pi, I_{Q_2}(\tau^\vee)) \cong I_{P_2}(\tau)^*.$$

*Proof.* By Frobenius reciprocity, we have

$$\text{Hom}_{PGL_3}(\Pi, I_{Q_2}(\tau^\vee)) = \text{Hom}_{L_2}(R_{Q_2}(\Pi), \tau^\vee).$$

The structure of  $R_{Q_2}(\Pi)$  as a representation of  $L_2 \times G_2$  has been determined in Proposition 4. Using the notation of that proposition, we now note the following lemma.

LEMMA 8.  $\text{Hom}_{L_2}(R_{Q_2}(\Pi), \tau^\vee) = \text{Hom}_{L_2}(\Pi_1, \tau^\vee)$ .

*Proof.* The central characters of  $L_2$  on  $\Pi/\Pi_2$  are  $|\cdot|$  and  $|\cdot|^3$  and these are different from  $|\chi_{\tau^\vee}| = |\cdot|^{-t}$ . Thus, we have

$$\text{Hom}_{L_2}(R_{Q_2}(\Pi), \tau^\vee) = \text{Hom}_{L_2}(\Pi_2, \tau^\vee).$$

Next, for any  $\tau^\vee$  as in Proposition 3, we can find an element  $z$  in the Bernstein center of  $GL_2$  such that  $z(\tau^\vee) = 1$  and  $z$  vanishes for all representations of  $GL_2$  with inducing parameters of type  $(\chi, |\cdot|^2)$  (cf. [MS97, pp. 104–105]). Such a  $z$  vanishes on  $\Pi_2/\Pi_1$  and thus we have

$$\text{Hom}(R_{Q_2}(\Pi), \tau^\vee) = \text{Hom}(\Pi_1, \tau^\vee). \quad \square$$

We now return to the proof of the proposition. By the lemma, we need to identify the  $G_2$ -module

$$\text{Hom}_{L_2}(\text{Ind}_{L_2 \times P_2}^{L_2 \times G_2}(C_c^\infty(GL_2)), \tau_{-3/2}^\vee).$$

Now if we express the maximal  $\tau_{-3/2}^\vee$ -isotypic quotient of  $\text{Ind}_{L_2 \times P_2}^{L_2 \times G_2}(C_c^\infty(GL_2))$  as  $V \otimes \tau_{-3/2}^\vee$  for some smooth  $G_2$ -module  $V$ , then

$$\text{Hom}_{L_2}(\text{Ind}_{L_2 \times P_2}^{L_2 \times G_2}(C_c^\infty(GL_2)), \tau_{-3/2}^\vee) \cong V^*.$$

It remains to prove that  $V \cong I_{P_2}(\tau)$ .

By [MVW87, Lemme, p. 59], the maximal  $\tau_{-3/2}^\vee$ -isotypic quotient of  $C_c^\infty(GL_2)$  is isomorphic to  $\tau_{3/2} \otimes \tau_{-3/2}^\vee$ . Hence, that of  $\text{Ind}_{P_2 \times L_2}^{G_2 \times L_2}(C_c^\infty(GL_2))$  is  $I_{P_2}(\tau) \otimes \tau_{-3/2}^\vee$ . In other words,  $V \cong I_{P_2}(\tau)$  and the proposition is proved.  $\square$

COROLLARY 9. Assume that  $\pi \hookrightarrow I_{Q_2}(\tau^\vee)$  as above. Then:

- i)  $\theta(\pi)$  is a quotient of  $I_{P_2}(\tau)$ , in particular it has finite length and

$$\Theta(\pi) \subset \{\text{the irreducible quotients of } I_{P_2}(\tau)\};$$

- ii) for any irreducible quotient  $\sigma$  of  $I_{P_2}(\tau)$ , there is a non-zero  $PGL_3 \times G_2$ -equivariant map  $\Pi \rightarrow I_{Q_2}(\tau^\vee) \otimes \sigma$ .

*Proof.* i) Observe that

$$\theta(\pi)^* = \text{Hom}_{PGL_3}(\Pi, \pi) \subset \text{Hom}_{PGL_3}(\Pi, I_{Q_2}(\tau^\vee)) = I_{P_2}(\tau)^*$$

so that  $\theta(\pi)^\vee \subset I_{P_2}(\tau)^\vee$ . Hence  $\theta(\pi)^\vee$  is admissible, and thus so is  $\theta(\pi)$ . The result follows.

- ii) The isomorphism of the proposition implies that there is a non-zero  $G_2$ -equivariant map

$$\sigma^\vee \hookrightarrow \text{Hom}_{PGL_3}(\Pi, I_{Q_2}(\tau^\vee)).$$

Since

$$\text{Hom}_{G_2}(\sigma^\vee, \text{Hom}_{PGL_3}(\Pi, I_{Q_2}(\tau^\vee))) \cong \text{Hom}_{G_2 \times PGL_3}(\Pi, \sigma \otimes I_{Q_2}(\tau^\vee)),$$

the result follows.  $\square$

For  $\sigma$  an irreducible smooth representation of  $G_2$ , we may define  $\theta'(\sigma)$  and  $\Theta'(\sigma)$  as in the case of  $PGL_3$ . We have the following.



PROPOSITION 10. *Let  $\tau$  as in Proposition 3 and assume that  $t > 0$ . Then  $I_{P_2}(\tau)$  has a unique irreducible quotient  $\sigma$ . Moreover, ignoring multiplicities, we have  $\Theta'(\sigma) \subset \{\pi, \pi^\vee\}$ .*

*Proof.* It is not difficult to check that  $I_{P_2}(\tau)$  is a quotient of an induced representation of Langlands type (i.e. a representation induced from a quasi-tempered representation of a Levi subgroup, whose central character is in the relevant positive chamber). In particular,  $I_{P_2}(\tau)$  has a unique irreducible quotient  $\sigma$ . Further, this unique irreducible quotient  $\sigma$  is the image of a standard intertwining map  $I_{P_2}(\tau) \rightarrow I_{P_2}(\tau^\vee)$ . From this, we deduce that  $\sigma$  is self-contragredient and is the unique irreducible submodule of  $I_{P_2}(\tau^\vee)$ .

Now, for any representation  $\delta$  of  $PGL_3$ , we have

$$\text{Hom}_{PGL_3 \times G_2}(\Pi, \delta \otimes \sigma) \subset \text{Hom}_{PGL_3 \times G_2}(\Pi, \delta \otimes I_{P_2}(\tau^\vee)).$$

Using Proposition 6, it is not difficult to estimate the latter space. In particular, we see that the latter space is zero unless  $\delta = \pi$  or  $\pi^\vee$ , which gives the desired upper bound on  $\Theta'(\sigma)$ . We remark that we do not have an analog of Proposition 7 because the analog of Lemma 8 is not true in general. □

The following is the main result of this section.

THEOREM 11. *i) Let  $\pi$  be an irreducible representation of  $PGL_3$  which is not square-integrable. Suppose that  $\pi \hookrightarrow I_{Q_2}(\tau^\vee)$  with  $\tau$  as in Proposition 3. Then*

$$\Theta(\pi) = \Theta(\pi^\vee) = \{\text{the irreducible quotients of } I_{P_2}(\tau)\}.$$

*Further, the correspondence is functorial for  $\pi$ .*

*ii) Suppose that  $t > 0$ . If  $\sigma$  is the unique irreducible quotient of  $I_{P_2}(\tau)$ , then*

$$\Theta'(\sigma) = \{\pi, \pi^\vee\} \quad \text{if } \pi \neq \pi^\vee,$$

*and*

$$\Theta'(\sigma) = \{\pi\} \quad \text{if } \pi = \pi^\vee.$$

*Proof.* i) After Corollary 9, part i, we need to show that if  $\sigma$  is any irreducible quotient of  $I_{P_2}(\tau)$ , then  $\pi \otimes \sigma$  is a quotient of  $\Pi$ . By Corollary 9, part ii, there is a non-zero map  $f : \Pi \rightarrow I_{Q_2}(\tau^\vee) \otimes \sigma$ . If  $I_{Q_2}(\tau^\vee)$  is irreducible and, hence, equal to  $\pi$ , then we are done. If  $I_{Q_2}(\tau^\vee)$  is reducible, then  $t > 0$ . However, by Proposition 10, any irreducible quotient of the image of  $f$  must be  $\pi \otimes \sigma$  or  $\pi^\vee \otimes \sigma$ , i.e.  $\sigma \in \Theta(\pi) = \Theta(\pi^\vee)$ . Thus the first statement of part i is proved.

To show that the correspondence is functorial for  $\pi$ , we need to show that the  $L$ -parameters match up. This is straightforward except in case ii of Proposition 3 when  $s_2 = 0$ , so that the representation  $\sigma \in \Theta(\pi)$  is a Langlands quotient of a representation induced from  $P_1$ . We leave the verification to the reader.

ii) This follows from part i and Proposition 10. □

For the purpose of global applications, it is useful to note the following corollary.

COROLLARY 12. *Let  $\pi(s)$  be the unramified representation of  $PGL_3$  attached to the semisimple conjugacy class of  $s \in SL_3(\mathbb{C})$ . Similarly, let  $\pi(i(s))$  be the unramified representation of  $G_2$  attached to the conjugacy class of  $i(s) \in G_2(\mathbb{C})$ . Then  $\Theta(\pi(s)) = \{\pi(i(s))\}$ .*

Finally, observe that  $\Theta(\pi)$  is a singleton set if and only if  $I_{P_2}(\tau)$  has a unique irreducible quotient. If  $t > 0$ , we have observed that this is always the case. If  $t = 0$ , then we need to know whether  $I_{P_2}(\tau)$  is irreducible. This reducibility question has been studied in [Key82, Mui97, Sha91]. Together with their results, Theorem 11 implies the following.

COROLLARY 13. Suppose that  $\pi \hookrightarrow I_{Q_2}(\tau^\vee)$  with  $\tau$  as in Proposition 3. Then  $\Theta(\pi)$  is a singleton set except in the following cases where it has two elements:

- i)  $\tau$  is a self-dual unitarizable supercuspidal representation whose central character is non-trivial;
- ii)  $\tau = \pi(\chi_1, \chi_2)$  with  $\chi_1 \neq \chi_2$  characters of order 2.

### 9. Discrete series

In this section, we determine the set  $\Theta(\pi)$  for  $\pi$  a non-cuspidal discrete series representation. Recall that  $\pi$  is either the Steinberg representation  $St$  or its twist  $St_\chi$  by a cubic character  $\chi$ . The pair  $\{\chi, \chi^{-1}\}$  of cubic characters determine a Galois cubic extension  $E$  of  $F$  by local class field theory. The field  $E$  then determines a square-integrable representation  $\pi(E)$  of  $G_2$  (cf. [Mui97, Proposition 4.2(i)] where this representation was denoted by  $\pi(\chi) = \pi(\chi^{-1})$ ).

THEOREM 14. Let  $\chi$  be a cubic character with corresponding cyclic cubic extension  $E$ . Then

$$\Theta(St_\chi) = \Theta(St_{\chi^2}) = \{\pi(E)\}.$$

*Proof.* Let  $st$  denote the Steinberg representation of  $L_2$  and let  $st_\chi$  be the twisted representation  $st \otimes \chi(\det)$ . Then  $St_\chi$  is a submodule of  $I_{Q_2}(st_\chi, \frac{1}{2})$ . Thus,

$$\theta(St_\chi)^* \hookrightarrow \text{Hom}_{PGL_3}(\Pi, I_{Q_2}(st_\chi, \frac{1}{2})).$$

As in the proof of Proposition 7, we see by Proposition 4 that

$$\text{Hom}_{PGL_3}(\Pi, I_{Q_2}(st_\chi, \frac{1}{2})) \cong \text{Hom}_{L_2}(\Pi_1, st_\chi \otimes |\det|^{1/2}) \cong I_{P_2}(st_{\chi^2}, -\frac{1}{2})^*.$$

In particular,  $\theta(St_\chi)$  is a quotient of  $I_{P_2}(st_{\chi^2}, -\frac{1}{2})$ . Since  $\pi(E)$  is the unique irreducible quotient of  $I_{P_2}(st_{\chi^2}, -\frac{1}{2})$  [Mui97, Proposition 4.2(ii)], we have  $\Theta(St_\chi) \subset \{\pi(E)\}$ .

Further, as in Corollary 9(ii), there is a non-zero  $G_2 \times PGL_3$ -equivariant map

$$f : \Pi \longrightarrow \pi(E) \otimes I_{Q_2}(st_\chi, \frac{1}{2}),$$

and we claim that its image is the submodule  $\pi(E) \otimes St_\chi$ . If not, then  $\pi(E)$  will be an element of  $\Theta(J_2(st_\chi, \frac{1}{2}))$ . However, Theorem 11 shows that  $\Theta(J_2(st_\chi, \frac{1}{2})) = \{J_{P_2}(st_\chi, \frac{1}{2})\}$ . With this contradiction, we see that  $\Theta(St_\chi) = \{\pi(E)\}$  and the theorem is proved.  $\square$

THEOREM 15. We have

$$\Theta(St) = \{\pi(1)', \pi_{sc}[1]\}.$$

Here,  $\pi(1)'$  is a generic discrete series representation of  $G_2$  defined in [Mui97, Proposition 4.3(i)] and  $\pi_{sc}[1]$  is a non-generic supercuspidal representation defined in [HMS98].

This is certainly the most difficult part of this paper. The rest of the section is concerned with its proof. We need the following lemma, which will be used repeatedly.

LEMMA 16.

- i) Under  $\Theta$  and  $\Theta'$ , we have the correspondences

$$\{1\} \leftrightarrow \{J_{P_1}(\pi(1, 1), 1)\} \quad \text{and} \quad \{J_1(st, \frac{1}{2}), J_2(st, \frac{1}{2})\} \leftrightarrow \{J_{P_2}(st, \frac{1}{2})\}.$$

- ii)  $J_{P_1}(st, \frac{1}{2}) \notin \Theta(St)$ .

*Proof.* Part i is a special case of Theorem 11. The proof of part ii is quite involved. By the proof of [Mui97, Proposition 4.3],  $J_{P_1}(st, \frac{1}{2})$  is a submodule of  $I_{P_2}(\pi(| \cdot |^{-1}, | \cdot |))$ . Thus

$$\text{Hom}_{PGL_3 \times G_2}(\Pi, St \otimes J_{P_1}(st, \frac{1}{2})) \subset \text{Hom}_{PGL_3 \times M_2}(R_{P_2}(\Pi), St \otimes \pi(| \cdot |^{-1}, | \cdot |))$$

and it suffices to show that the latter space is zero. By Proposition 6, it suffices to show that the following spaces are zero:

$$\begin{aligned} & \text{Hom}(\text{Ind}_{Q_i \times GL_2}^{PGL_3 \times GL_2} C_c^\infty(GL_2), St \otimes \pi(| \cdot |^{-3/2}, | \cdot |^{1/2})), \quad i = 1, 2; \\ & \text{Hom}(\text{Ind}_{Q_0 \times B'}^{PGL_3 \times GL_2} C_c^\infty(GL_1), St \otimes \pi(| \cdot |^{-3/2}, | \cdot |^{1/2})); \\ & \text{Hom}(\Pi(PGL_6), St \otimes \pi(| \cdot |^{-1}, | \cdot |)). \end{aligned}$$

The vanishing of the first two terms above can be easily checked using Frobenius reciprocity. Indeed, the first term is equal to

$$\text{Hom}_{GL_2 \times GL_2}(C_c^\infty(GL_2), st \otimes \pi(| \cdot |^{-3/2}, | \cdot |^{1/2})) = 0,$$

and the second term is equal to

$$\text{Hom}_{Q_0 \times B'}(C_c^\infty(GL_1), 1 \otimes (| \cdot | \times | \cdot |^{-2})) \oplus \text{Hom}_{Q_0 \times B'}(C_c^\infty(GL_1), 1 \otimes (| \cdot |^{-1} \times 1)) = 0.$$

The last term can be treated with our methods. However, in order to avoid repetitive arguments, we shall relate it to a special case of the Howe correspondence as follows.

*Remark.* The minimal representation  $\Pi(PGL_6)$  is the Howe lift of the trivial representation of  $GL_1$ . In particular, the correspondence for the dual pair  $PGL_3 \times PGL_2 \subset PGL_6$  is a special case of the Howe correspondence for the dual pair  $GL_3 \times GL_2$ . Since a discrete series representation of  $GL_n$  does not appear in the Howe correspondence with  $GL_m$  unless  $m \geq n$ , the Steinberg representation (or any supercuspidal representation) of  $PGL_3$  cannot be a quotient of  $\Pi(PGL_6)$ .

The lemma is proved. □

We begin the proof of Theorem 15 by showing that  $\Theta(St)$  contains  $\pi_{sc}[1]$ . By Frobenius reciprocity and Proposition 5, we see that

$$\text{Hom}_{PGL_3 \times G_2}(\Pi, I_{Q_0}(\rho_0) \otimes \pi_{sc}[1]) \supset \text{Hom}_{G_2}(\Pi(D_4), \pi_{sc}[1]).$$

Now the restriction of  $\Pi(D_4)$  to  $G_2$  has been determined completely in [HMS98]. Their result states that

$$\Pi(D_4) = J_{P_1}(\pi(1, 1), 1) \oplus 2 \cdot J_{P_2}(st, \frac{1}{2}) \oplus \pi_{sc}[1].$$

Hence, we deduce that there is a non-zero map  $\Pi \rightarrow I_{Q_0}(\rho_0) \otimes \pi_{sc}[1]$  so that  $\Theta'(\pi_{sc}[1])$  contains some irreducible subquotient of  $I_{Q_0}(\rho_0)$ . Since the irreducible subquotients of  $I_{Q_0}(\rho_0)$  are precisely  $St$  and the representations of  $PGL_3$  mentioned in Lemma 16, part i, the lemma implies that  $St \in \Theta'(\pi_{sc}[1])$  and thus  $\pi_{sc}[1] \in \Theta(St)$ .

Next we find an upper bound for  $\Theta(St)$ . Since  $St$  is a submodule of  $I_{Q_2}(st, \frac{1}{2})$ , we have

$$\theta(St)^* \subset \text{Hom}_{PGL_3}(\Pi, I_{Q_2}(st, \frac{1}{2})) = \text{Hom}_{L_2}(R_{Q_2}(\Pi), st \otimes |\det|^{1/2}) = W.$$

Since  $R_{Q_2}(\Pi)$  has a natural filtration by Proposition 4, this induces a filtration on  $W$ :

$$0 = W_3 \subset W_2 \subset W_1 \subset W_0 = W,$$

such that

$$\begin{aligned} W_2/W_3 & \cong \text{Hom}_{L_2}(\Pi_3/\Pi_2, st \otimes |\det|^{1/2}), \\ W_1/W_2 & \subset \text{Hom}_{L_2}(\Pi_2/\Pi_1, st \otimes |\det|^{1/2}), \\ W_0/W_1 & \subset \text{Hom}_{L_2}(\Pi_1, st \otimes |\det|^{1/2}). \end{aligned}$$

Being a subspace of  $W$ ,  $\theta(St)^*$  inherits a filtration as well.

Now we investigate each of the terms above. The study of the last two terms is similar to the proof of Proposition 7. We simply state the result as follows.

LEMMA 17.

- i)  $\text{Hom}_{L_2}(\Pi_1, st \otimes |\det|^{1/2}) = I_{P_2}(st, -\frac{1}{2})^*$ .
- ii)  $W_1/W_2 = \text{Hom}_{L_2}(\Pi_2/\Pi_1, st \otimes |\det|^{1/2}) = 0$ .

We now consider the space

$$\text{Hom}_{L_2}(\Pi_3/\Pi_2, st \otimes |\det|^{1/2}) = \text{Hom}_{L_2}(\Pi(\mathfrak{L}_2), st).$$

Recall that  $\mathfrak{L}_2$  has derived group  $D_5$  and  $\Pi(\mathfrak{L}_2)$  is the minimal representation of  $D_5$  with the center of  $\mathfrak{L}_2$  acting trivially. Hence, we need to study the dual pair  $PGL_2 \times G_2 \subset D_5$ . Using the results of [MS97, § 2] and particularly [MS97, Proposition 2.3], we obtain the following.

LEMMA 18. *There is a short exact sequence of representations of  $G_2$ :*

$$0 \longrightarrow W'_1 \longrightarrow \text{Hom}_{L_2}(\Pi(\mathfrak{L}_2), st) \longrightarrow W'_2 \longrightarrow 0,$$

with

$$W'_1 \subset \Pi(D_4)^* \quad \text{and} \quad W'_2 \subset I_{P_1}(|\det|^{-1/2})^*.$$

Here  $\Pi(D_4)$  is the minimal representation of  $D_4$ .

Since  $\theta(St)^*$  is a subspace of  $W$ , the above considerations imply that there is a filtration

$$0 \subset V_3 \subset V_2 \subset V_1 = \theta(St)^\vee$$

such that

$$\begin{aligned} V_3 &\subset \Pi(D_4)^\vee, \\ V_2/V_3 &\subset I_{P_1}(|\det|^{-1/2})^\vee, \\ V_1/V_2 &\subset I_{P_2}(st, -\frac{1}{2})^\vee. \end{aligned}$$

In particular, we see that  $\theta(St)$  is of finite length. We need to identify each of these subquotients as much as possible.

Together with the fact that  $\pi_{\text{sc}}[1] \in \Theta(St)$ , Lemma 16 implies that  $V_3 = \pi_{\text{sc}}[1]$ . We now consider  $V_2/V_3$ . Since  $V_3$  is supercuspidal,  $V_2/V_3$  is actually a submodule of  $\theta(St)^\vee$ . If  $V_2/V_3$  is non-zero, then some irreducible quotient of  $I_{P_1}(|\det|^{-1/2})$  will be an element of  $\Theta(St)$ . Now the irreducible constituents of  $I_{P_1}(|\det|^{-1/2})$  are precisely the three representations of  $G_2$  in Lemma 16 [Mui97, Proposition 4.3]; in fact,  $J_{P_1}(st, 1/2)$  is the unique irreducible quotient. Hence, Lemma 16 implies that  $V_2/V_3 = 0$ .

By the above, we see that

$$\theta(St) = \pi_{\text{sc}}[1] \oplus (\text{a quotient of } I_{P_2}(st, -\frac{1}{2})).$$

We claim that the second term is non-zero. This follows from Corollary 20 (to be proved in the next section), which implies that  $\theta(St)$  has non-zero Whittaker functionals, and the fact that  $\pi_{\text{sc}}[1]$  is non-generic. Since  $I_{P_2}(st, -\frac{1}{2})$  has  $\pi[1]'$  as its unique irreducible quotient (cf. [Mui97, Proposition 4.3] and its proof), we conclude that  $\Theta(St) = \{\pi_{\text{sc}}[1], \pi[1]'\}$ . The theorem is proved.

### 10. Whittaker vectors

Before we come to the study of  $\Theta(\pi)$  for supercuspidal  $\pi$ , we need to establish some results on Whittaker vectors. The results of this section are independent of those in §§ 8 and 9 and thus could have been treated after § 7. Suppose that  $G \times G' \subset H$  is one of the following dual pairs, with  $G$  of

smaller dimension:

$$\begin{aligned} PGL_3 \times G_2 &\subset E_6 \\ G_2 \times PGSp_6 &\subset E_7. \end{aligned}$$

The second dual pair is not the subject matter of this paper, but will be crucial in the next section. Let  $U$  and  $U'$  be a maximal unipotent subgroup of  $G$  and  $G'$ , respectively, and let  $\psi$  and  $\psi'$  be unitary characters of  $U$  and  $U'$  which are in general position. Also, let  $\Pi$  be the minimal representation of  $H$ . We have the following proposition.

PROPOSITION 19. *As representations of  $G$ ,  $\Pi_{U',\psi'} \cong \text{ind}_U^G \psi$ , the Gelfand–Graev representation of  $G$ .*

*Proof.* This is similar to the proof for the finite field analogue [Gan99, Theorem 7.1]. We shall give a sketch for the case  $G \times G' = PGL_3 \times G_2$ . Let  $\psi'_2 = \psi'|_{N_2}$ . Then  $\psi'_2$  is a non-trivial character of  $N_2/(N_2 \cap N_1) \cong \mathbb{G}_a$ . Let  $N_\alpha \cong \mathbb{G}_a$  be the unipotent radical of the Borel subgroup  $M_2 \cap P_1$  of  $M_2$ . Then  $N_\alpha$  normalizes  $N_2$  and fixes the character  $\psi'_2$ . Thus  $\Pi_{N_2,\psi'_2}$  is naturally a representation of  $N_\alpha \times PGL_3$ . Let  $\psi_\alpha = \psi'|_{N_\alpha}$ ; it is a non-trivial character of  $N_\alpha$  and  $\Pi_{U',\psi'} = (\Pi_{N_2,\psi'_2})_{N_\alpha,\psi_\alpha}$ .

Using [MS97, Theorem 6.1], we see that there is an isomorphism of  $N_\alpha \times PGL_3$ -modules  $\Pi_{N_2,\psi'_2} \cong C_c^\infty(\mathcal{N})$  where  $\mathcal{N}$  is the set of  $3 \times 3$  nilpotent matrices. The action of  $PGL_3$  on  $C_c^\infty(\mathcal{N})$  is the action induced by conjugation whereas the action of  $b \in N_\alpha$  is induced by the action  $X \mapsto X + b \cdot X^2$ .

Now  $\mathcal{N}$  possesses a natural stratification by the rank of matrices. Thus  $C_c^\infty(\mathcal{N})$  has a natural filtration whose successive quotients are  $C_c^\infty(\mathcal{N}_i)$ , where  $\mathcal{N}_i$  is the set of nilpotent matrices of rank  $i$  ( $i = 0, 1$  or  $2$ ). Since  $N_\alpha$  acts trivially on  $\mathcal{N}_i$  for  $i \leq 1$ , we conclude that  $\Pi_{U',\psi'} \cong C_c^\infty(\mathcal{N}_2)_{N_\alpha,\psi_\alpha}$ . To prove the proposition, we need to identify the latter module as the Gelfand–Graev representation of  $PGL_3$ .

The action of  $N_\alpha \times PGL_3$  on  $\mathcal{N}_2$  is clearly transitive. Let

$$X_0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \in \mathcal{N}_2.$$

Without loss of generality, take  $U$  to be the group of upper triangular unipotent matrices in  $PGL_3$ . The stabilizer of  $X_0$  in  $N_\alpha \times PGL_3$  is contained in the subgroup  $N_\alpha \times U$  and the  $N_\alpha \times U$ -orbit  $\mathcal{N}'_2$  of  $X_0$  consists of the elements

$$X_b = \begin{pmatrix} 0 & 1 & b \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

Indeed, the action of

$$b \times \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \in N_\alpha \times U$$

is via  $X_0 \mapsto X_{x-y-b}$ .

The above discussion implies that

$$C_c^\infty(\mathcal{N}_2) \cong \text{ind}_{N_\alpha \times U}^{N_\alpha \times PGL_3} C_c^\infty(\mathcal{N}'_2)$$

where the action of  $N_\alpha \times U$  on  $C_c^\infty(\mathcal{N}'_2)$  is induced by the geometric action above. From this, we conclude that

$$C_c^\infty(\mathcal{N}_2)_{N_\alpha,\psi_\alpha} \cong \text{ind}_U^{PGL_3} (C_c^\infty(\mathcal{N}'_2)_{N_\alpha,\psi_\alpha}) \cong \text{ind}_U^{PGL_3} \psi$$

where  $\psi$  is the character of  $U$  given by

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto \psi_\alpha(y - x).$$

This proves the proposition. □

COROLLARY 20.

- i) If  $\pi^\vee$  is non-generic, then  $\theta(\pi)$  does not have any generic subquotients.
- ii) If  $\pi^\vee$  is generic, then  $\theta(\pi)$  has a unique irreducible generic subquotient. In particular,  $\theta(\pi)$  is non-zero.

### 11. Supercuspidal representations

In this section, we give the proof of Theorem 2. For convenience, let us repeat the statement of the theorem.

THEOREM 21. *Let  $\pi$  be a supercuspidal representation of  $PGL_3$ . Then:*

- i)  $\theta(\pi)$  is supercuspidal and has a unique generic summand;
- ii) if  $\pi$  is non-self-dual, then  $\theta(\pi)$  is irreducible.

Recall that we have a surjection  $\Pi \rightarrow \pi \otimes \theta(\pi)$ . By the exactness of the Jacquet functors, we have a surjection  $R_{P_2}(\Pi) \rightarrow \pi \otimes R_{P_2}(\theta(\pi))$ . By Proposition 6,  $\pi \otimes R_{P_2}(\theta(\pi))$  must be a quotient of  $\Pi(PGL_6)$ . However, following the remark in the proof of Lemma 16, we know that a square-integrable representation of  $PGL_3$  does not occur as a quotient of  $\Pi(PGL_6)$ . Thus  $R_{P_2}(\theta(\pi)) = 0$ .

We now show that  $R_{P_1}(\theta(\pi)) = 0$ . There is a surjection  $R_{P_1}(\Pi) \rightarrow \pi \otimes R_{P_1}(\theta(\pi))$  and, by the above, we know that  $R_{P_1}(\theta(\pi))$  is a supercuspidal representation of  $M_1$ . If it is not zero, then it is generic and so we have a surjection

$$R_{P_1}(\Pi)_{N_\beta, \psi_\beta} \rightarrow \pi \otimes R_{P_1}(\theta(\pi))_{N_\beta, \psi_\beta}.$$

Here,  $N_\beta \cong N_2/(N_1 \cap N_2) \cong \mathbb{G}_a$  is the unipotent radical of the Borel subgroup  $M_1 \cap P_2$  of  $M_1$ , and  $\psi_\beta$  is a non-trivial character of  $N_\beta$ . It remains to see that  $R_{P_1}(\Pi)_{N_\beta, \psi_\beta}$  does not contain supercuspidal representations of  $PGL_3$  as subquotients.

Next, note that  $(\Pi_{N_1})_{N_\beta, \psi_\beta}$  is isomorphic to  $(\Pi_{N_2, \psi'_2})_{N_\alpha}$ , where  $\psi'_2$  is the character of  $N_2$  as in the proof of Proposition 19. Since  $\Pi_{N_2, \psi'_2}$  was calculated there, we see that as representations of  $PGL_3$ ,  $R_{P_1}(\theta(\pi))_{N_\beta, \psi_\beta} \cong C_c^\infty(\mathcal{N})_{N_\alpha}$  where  $\mathcal{N}$  is the space of nilpotent  $3 \times 3$  matrices and  $N_\alpha$  is the unipotent radical of the Borel subgroup  $M_2 \cap P_1$  of  $M_2$ . Since  $C_c^\infty(\mathcal{N})$  has a natural filtration whose successive quotients are  $C_c^\infty(\mathcal{N}_i)$ , where  $\mathcal{N}_i$  is the set of nilpotent matrices of rank  $i$  ( $i = 0, 1, 2$ ), it is easy to see that

$$\begin{aligned} C_c^\infty(\mathcal{N}_0)_{N_\alpha} &\cong 1, \\ C_c^\infty(\mathcal{N}_1)_{N_\alpha} &\cong \text{Ind}_{Q_0}^{PGL_3} C_c^\infty(GL_1), \\ C_c^\infty(\mathcal{N}_2)_{N_\alpha} &\cong \text{ind}_U^{PGL_3} 1, \end{aligned}$$

where  $U$  is the unipotent radical of  $Q_0$ . In particular, these spaces do not contain supercuspidal representations of  $PGL_3$  as subquotients and thus  $R_{P_1}(\theta(\pi)) = 0$ .

We have shown that  $\theta(\pi)$  is supercuspidal and thus  $\theta(\pi)$  is semi-simple. Since  $\pi$  is supercuspidal, so is  $\pi^\vee$  and thus  $\pi^\vee$  is generic. Corollary 20 now immediately implies part i.

Finally, we assume that  $\pi$  is not self-dual. To show that  $\theta(\pi)$  is irreducible, it suffices to show that every summand of  $\theta(\pi)$  is generic. The key input here is a wonderful idea of Muic–Savin which was used in [MS00] for the study of symplectic-orthogonal theta lifts. Consider the dual pair  $G_2 \times PGSp_6 \subset E_7$ , and recall that  $PGSp_6$  has a Siegel parabolic subgroup  $P$  whose Levi subgroup is  $GL_3$ . If  $\Pi(E_7)$  denotes the minimal representation of  $E_7$ , then [MS97, Theorem 1.1] shows that  $\Pi$  is a  $PGL_3 \times G_2$ -quotient of  $R_P(\Pi(E_7))$ . Now suppose that  $\pi \otimes \sigma$  is a quotient of  $\Pi$ , with  $\sigma$  irreducible supercuspidal. Then, by Frobenius reciprocity, there is a non-zero map  $\Pi(E_7) \rightarrow I_P(\pi) \otimes \sigma$ . The key observation is that, when  $\pi$  is not self-dual,  $I_P(\pi)$  is irreducible and thus generic. However, by Proposition 19 applied to the dual pair  $G_2 \times PGSp_6$ , we deduce that  $\sigma^\vee$  is generic and thus so is  $\sigma$  (since  $\sigma$  is unitarizable). The proof of the theorem is now complete.

### 12. Depth zero lifts and a conjecture

The conjecture of Gross and Savin describes the theta correspondence in terms of Langlands parameters. However, it is natural to ask for an alternative characterization of  $\theta(\pi)$  in terms of  $\pi$ , since the Langlands parametrization for supercuspidal representations is not known for  $G_2$ . In this section, we determine  $\theta(\pi)$  for  $\pi$  a depth zero supercuspidal representation and state a conjecture in the case when  $\pi$  has positive depth, in terms of the construction of supercuspidal representations via compact induction.

Let us now assume that  $p \geq 5$  and suppose that  $\pi$  has depth zero. By results of Moy and Prasad [MP96],  $\pi \cong \text{ind}_K^{PGL_3} \sigma$ , where  $K$  is a hyperspecial maximal compact subgroup of  $PGL_3$  with first principal congruence subgroup  $K_1$  and  $\sigma$  is a cuspidal representation of  $K/K_1 = PGL_3(\mathbb{F}_q)$ . Fix an isomorphism  $\mathbb{Z}_{(p)}/\mathbb{Z} \cong \mathbb{F}_p^\times$ , where  $\mathbb{Z}_{(p)}$  is the localization of  $\mathbb{Z}$  at the prime  $p$ . Then, relative to this isomorphism, the irreducible generic representations of a connected reductive group  $G$  of adjoint type over  $\mathbb{F}_q$  can be parametrized by semisimple conjugacy classes of the dual group  $G^\vee$  [Car85]. In any case, let us write  $\sigma = \sigma(s)$  and  $\pi = \pi(s)$  if  $\sigma$  corresponds to the class of  $s$  in the above parametrization. Since  $\sigma$  is cuspidal,  $s$  is regular and elliptic.

Via the natural inclusion  $i : SL_3 \hookrightarrow G_2$ , the element  $i(s)$  is still regular and elliptic in  $G_2(\mathbb{F}_q)$ , and hence determines a cuspidal generic representation  $\sigma'(i(s))$  of  $G_2(\mathbb{F}_q)$ . Let  $\pi'(i(s)) = \text{ind}_{K'}^{G_2} \sigma'(i(s))$ , where  $K'$  is a hyperspecial maximal compact subgroup of  $G_2$ . Then  $\pi'(i(s))$  is an irreducible generic supercuspidal representation of  $G_2$  and we have the following.

PROPOSITION 22. *We have  $\theta(\pi(s)) = \pi'(i(s))$ .*

*Proof.* The dual pair  $PGL_3 \times G_2 \subset H$  can be defined over  $\mathbb{Z}$  and we may take  $K, K'$  and  $K_H$  to be the hyperspecial maximal compact subgroups corresponding to the respective groups of integer points. Then  $K_H \cap (PGL_3 \times G_2) = K \times K'$ . Let  $K_1$  be the first principal congruence subgroup of  $K_H$ . Then as a representation of  $K_H/K_1 \cong E_6(\mathbb{F}_q)$ ,  $\Pi^{K_1} \cong 1 \oplus R$ , where  $R$  denotes the reflection representation of  $E_6(\mathbb{F}_q)$  (cf. [Sav96] and [Gan99]). Hence,  $R$  is a  $K_H$ -equivariant quotient of  $\Pi$ .

It was shown in [Gan99, Theorem 10.2] that  $\sigma(s) \otimes \sigma'(i(s))$  is a  $K \times K'$ -equivariant quotient of  $R$ . Note that since  $\text{ind}_{K \times K'}^{PGL_3 \times G_2} \sigma(s) \otimes \sigma'(i(s))$  is irreducible, we have

$$\text{ind}_{K \times K'}^{PGL_3 \times G_2} \sigma(s) \otimes \sigma'(i(s)) = \text{Ind}_{K \times K'}^{PGL_3 \times G_2} \sigma(s) \otimes \sigma'(i(s)).$$

Hence, the result follows by Frobenius reciprocity. □

We now describe a conjecture for  $\pi$  of positive depth. Given a pair  $(T, \chi)$ , where  $T$  is an anisotropic maximal torus of  $PGL_3$  and  $\chi$  an admissible character of  $T$ , we can construct a supercuspidal representation  $\pi_{T, \chi}$  of  $PGL_3$ . Moreover, every supercuspidal representation of  $PGL_3$  arises in this way. Note that  $T$  is of the form  $K^\times/k^\times$  for some cubic field extension  $K/k$ . For simplicity, let us assume that  $K$  is Galois over  $k$ .

Now there is a natural conjugacy class of embedding of  $T$  into  $G_2$ . More precisely, via the isomorphism  $K^\times/k^\times \cong K_{\mathbb{N}=1}$ , one can embed  $T$  naturally into  $SL_3$ , and then into  $G_2$  via the natural conjugacy class of embedding  $SL_3 \hookrightarrow G_2$ . Hence,  $T$  can be regarded as an anisotropic maximal torus of  $G_2$ . By the recent results of [Yu01], one can construct a supercuspidal representation  $\pi'_{T,\chi}$  of  $G_2$  out of the data  $(T, \chi)$ . It is natural to make the following conjecture.

CONJECTURE.  $\theta(\pi_{T,\chi}) = \pi'_{T,\chi}$ .

Using the result of [Sav96], it may be possible to verify the conjecture in the case when  $K$  is the unramified cubic extension of  $k$ .

REFERENCES

- Bus01 C. Bushnell, *Representations of reductive  $p$ -adic groups: localization of Hecke algebras and applications*, J. London Math. Soc. **63** (2001), 364–386.
- BZ76 J. Bernstein and A. Zelevinsky, *Representations of the group  $GL_n(F)$ , where  $F$  is a non-archimedean local field*, Russian Math. Surveys **31** (1976), 1–68.
- Car85 R. Carter, *Finite groups of Lie type* (Wiley, 1985).
- Gan99 W. T. Gan, *Exceptional Howe correspondences over finite fields*, Compositio Math. **118** (1999), 323–344.
- GS97 B. H. Gross and G. Savin, *The dual pair  $PGL_3 \times G_2$* , Canadian Math. Bull. **40** (1997), 376–384.
- HMS98 J.-S. Huang, K. Magaard and G. Savin, *Unipotent representations of  $G_2$  arising from the minimal representation of  $D_4^E$* , J. Reine Angew Math. **500** (1998), 65–81.
- Key82 C. D. Keys, *On the decomposition of reducible principal series representations of  $p$ -adic Chevalley groups*, Pacific J. Math. **101** (1982), 351–388.
- MP96 A. Moy and G. Prasad, *Jacquet functors and unrefined minimal  $K$ -types*, Comment. Math. Helvetici **71** (1996), 96–121.
- MS97 K. Magaard and G. Savin, *Exceptional  $\Theta$ -correspondences I*, Compositio Math. **107** (1997), 89–123.
- Mui97 G. Muic, *The unitary dual of  $p$ -adic  $G_2$* , Duke Math. J. **90** (1997), 465–493.
- MS00 G. Muic and G. Savin, *Symplectic-orthogonal theta lifts of generic discrete series*, Duke Math. J. **101** (2000), 317–333.
- MVW87 C. Moeglin, M.-F. Vigneras and J.-L. Waldspurger, *Correspondances de Howe sur un corps  $p$ -adique*, Lecture Notes in Mathematics, vol. 1291 (Springer, 1987).
- Sav94 G. Savin, *Dual pair  $G_J \times PGL_2$* , Invent. Math. **118** (1994), 141–160.
- Sav96 G. Savin,  *$K$ -types of minimal representation ( $p$ -adic case)*, Glasnik Math. **31** (1996), 93–99.
- Sav99 G. Savin, *A class of supercuspidal representations of  $G_2(k)$* , Canadian Math. Bulletin **42** (1999), 393–400.
- Sha91 F. Shahidi, *Langlands’ conjecture on Plancherel measures for  $p$ -adic groups*, in *Harmonic analysis on reductive groups*, Progress in Mathematics, vol. 101 (Birkhäuser, 1991), 277–295.
- Yu01 J.-K. Yu, *Construction of tame supercuspidal representations*, J. Amer. Math. Soc. **14** (2001), 579–622.

Wee Teck Gan [wgan@math.ucsd.edu](mailto:wgan@math.ucsd.edu)

Department of Mathematics, Princeton University, Princeton, NJ 08544, USA

*Current address:* Department of Mathematics, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

Gordan Savin [savin@math.utah.edu](mailto:savin@math.utah.edu)

Department of Mathematics, University of Utah, Salt Lake City, UT 84112, USA