

# PERTURBATIONS OF NORM-ADDITIVE MAPS BETWEEN CONTINUOUS FUNCTION SPACES

LONGFA SUN AND YINGHUA SUN

*Hebei Key Laboratory of Physics and Energy Technology, School of Mathematics and  
Physics, North China Electric Power University, Baoding, P. R. China*

Corresponding author: Longfa Sun, email: [sun.longfa@ncepu.edu.cn](mailto:sun.longfa@ncepu.edu.cn)

(Received 7 September 2023)

*Abstract* Let  $X, Y$  be two locally compact Hausdorff spaces and  $T : C_0(X) \rightarrow C_0(Y)$  be a standard surjective  $\varepsilon$ -norm-additive map, i.e.

$$\| \|T(f) + T(g)\| - \|f + g\| \| \leq \varepsilon, \text{ for all } f, g \in C_0(X).$$

Then there exist a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon, \text{ for all } y \in Y, f \in C_0(X).$$

The estimate ' $\frac{3}{2}\varepsilon$ ' is optimal. And this result can be regarded as a new nonlinear extension of the Banach–Stone theorem.

*Keywords:* Banach–Stone theorem; norm-additive maps; linear isometries; continuous function spaces; stability

*2020 Mathematics subject classification:* Primary 46B04; 46B20  
Secondary 46E15

## 1. Introduction

Let  $X$  be a locally compact Hausdorff space. The space  $C_0(X)$  will stand for the Banach space of all continuous real-valued functions which vanish at infinity on  $X$  (i.e.  $\{x \in X : |f(x)| \geq \varepsilon\}$  is compact in  $X$  for every  $f \in C_0(X)$  and every  $\varepsilon > 0$ ) equipped with the supremum norm. The following result is well-known as the Banach–Stone theorem (see [2, 3, 21]).

**Theorem 1.1.** *Let  $X, Y$  be two locally compact Hausdorff spaces and  $T : C_0(X) \rightarrow C_0(Y)$  be a linear surjective isometry. Then there exists a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that*



$$T(f)(y) = \lambda(y)f(\varphi(y)), \text{ for all } y \in Y, f \in C_0(X).$$

The Banach–Stone theorem describes a deep fact that the linear metric structure of  $C_0(X)$  determines the topology of  $X$ . And it has found a large number of generalizations and variants in many different contexts (see [15] for a survey of corresponding results). The classical Mazur–Ulam theorem [18] states that every standard surjective isometry between two real Banach spaces must be linear. Thus, the existence of a standard surjective isometry between  $C_0(X)$  and  $C_0(Y)$  can also guarantee that  $X$  and  $Y$  are homeomorphic. Instead of isometries, Amir and Cambern investigated the linear isomorphisms between  $C_0(X)$  and  $C_0(Y)$ , where  $X, Y$  are compact Hausdorff spaces or locally compact Hausdorff spaces ([1, 5–7]). They showed that if the linear isomorphism  $T : C_0(X) \rightarrow C_0(Y)$  satisfies that  $\|T\| \cdot \|T^{-1}\| < 2$ , then the underlying spaces  $X$  and  $Y$  are homeomorphic, and the universal constant ‘2’ is optimal (see [9]).

In another direction, the nonlinear extension of the Banach–Stone theorem has attracted a large number of mathematicians’ attention (see [11–14, 17, 23]). Recently, Galego and Porto da Silva [14] studied the bijective coarse quasi-isometries between  $C_0(X)$  and  $C_0(Y)$  and they obtained an optimal nonlinear extension of the Banach–Stone theorem.

Let  $E, F$  be two Banach spaces. A map  $T : E \rightarrow F$  is said to be a coarse quasi-isometry (or coarse  $(M, \varepsilon)$ -quasi-isometry) for some constants  $M \geq 1$  and  $\varepsilon \geq 0$  provided

$$\frac{1}{M} \|u - v\| - \varepsilon \leq \|T(u) - T(v)\| \leq M \|u - v\| + \varepsilon,$$

for all  $u, v \in E$ .  $T$  is called an  $\varepsilon$ -isometry when  $M = 1$  and an isometry when  $M = 1$  and  $\varepsilon = 0$ . If  $T(0) = 0$ , then  $T$  is called standard.

**Theorem 1.2. (Galego-Porto da Silva).** *Let  $X, Y$  be two locally compact Hausdorff spaces and  $T : C_0(X) \rightarrow C_0(Y)$  be a standard bijective map such that both  $T$  and  $T^{-1}$  are coarse  $(M, \varepsilon)$ -quasi-isometries with  $M < \sqrt{2}$ . Then there exists a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that*

$$|MT(f)(y) - \lambda(y)f(\varphi(y))| \leq (M^2 - 1)\|f\| + \Delta\varepsilon, \text{ for all } y \in Y, f \in C_0(X),$$

where  $\Delta$  does not depend on  $f$  and  $y$ .

The upper bound  $\sqrt{2}$  on  $M$  is optimal even in the linear case when  $T$  is a linear isomorphism [9]. When  $M = 1$ , i.e.  $T : C_0(X) \rightarrow C_0(Y)$  is a bijective standard  $\varepsilon$ -isometry, Theorem 1.2 yields that

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq 2\varepsilon, \text{ for all } y \in Y, f \in C_0(X).$$

As it follows from a result of Omladič and Šemrl [20],  $T$  can be weakened as a standard surjective  $\varepsilon$ -isometry and the estimate ‘ $2\varepsilon$ ’ is optimal (see, also, [4, p. 360]).

In view of geometry, the isometry  $T$  from Banach space  $E$  to another Banach space  $F$  preserves the length of one diagonal of the parallelogram generated by two vectors. But, one may ask what happens if  $T$  preserves the length of another diagonal of the parallelogram instead, that is,

$$\|T(u) + T(v)\| = \|u + v\|, \text{ for all } u, v \in E.$$

By letting  $g = -f$  in the above equation, it is clear that  $T$  is an isometry with  $T(-f) = -T(f)$  and  $T(0) = 0$ . Thus, the Banach–Stone theorem (Theorem 1.1) still holds when  $T$  is surjective. Such transformations are called norm-additive maps and have stronger properties than isometries when the domain is symmetric. And these maps have been studied recently in [8, 10, 16, 19, 22].

Let  $E, F$  be two Banach spaces and  $T : E \rightarrow F$  be a map,  $\varepsilon \geq 0$ .  $T$  is called an  $\varepsilon$ -norm-additive map provided

$$\left| \|T(u) + T(v)\| - \|u + v\| \right| \leq \varepsilon, \text{ for all } u, v \in E.$$

In this paper, we mainly study the properties of the  $\varepsilon$ -norm-additive map between  $C_0(X)$  and  $C_0(Y)$  which is a natural and interesting generalization of norm-additive map to the perturbed case. It is worth noting that although the proof of Theorem 1.2 for the sharp estimate ‘ $2\varepsilon$ ’ of the surjective  $\varepsilon$ -isometries is very skilful [14], it cannot be applied to  $\varepsilon$ -norm-additive mappings for hunting the sharp estimate because the  $\varepsilon$ -norm-additive mapping between  $C_0(X)$  and  $C_0(Y)$  may be a strictly  $2\varepsilon$ -isometry (see Example 2.3).

We mainly prove that if  $X, Y$  are two locally compact Hausdorff spaces and  $T : C_0(X) \rightarrow C_0(Y)$  is a standard surjective  $\varepsilon$ -norm-additive map, then there exists a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon, \text{ for all } y \in Y, f \in C_0(X).$$

The constant ‘ $\frac{3}{2}$ ’ is optimal.

## 2. Main results

We start this section with the following observation which reveals the relationship between  $\varepsilon$ -isometries and  $\varepsilon$ -norm-additive maps on Banach spaces.

**Proposition 2.1.** *Suppose that  $E$  and  $F$  are Banach spaces and  $T : E \rightarrow F$  is an  $\varepsilon$ -norm-additive map. Then  $T$  is a  $2\varepsilon$ -isometry.*

**Proof.** For any  $u \in E$ , by the definition of  $T$ , we have

$$\left| \|T(u) + T(-u)\| - \|u - u\| \right| \leq \varepsilon,$$

i.e.

$$\|T(u) + T(-u)\| \leq \varepsilon.$$

For  $u, v \in E$ , we obtain that

$$\begin{aligned} \|T(u) - T(v)\| &= \|(T(u) + T(-u)) - (T(-u) + T(v))\| \\ &\leq \|T(-u) + T(v)\| + \|T(u) + T(-u)\| \\ &\leq \|u - v\| + 2\varepsilon, \end{aligned}$$

and

$$\|T(u) - T(v)\| = \|(T(u) + T(-u)) - (T(-u) + T(v))\|$$

$$\begin{aligned} &\geq \|T(-u) + T(v)\| - \|T(u) + T(-v)\| \\ &\geq \|u - v\| - 2\varepsilon. \end{aligned}$$

Thus,  $T$  is a  $2\varepsilon$ -isometry and the proof is complete. □

Although every  $\varepsilon$ -norm-additive map is actually a  $2\varepsilon$ -isometry, the converse is not true in general.

**Example 2.2.** Define  $T : c_0 \rightarrow c_0$  by  $T(u) = (\|u\|, u_1, u_2, \dots)$  for  $u = (u_n)_{n=1}^\infty \in c_0$ . Then  $T$  is a standard  $(0-)$ isometry, but it is not a  $\delta$ -norm-additive map for any  $\delta \geq 0$ .

The following example shows that the constant ‘2’ in Proposition 2.1 is sharp.

**Example 2.3.** Let  $X = \{a\}$ , then  $C(X) = \mathbb{R}$ . Fix  $\varepsilon > 0$ , define  $T : \mathbb{R} \rightarrow \mathbb{R}$  by

$$T(u) = \begin{cases} 0 & \text{if } u = 0, \\ -\frac{\varepsilon}{2} & \text{if } u = \varepsilon, \\ u + \frac{\varepsilon}{2} & \text{if } u \neq 0, \varepsilon. \end{cases}$$

Then  $T$  is a standard  $\varepsilon$ -norm-additive map. Note that  $||T(2\varepsilon) - T(\varepsilon)| - |2\varepsilon - \varepsilon|| = 2\varepsilon$ . This and Proposition 2.1 together show that  $T$  is a strictly  $2\varepsilon$ -isometry.

Let  $X, Y$  be two locally compact Hausdorff spaces and  $T : C_0(X) \rightarrow C_0(Y)$  be a standard surjective  $\varepsilon$ -norm-additive map. Combining Proposition 2.1, Theorem 1.2 and the Omladič–Šemrl’s theorem [20], we have the following result.

**Theorem 2.4.** *Let  $T : C_0(X) \rightarrow C_0(Y)$  be a standard surjective  $\varepsilon$ -norm-additive map. Then there exists a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that*

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq 4\varepsilon, \text{ for all } y \in Y, f \in C_0(X).$$

However, the constant in the estimate above is not the best. Our main goal is to obtain a sharp version of the above theorem by reducing  $4\varepsilon$  to  $\frac{3}{2}\varepsilon$ . And then we will show that  $\frac{3}{2}\varepsilon$  is optimal. To begin with, we establish the following useful lemmas.

**Lemma 2.5.** *Let  $x \in X$  and  $f_1, f_2, \dots, f_n \in C_0(X)$  with  $(f_1(x), f_2(x), \dots, f_n(x)) \neq 0$ . Then there exists a  $g \in C_0(X)$  with  $g(z) \leq 0$  for all  $z \in X$  such that*

$$\|g\| = -g(x) \text{ and } \|g + f_i\| = -g(x) - f_i(x), \forall i \in \{1, 2, \dots, n\}.$$

**Proof.** Let  $I = [-\|f_1\|, \|f_1\|] \times [-\|f_2\|, \|f_2\|] \times \dots \times [-\|f_n\|, \|f_n\|] \subset \mathbb{R}^n$ . Define  $F : X \rightarrow \mathbb{R}^n$  by

$$F(z) = (f_1(z), f_2(z), \dots, f_n(z)), \forall z \in X.$$

Then  $F$  is well defined and continuous. Let  $\alpha = \max_{1 \leq i \leq n} \|f_i\|$ . For every  $(u_1, u_2, \dots, u_n) \in I$ , let

$$h((u_1, u_2, \dots, u_n)) = \max_{1 \leq i \leq n} \{f_i(x) - u_i - 3\alpha, -3\alpha\}.$$

Then  $h : I \rightarrow \mathbb{R}$  is well defined. It is clear that  $h$  is continuous and  $-3\alpha \leq h(F(z)) \leq 0$  for every  $z \in X$ . By the Urysohn lemma, there exists a continuous function  $P : I \rightarrow [0, 1]$  such that  $P((0, 0, \dots, 0)) = 0$  and  $P(F(x)) = 1$ . Define  $g : X \rightarrow \mathbb{R}$  by

$$g(z) = (P \cdot h)(F(z)) (= P(F(z)) \cdot h(F(z))), \forall z \in X.$$

Then  $g$  is well defined. Since  $P, h, F$  are continuous,  $g$  is also continuous.

We assert that  $g \in C_0(X)$ . For any convergent net  $\{x_\lambda\}_{\lambda \in \Lambda}$  with  $x_\lambda$  converges to infinity, we have  $F(x_\lambda) \rightarrow (0, 0, \dots, 0) \in \mathbb{R}^n$ . Hence  $P(F(x_\lambda)) \rightarrow 0$ . Note that  $|h(F(x_\lambda))| \leq 3\alpha$  for any  $\lambda \in \Lambda$ , we have  $g(x_\lambda) \rightarrow 0$ . Thus  $g \in C_0(X)$ .

For every  $z \in X$ ,  $0 \leq P(F(z)) \leq 1$  and  $-3\alpha \leq h(F(z)) \leq 0$ . Thus  $-3\alpha \leq g(z) \leq 0$  for every  $z \in X$ . Note that  $g(x) = -3\alpha$ , then  $\|g\| = -g(x)$ . For every  $i \in \{1, 2, \dots, n\}$  and every  $z \in X$ , we have

$$\begin{aligned} \alpha &\geq g(z) + f_i(z) = P(F(z)) \cdot h(F(z)) + f_i(z) \\ &\geq h(F(z)) + f_i(z) \geq f_i(x) - f_i(z) - 3\alpha + f_i(z) \\ &= f_i(x) - 3\alpha = g(x) + f_i(x). \end{aligned}$$

Then

$$|g(z) + f_i(z)| \leq 3\alpha - f_i(x) = -g(x) - f_i(x), \forall z \in X.$$

This implies that  $\|g + f_i\| = -g(x) - f_i(x)$  and the proof is complete. □

Similar to Lemma 2.5, we have the following lemma.

**Lemma 2.6.** *Let  $x \in X$  and  $f_1, f_2, \dots, f_n \in C_0(X)$  with  $(f_1(x), f_2(x), \dots, f_n(x)) \neq 0$ . Then there exists a  $g \in C_0(X)$  with  $g(z) \geq 0$  for all  $z \in X$  such that*

$$\|g\| = g(x) \text{ and } \|g + f_i\| = g(x) + f_i(x), \forall i \in \{1, 2, \dots, n\}.$$

For every  $x \in X$  and  $f \in C_0(X)$ , let

$$\begin{aligned} P(f, x) &= \{(y, \lambda) \in Y \times \{\pm 1\} : \lambda T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon\}, \\ \bar{P}(f, x) &= \{(y, \lambda) \in Y \times \{\pm 1\} : \lambda T(f)(y) \leq f(x) + \frac{3}{2}\varepsilon\}. \end{aligned}$$

Put

$$P_+(x) = \bigcap_{f \in C_0(X), f(x) \geq 0} P(f, x), \quad P_-(x) = \bigcap_{f \in C_0(X), f(x) \leq 0} \bar{P}(f, x).$$

**Lemma 2.7.** *For every  $x \in X$ ,  $P_+(x)$  is non-empty.*

**Proof.** Fix  $x \in X$ , the proof is divided into three steps.

Step I. We prove that for  $f \in C_0(X)$  with  $f(x) > \frac{3}{2}\varepsilon$ ,  $P(f, x)$  is a non-empty compact set. Let

$$P_1 = \{y \in Y : T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon\}, \quad P_2 = \{y \in Y : -T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon\}.$$

Since  $T$  is an  $\varepsilon$ -norm-additive map,

$$2\|T(f)\| - \|f\| = \|T(f) + T(f)\| - \|f + f\| \leq \varepsilon,$$

i.e.

$$\|f\| - \frac{1}{2}\varepsilon \leq \|T(f)\| \leq \|f\| + \frac{1}{2}\varepsilon.$$

This yields that at least one of the  $P_1, P_2$  is non-empty. Note that  $f(x) > \frac{3}{2}\varepsilon$ ,  $P_1$  and  $P_2$  are two compact sets. Since  $P(f, x) = P_1 \times \{1\} \cup P_2 \times \{-1\}$ ,  $P(f, x)$  is a non-empty compact subset of  $Y \times \{\pm 1\}$ .

Step II. Fix a  $f_0 \in C_0(X)$  with  $f_0(x) > \frac{3}{2}\varepsilon$ , we prove that  $P(f_0, x) \cap P(f, x)$  is a non-empty compact set for every  $f \in C_0(X)$  with  $f(x) \geq 0$ . It is clear that  $P(f, x)$  is closed in  $Y \times \{\pm 1\}$  and hence  $(P(f_0, x) \cap P(f, x)) \subset P(f_0, x)$  is compact. It remains to prove that  $P(f_0, x) \cap P(f, x)$  is non-empty. By Lemma 2.5, there exists a  $g \in C_0(X)$  such that

$$\|g\| = -g(x), \quad \|g + f\| = -g(x) - f(x), \quad \|g + f_0\| = -g(x) - f_0(x).$$

Since  $T$  is an  $\varepsilon$ -norm-additive map, one has

$$\|T(g)\| \geq \|g\| - \frac{1}{2}\varepsilon, \quad \|f + g\| + \varepsilon \geq \|T(f) + T(g)\|, \quad \|f_0 + g\| + \varepsilon \geq \|T(f_0) + T(g)\|.$$

Pick  $(y, \lambda) \in Y \times \{\pm 1\}$  such that  $\|T(g)\| = \lambda T(g)(y)$ . Then

$$\begin{aligned} -g(x) - f_0(x) + \varepsilon &= \|g + f_0\| + \varepsilon \geq \|T(g) + T(f_0)\| \geq \lambda T(g)(y) + \lambda T(f_0)(y) \\ &\geq \|g\| - \frac{1}{2}\varepsilon + \lambda T(f_0)(y) = -g(x) - \frac{1}{2}\varepsilon + \lambda T(f_0)(y). \end{aligned}$$

This implies  $-\lambda T(f_0)(y) \geq f_0(x) - \frac{3}{2}\varepsilon$ . Similarly, we can get  $-\lambda T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon$ . Thus  $(y, -\lambda) \in P(f_0, x) \cap P(f, x)$  and  $P(f_0, x) \cap P(f, x)$  is a non-empty compact subset of  $Y \times \{\pm 1\}$ .

Step III. We prove that the set family  $\{P(f, x) \cap P(f_0, x)\}_{f \in C_0(X), f(x) \geq 0}$  has the finite intersection property. Fix  $f_1, f_2, \dots, f_n \in C_0(X)$  with  $f_i(x) \geq 0$  for every  $i \in \{1, 2, \dots, n\}$ , by Lemma 2.5, there exists a  $g \in C_0(X)$  such that

$$\|g\| = -g(x) \quad \text{and} \quad \|g + f_i\| = -g(x) - f_i(x), \quad \forall i \in \{0, 1, 2, \dots, n\}.$$

Pick  $(y, \lambda) \in Y \times \{\pm 1\}$  such that  $\|T(g)\| = \lambda T(g)(y)$ . For every  $i \in \{1, 2, \dots, n\}$ , by the same argument as in Step II, we obtain

$$(y, -\lambda) \in P(f_i, x) \cap P(f_0, x).$$

Thus

$$\bigcap_{f \in C_0(X), f(x) \geq 0} (P(f, x) \cap P(f_0, x)) \neq \emptyset.$$

Note that

$$P_+(x) = \bigcap_{f \in C_0(X), f(x) \geq 0} P(f, x) = \bigcap_{f \in C_0(X), f(x) \geq 0} (P(f, x) \cap P(f_0, x)).$$

The proof is complete. □

The following Lemma 2.8 is analogous to Lemma 2.7 and we omit its proof.

**Lemma 2.8.** *For every  $x \in X$ ,  $P_-(x)$  is non-empty.*

**Remark 2.9.** Fix  $x \in X$ , it is not difficult to verify that if  $(y, \lambda) \in P_+(x)$ , then  $(y, -\lambda) \notin P_+(x)$ . Suppose on the contrary that  $(y, \lambda), (y, -\lambda) \in P_+(x)$ , pick  $f \in C_0(X)$  with  $f(x) > \frac{3}{2}\varepsilon$ , then

$$\lambda T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon > 0, \quad -\lambda T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon > 0.$$

This leads to a contradiction. By a similar argument as above, we can see that if  $(y, \lambda) \in P_-(x)$ , then  $(y, -\lambda) \notin P_-(x)$ .

From now on, we assume that  $T : C_0(X) \rightarrow C_0(Y)$  is a standard surjective  $\varepsilon$ -norm-additive map. For every  $y \in Y$  and  $g \in C_0(Y)$ , define

$$Q(g, y) = \{(x, \lambda) \in X \times \{\pm 1\} : \lambda f(x) \geq g(y) - \frac{3}{2}\varepsilon, f \in T^{-1}(g)\},$$

$$\bar{Q}(g, y) = \{(x, \lambda) \in X \times \{\pm 1\} : \lambda f(x) \leq g(y) + \frac{3}{2}\varepsilon, f \in T^{-1}(g)\}.$$

Put

$$Q_+(y) = \bigcap_{g \in C_0(Y), g(y) \geq 0} Q(g, y), \quad Q_-(y) = \bigcap_{g \in C_0(Y), g(y) \leq 0} \bar{Q}(g, y).$$

**Lemma 2.10.** *For every  $y \in Y$ ,  $Q_+(y)$ ,  $Q_-(y)$  both are non-empty.*

**Proof.** Let  $y \in Y$ , we just prove  $Q_+(y)$  is non-empty, the case for  $Q_-(y)$  is similar. The proof is divided into three steps.

Step I. Assume that  $g \in C_0(Y)$  with  $g(y) > \frac{3}{2}\varepsilon$ . We first prove  $Q(g, y)$  is a non-empty compact set. By Lemma 2.5, there exists  $\bar{g} \in C_0(Y)$  such that

$$\|\bar{g}\| = -\bar{g}(y) \quad \text{and} \quad \|\bar{g} + g\| = -\bar{g}(y) - g(y).$$

Since  $T$  is surjective, there exists  $h \in C_0(X)$  such that  $T(h) = \bar{g}$ . Find  $(x, \lambda) \in X \times \{\pm 1\}$  such that  $\|h\| = \lambda h(x)$ . For every  $f \in T^{-1}(g)$ ,

$$\begin{aligned} \|\bar{g}\| - \frac{1}{2}\varepsilon + \lambda f(x) &\leq \|h\| + \lambda f(x) = \lambda(h(x) + f(x)) \\ &\leq \|h + f\| \leq \|T(h) + T(f)\| + \varepsilon \end{aligned}$$

$$\begin{aligned} &= \|\bar{g} + g\| + \varepsilon = -\bar{g}(y) - g(y) + \varepsilon \\ &= \|\bar{g}\| - g(y) + \varepsilon. \end{aligned}$$

Thus  $-\lambda f(x) \geq g(y) - \frac{3}{2}\varepsilon$ . This implies that  $(x, -\lambda) \in Q(g, y)$  and  $Q(g, y)$  is non-empty. For every  $f \in T^{-1}(g)$ , let

$$\begin{aligned} R(f)_1 &= \{x \in X : f(x) \geq g(y) - \frac{3}{2}\varepsilon\}, \quad R(f)_2 = \{x \in X : -f(x) \geq g(y) - \frac{3}{2}\varepsilon\}, \\ R(f) &= \{(x, \lambda) \in X \times \{\pm 1\} : \lambda f(x) \geq g(y) - \frac{3}{2}\varepsilon\}. \end{aligned}$$

By the above argument,  $R(f) \neq \emptyset$  and  $R(f) = R(f)_1 \times \{1\} \cup R(f)_2 \times \{-1\}$ . Since  $g(y) > \frac{3}{2}\varepsilon$ ,  $R(f)_1, R(f)_2$  are compact. By the Tychonoff theorem,  $R(f)$  is a non-empty compact subset of  $X \times \{\pm 1\}$ . Note that  $Q(g, y) = \bigcap_{f \in T^{-1}(g)} R(f)$ , this implies that  $Q(g, y)$  is a non-empty compact set.

Step II. Fix  $g_0 \in C_0(Y)$  with  $g_0(y) > \frac{3}{2}\varepsilon$ , we will show that for every  $g \in C_0(Y)$  with  $g(y) \geq 0$ ,  $Q(g_0, y) \cap Q(g, y)$  is a non-empty compact set. Note that  $Q(g, y)$  is closed in  $X \times \{\pm 1\}$ . Hence  $(Q(g_0, y) \cap Q(g, y)) \subset Q(g_0, y)$  is compact. By Lemma 2.5, there exists a  $\bar{g} \in C_0(Y)$  such that

$$\|\bar{g}\| = -\bar{g}(y), \|\bar{g} + g\| = -\bar{g}(y) - g(y), \|\bar{g} + g_0\| = -\bar{g}(y) - g_0(y).$$

Pick  $h \in C_0(X)$  such that  $T(h) = \bar{g}$ . Find  $\{x, \lambda\} \in X \times \{\pm 1\}$  such that  $\|h\| = \lambda h(x)$ . For every  $f \in T^{-1}(g)$ ,

$$\begin{aligned} \|\bar{g}\| - \frac{1}{2}\varepsilon + \lambda f(x) &\leq \|h\| + \lambda f(x) = \lambda(h(x) + f(x)) \\ &\leq \|h + f\| \leq \|T(h) + T(f)\| + \varepsilon \\ &= \|\bar{g} + g\| + \varepsilon = -\bar{g}(y) - g(y) + \varepsilon \\ &= \|\bar{g}\| - g(y) + \varepsilon. \end{aligned}$$

This implies that  $-\lambda f(x) \geq g(y) - \frac{3}{2}\varepsilon$  and  $(x, -\lambda) \in Q(g, y)$ . Similarly, we can show  $(x, -\lambda) \in Q(g_0, y)$ . Therefore,  $Q(g_0, y) \cap Q(g, y)$  is a non-empty compact set.

Step III. We check that the set family  $\{Q(g, y) \cap Q(g_0, y)\}_{g \in C_0(Y), g(y) \geq 0}$  has the finite intersection property. Fix  $g_1, g_2, \dots, g_n \in C_0(Y)$  with  $g_i(y) \geq 0$  for every  $i \in \{1, 2, \dots, n\}$ , by Lemma 2.5, there exists a  $\bar{g} \in C_0(Y)$  such that

$$\|\bar{g}\| = -\bar{g}(y), \|\bar{g} + g_i\| = -\bar{g}(y) - g_i(y), \forall i \in \{0, 1, 2, \dots, n\}.$$

Pick  $h \in C_0(X)$  such that  $T(h) = \bar{g}$ . Find  $\{x, \lambda\} \in X \times \{\pm 1\}$  such that  $\|h\| = \lambda h(x)$ . By the same argument as in Step II, we have

$$(x, -\lambda) \in Q(g_0, y) \quad \text{and} \quad (x, -\lambda) \in Q(g_i, y), \quad \forall i \in \{1, 2, \dots, n\}.$$



Thus

$$\bigcap_{g \in C_0(Y), g(y) \geq 0} (Q(g, y) \cap Q(g_0, y)) \neq \emptyset.$$

Note that

$$Q_+(y) = \bigcap_{g \in C_0(Y), g(y) \geq 0} Q(g, y) = \bigcap_{g \in C_0(Y), g(y) \geq 0} (Q(g, y) \cap Q(g_0, y)).$$

Then  $Q_+(y)$  is non-empty and the proof is complete. □

**Lemma 2.11.** *For every  $x \in X$ ,  $P_+(x)$  is a singleton.*

**Proof.** Let  $x \in X$ , by Lemma 2.7,  $P_+(x)$  is non-empty. Let  $(y, \lambda) \in P_+(x)$ , by Lemma 2.10,  $Q_+(y)$  and  $Q_-(y)$  are non-empty. For every  $f \in C_0(X)$  with  $f(x) > 3\varepsilon$ , we have

$$\lambda T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon.$$

If  $\lambda = 1$ , then  $T(f)(y) \geq f(x) - \frac{3}{2}\varepsilon > 0$ . For every  $(x', \mu) \in Q_+(y)$  and every  $f \in C_0(X)$  with  $f(x) > 3\varepsilon$ , one has

$$\mu f(x') \geq T(f)(y) - \frac{3}{2}\varepsilon \geq f(x) - 3\varepsilon > 0. \tag{2.1}$$

We assert that  $x = x'$ . Suppose on the contrary that  $x \neq x'$ , by the Urysohn lemma, there exists a  $f \in C_0(X)$  with  $f(x) > 3\varepsilon$  and  $f(x') = 0$ . This contradicts to (2.1). And hence we have  $x = x'$  and  $\mu = 1$ . Thus  $Q_+(y) = \{(x, \lambda)\}$ .

If  $\lambda = -1$ , then  $T(f)(y) \leq -(f(x) - \frac{3}{2}\varepsilon) < 0$ . For every  $(x', \mu) \in Q_-(y)$  and every  $f \in C_0(X)$  with  $f(x) > 3\varepsilon$ , one has

$$\mu f(x') \leq T(f)(y) + \frac{3}{2}\varepsilon \leq -(f(x) - \frac{3}{2}\varepsilon) + \frac{3}{2}\varepsilon = -f(x) + 3\varepsilon < 0.$$

By the same argument as above, we have  $x = x'$  and  $\mu = -1$ . Thus  $Q_-(y) = \{(x, \lambda)\}$ .

We assert that  $P_+(x)$  is a singleton. Suppose on the contrary that there exist  $(y_1, \lambda_1) \neq (y_2, \lambda_2) \in P_+(x)$ . By Remark 2.9,  $y_1 \neq y_2$ . Without loss of generality, we assume that  $\lambda_1 = 1$ . Then  $Q_+(y_1) = \{(x, 1)\}$ . By the Urysohn lemma, there exists  $g \in C_0(Y)$  such that  $g(y_1) > 3\varepsilon$  and  $g(y_2) = 0$ . Then for every  $f \in C_0(X)$  with  $T(f) = g$ , we have

$$f(x) \geq T(f)(y_1) - \frac{3}{2}\varepsilon = g(y_1) - \frac{3}{2}\varepsilon > 0.$$

Thus

$$0 = \lambda_2 g(y_2) = \lambda_2 T(f)(y_2) \geq f(x) - \frac{3}{2}\varepsilon \geq g(y_1) - 3\varepsilon > 0.$$

This leads to a contradiction. Hence  $P_+(x)$  is a singleton and the proof is complete. □

Similar to Lemma 2.11, we have the following result.

**Lemma 2.12.** *For every  $x \in X$ ,  $P_-(x)$  is a singleton.*

**Lemma 2.13.** *For every  $x \in X$ ,  $P_+(x) = P_-(x)$ .*

**Proof.** Fix  $x \in X$ , by Lemmas 2.11 and 2.12, there exist  $y_1, y_2 \in Y$  and  $\lambda_1, \lambda_2 \in \{\pm 1\}$  such that  $P_+(x) = (y_1, \lambda_1)$  and  $P_-(x) = (y_2, \lambda_2)$ . According to the proofs of Lemma 2.11 and 2.12, we have

$$(x, \lambda_1) = \begin{cases} Q_+(y_1), & \text{if } \lambda_1 = 1, \\ Q_-(y_1), & \text{if } \lambda_1 = -1, \end{cases} \quad \text{and} \quad (x, \lambda_2) = \begin{cases} Q_-(y_2), & \text{if } \lambda_2 = 1, \\ Q_+(y_2), & \text{if } \lambda_2 = -1. \end{cases}$$

We assert that  $y_1 = y_2$ . Suppose on the contrary that  $y_1 \neq y_2$ , by the Urysohn lemma, there exists a  $g \in C_0(Y)$  such that

$$\lambda_1 g(y_1) > \lambda_2 g(y_2) + 3\varepsilon > 6\varepsilon. \tag{2.2}$$

For every  $f \in C_0(X)$  with  $T(f) = g$ ,

$$\begin{cases} f(x) \geq g(y_1) - \frac{3}{2}\varepsilon & \text{if } \lambda_1 = 1, \\ -f(x) \leq g(y_1) + \frac{3}{2}\varepsilon & \text{if } \lambda_1 = -1. \end{cases}$$

Thus

$$f(x) \geq \lambda_1 g(y_1) - \frac{3}{2}\varepsilon. \tag{2.3}$$

On the other hand, for every  $f \in C_0(X)$  with  $T(f) = g$ ,

$$\begin{cases} f(x) \leq -g(y_2) + \frac{3}{2}\varepsilon & \text{if } \lambda_2 = 1, \\ -f(x) \geq -g(y_2) - \frac{3}{2}\varepsilon & \text{if } \lambda_2 = -1. \end{cases}$$

Thus

$$f(x) \leq -\lambda_2 g(y_2) + \frac{3}{2}\varepsilon. \tag{2.4}$$

Combining (2.2), (2.3) and (2.4), we have

$$-\lambda_2 g(y_2) + \frac{3}{2}\varepsilon \geq f(x) \geq \lambda_1 g(y_1) - \frac{3}{2}\varepsilon > \lambda_2 g(y_2) + \frac{3}{2}\varepsilon.$$

This implies that  $\lambda_2 g(y_2) < 0$ . By (2.2),  $\lambda_2 g(y_2) > 0$ . This leads to a contradiction and hence  $y_1 = y_2$ .

Next we prove that  $\lambda_1 = \lambda_2$ . Choose  $f \in C_0(X)$  such that  $f(x) > 3\varepsilon$ , then

$$\lambda_1 T(f)(y_1) \geq f(x) - \frac{3}{2}\varepsilon > \frac{3}{2}\varepsilon, \quad \lambda_2 T(-f)(y_1) \leq -f(x) + \frac{3}{2}\varepsilon < -\frac{3}{2}\varepsilon.$$

If  $\lambda_1 \neq \lambda_2$ , then

$$\varepsilon \geq \|T(f) + T(-f)\| \geq |T(f)(y_1) + T(-f)(y_1)| > 3\varepsilon.$$

This leads to a contradiction. Hence  $\lambda_1 = \lambda_2$  and  $P_+(x) = P_-(x)$ . The proof is complete.  $\square$

By a similar argument as Lemmas 2.11, 2.12 and 2.13, we can show the following result. To simplify this article, we omit its proof.

**Lemma 2.14.** *For every  $y \in Y$ ,  $Q_+(y) = Q_-(y)$  is a singleton.*

**Remark 2.15.** For every  $x \in X$ ,  $y \in Y$  and  $\lambda \in \{\pm 1\}$ , by Lemmas 2.11, 2.12, 2.13 and 2.14, one has

$$\{(x, \lambda)\} = Q_+(y) \iff \{(y, \lambda)\} = P_+(x).$$

Now we are ready to show the main result of this paper.

**Theorem 2.16.** *Let  $T : C_0(X) \rightarrow C_0(Y)$  be a standard surjective  $\varepsilon$ -norm-additive map. Then there exists a homeomorphism  $\varphi : Y \rightarrow X$  and a continuous function  $\lambda : Y \rightarrow \{\pm 1\}$  such that*

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon, \text{ for all } y \in Y, f \in C_0(X).$$

**Proof.** For  $y \in Y$ , by Lemma 2.14, there exist  $x \in X$ ,  $\lambda \in \{\pm 1\}$  such that

$$Q_+(y) = \{(x, \lambda)\}. \tag{2.5}$$

Define

$$\varphi(y) = x, \quad \lambda(y) = \lambda,$$

where  $x, y, \lambda$  satisfy (2.5). Then  $\varphi : Y \rightarrow X$  and  $\lambda : Y \rightarrow \{\pm 1\}$  are well defined. It follows from Remark 2.15 that  $\varphi$  is bijective. In what follows, we show that

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon, \text{ for all } y \in Y, f \in C_0(X). \tag{2.6}$$

Given  $y \in Y$  and  $f \in C_0(X)$ . The proof is divided into two cases.

Case I.  $f(\varphi(y)) \geq 0$ . By Remark 2.15, we have  $\{(y, \lambda(y))\} = P_+(\varphi(y))$ . Then

$$\lambda(y)T(f)(y) \geq f(\varphi(y)) - \frac{3}{2}\varepsilon. \tag{2.7}$$

We assert that

$$\lambda(y)T(f)(y) \leq f(\varphi(y)) + \frac{3}{2}\varepsilon. \tag{2.8}$$

Suppose on the contrary that  $\lambda(y)T(f)(y) > f(\varphi(y)) + \frac{3}{2}\varepsilon \geq 0$ . If  $\lambda(y) = 1$ , we have  $\{(\varphi(y), 1\} = Q_+(y)$  and

$$f(\varphi(y)) \geq T(f)(y) - \frac{3}{2}\varepsilon > f(\varphi(y)).$$

This leads to a contradiction. If  $\lambda(y) = -1$ , we have  $T(f)(y) < 0$  and  $\{(\varphi(y), -1\} = Q_+(y) = Q_-(y)$ . Then

$$-f(\varphi(y)) \leq T(f)(y) + \frac{3}{2}\varepsilon < -f(\varphi(y)).$$

This is a contradiction. Combining (2.7) and (2.8), we have

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon.$$

Case II.  $f(\varphi(y)) \leq 0$ . By Remark 2.15 again, we have  $\{(y, \lambda(y))\} = P_+(\varphi(y)) = P_-(\varphi(y))$ . Then

$$\lambda(y)T(f)(y) \leq f(\varphi(y)) + \frac{3}{2}\varepsilon. \tag{2.9}$$

We assert that

$$\lambda(y)T(f)(y) \geq f(\varphi(y)) - \frac{3}{2}\varepsilon. \tag{2.10}$$

Suppose on the contrary that  $\lambda(y)T(f)(y) < f(\varphi(y)) - \frac{3}{2}\varepsilon \leq 0$ . If  $\lambda(y) = 1$ , we have  $\{(\varphi(y), 1\} = Q_+(y) = Q_-(y)$  and

$$f(\varphi(y)) \leq T(f)(y) + \frac{3}{2}\varepsilon < f(\varphi(y)).$$

This leads to a contradiction. If  $\lambda(y) = -1$ , we have  $Tf(y) > 0$  and  $\{(\varphi(y), -1\} = Q_+(y) = Q_-(y)$ . Then

$$-f(\varphi(y)) \geq T(f)(y) - \frac{3}{2}\varepsilon > -f(\varphi(y)).$$

This is a contradiction. Combining (2.9) and (2.10), we have

$$|T(f)(y) - \lambda(y)f(\varphi(y))| \leq \frac{3}{2}\varepsilon.$$

Next we prove that  $\varphi$  is a homeomorphism and  $\lambda$  is continuous. Suppose that  $\{y_\alpha\}_{\alpha \in \Lambda} \subset Y$  is a convergent net with  $y_\alpha \rightarrow y$ . Choose a compact neighbourhood  $U$  of  $y$ , without loss of generality, we can assume that  $\{y_\alpha\}_{\alpha \in \Lambda} \subset U$ . By the Urysohn lemma, there exists

a  $g \in C_0(Y)$  such that  $g|_U \equiv 3\varepsilon + 1$ . By (2.6), for any  $f \in C_0(X)$  with  $Tf = g$ , we obtain that

$$\begin{aligned} \frac{3}{2}\varepsilon &\geq |T(f)(y_\alpha) - \lambda(y_\alpha)f(\varphi(y_\alpha))| \\ &\geq |g(y_\alpha)| - |f(\varphi(y_\alpha))| \\ &= 3\varepsilon + 1 - |f(\varphi(y_\alpha))|. \end{aligned}$$

This implies that  $|f(\varphi(y_\alpha))| \geq \frac{3}{2}\varepsilon + 1$  and  $\{\varphi(y_\alpha)\}_{\alpha \in \Lambda}$  is contained in the compact set  $\{x \in X : |f(x)| \geq \frac{3}{2}\varepsilon + 1\}$ .

By (2.6), for  $\alpha \in \Lambda$ , we have

$$|Tf(y_\alpha) - \lambda(y_\alpha)f(\varphi(y_\alpha))| \leq \frac{3}{2}\varepsilon, \text{ for all } f \in C_0(X). \tag{2.11}$$

For any convergent subnet  $\{\varphi(y_{\alpha'})\}_{\alpha' \in \Lambda'}$  of  $\{\varphi(y_\alpha)\}_{\alpha \in \Lambda}$  with  $\varphi(y_{\alpha'}) \rightarrow x$ , let  $\{\lambda(y_{\alpha''})\}_{\alpha'' \in \Lambda''}$  be a convergent subnet of  $\{\lambda(y_{\alpha'})\}_{\alpha' \in \Lambda'}$  with  $\lambda(y_{\alpha''}) \rightarrow \lambda$ . By (2.11), we have

$$|T(f)(y) - \lambda f(x)| \leq \frac{3}{2}\varepsilon, \text{ for all } f \in C_0(X).$$

This implies that  $\{(x, \lambda)\} = Q_+(y)$  and hence  $\varphi(y) = x$ . Thus  $\varphi$  is continuous. By the same argument, we can get  $\varphi^{-1}$  is also continuous. Hence  $\varphi$  is a homeomorphism. For any convergent subset  $\{\lambda(y_{\alpha'})\}_{\alpha' \in \Lambda'}$  of  $\{\lambda(y_\alpha)\}_{\alpha \in \Lambda}$  with  $\lambda(y_{\alpha'}) \rightarrow \lambda$ , by (2.11) again, we have

$$|T(f)(y) - \lambda f(\varphi(y))| \leq \frac{3}{2}\varepsilon, \text{ for all } f \in C_0(X).$$

This implies that  $\{(\varphi(y), \lambda)\} = Q_+(y)$  and  $\lambda(y) = \lambda$ . Hence  $\lambda : Y \rightarrow \{\pm 1\}$  is continuous. The proof is complete.  $\square$

The following example shows that the estimate ‘ $\frac{3}{2}\varepsilon$ ’ in Theorem 2.16 is optimal.

**Example 2.17.** Let  $X = \{x_1, x_2\}$  with the discrete topology, then  $C(X) = \ell_\infty^2$ . Let

$$U_1 = \{(a, b) \in C(X) : b \leq -\frac{1}{2}\varepsilon, a + b \geq -\frac{1}{2}\varepsilon, (a, b) \neq (\varepsilon, -\frac{1}{2}\varepsilon)\}$$

and

$$U_2 = C(X) \setminus (U_1 \cup \{(0, 0), (\varepsilon, -\frac{1}{2}\varepsilon)\}).$$

Define  $T : C(X) \rightarrow C(X)$  by

$$T((a, b)) = \begin{cases} (0, 0) & (a, b) = (0, 0), \\ (\varepsilon, \varepsilon) & (a, b) = (\varepsilon, -\frac{1}{2}\varepsilon), \\ (a - \frac{1}{2}\varepsilon, b) & (a, b) \in U_1, \\ (a, b - \frac{1}{2}\varepsilon) & (a, b) \in U_2. \end{cases}$$

It is not difficult to verify that  $T$  is a standard surjective map. In what follows, we show that  $T$  is an  $\varepsilon$ -norm-additive map. By the definition of  $T$ ,

$$\|T(f) - f\| \leq \frac{1}{2}\varepsilon, \forall f \in C(X), f \neq (\varepsilon, -\frac{1}{2}\varepsilon).$$

Then for any  $f, g \in C(X)$  with  $f, g \neq (\varepsilon, -\frac{1}{2}\varepsilon)$ , we have

$$\begin{aligned} \|\|T(f) + T(g)\| - \|f + g\|\| &\leq \|(T(f) + T(g)) - (f + g)\| \\ &\leq \|T(f) - f\| + \|T(g) - g\| \leq \varepsilon. \end{aligned}$$

The left case is  $f = (\varepsilon, -\frac{1}{2}\varepsilon)$  and  $g \in C(X)$ . When  $g = (0, 0)$  or  $(\varepsilon, -\frac{1}{2}\varepsilon)$ , it is clear that

$$\|T(f) + T(g)\| = \|f + g\|.$$

Let  $g = (a, b) \in U_1$ , we have

$$|b + \varepsilon| \leq |b| \leq a + \frac{1}{2}\varepsilon.$$

Then

$$\begin{aligned} \|T(f) + T(g)\| &= \max\{|a + \frac{1}{2}\varepsilon|, |b + \varepsilon|\} = a + \frac{1}{2}\varepsilon, \\ \|f + g\| &= \max\{|a + \varepsilon|, |b - \frac{1}{2}\varepsilon|\} = a + \varepsilon. \end{aligned}$$

Thus

$$\|\|T(f) + T(g)\| - \|f + g\|\| = \frac{1}{2}\varepsilon.$$

Let  $g = (a, b) \in U_2$ , we have

$$\begin{aligned} \|\|T(f) + T(g)\| - \|f + g\|\| &= \|\|(a + \varepsilon, b + \frac{1}{2}\varepsilon)\| - \|(a + \varepsilon, b - \frac{1}{2}\varepsilon)\|\| \\ &\leq \|(a + \varepsilon, b + \frac{1}{2}\varepsilon) - (a + \varepsilon, b - \frac{1}{2}\varepsilon)\| \\ &= \|(0, \varepsilon)\| = \varepsilon. \end{aligned}$$

Therefore,  $T$  is an  $\varepsilon$ -norm-additive map. The homeomorphism  $\varphi : X \rightarrow X$  and  $\lambda : X \rightarrow \{\pm 1\}$  which satisfy Theorem 2.16 are

$$\varphi = Id_X, \lambda(x_i) = 1, \text{ for } i = 1, 2.$$

When  $f = (\varepsilon, -\frac{1}{2}\varepsilon)$ , then

$$|T(f)(x_2) - \lambda(x_2)f(\varphi(x_2))| = |\varepsilon - (-\frac{1}{2}\varepsilon)| = \frac{3}{2}\varepsilon.$$

This implies that the estimate  $\frac{3}{2}\varepsilon$  in Theorem 2.16 is optimal.

**Remark 2.18.** The assumption of surjectivity of  $T$  in Theorem 2.16 is essential in general. For instance, let  $X = \{a\}$ ,  $Y = \{b, c\}$  be two discrete topological spaces, then  $C(X) = \mathbb{R}$  and  $C(Y) = \ell_\infty^2$ . Define  $T : \mathbb{R} \rightarrow \ell_\infty^2$  by  $T(x) = (x, \sin x)$  for all  $x \in \mathbb{R}$ . Then  $T$  is a norm-additive map, but  $X$  and  $Y$  are not homeomorphic.

**Acknowledgements.** The authors are grateful to the referee and the editor for their constructive comments and helpful suggestions.

**Funding Statement.** Longfa Sun is supported by the National Natural Science Foundation of China (Grant no. 12101234), the Natural Science Foundation of Hebei Province (Grant no. A2022502010), the Fundamental Research Funds for the Central Universities (Grant no. 2023MS164) and the China Scholarship Council.

## References

- (1) D. Amir, On isomorphisms of continuous function spaces, *Israel J. Math.* **3**(4) (1965), 205–210.
- (2) S. Banach, *Théorie des Opérations Lineaires* Vol. 110, 2nd edn. (Chelsea Publishing Company, New York, 1963), pp. 259.
- (3) E. Behrends, *M-Structure and the Banach-Stone Theorem* Vol. 736, 1st edn. Lecture Notes in Mathematics (Springer-Verlag Heidelberg, Berlin, 1979), 220 pp. 4549.
- (4) Y. Benyamini and J. Lindenstrauss, Geometric nonlinear functional analysis I. *Amer. Math. Soc. Colloquium Publications*, Volume 48 (Amer. Math. Soc, Providence, RI, 2000).
- (5) M. Cambern, A generalized Banach-Stone theorem, *Proc. Amer. Math. Soc.* **17**(2) (1966), 396–400.
- (6) M. Cambern, On isomorphisms with small bound, *Proc. Amer. Math. Soc.* **18**(6) (1967), 1062–1066.
- (7) M. Cambern, Isomorphisms of  $C_0(Y)$  onto  $C_0(X)$ , *Pacific J. Math.* **35**(2) (1970), 307–312.
- (8) L. Chen, Y. Dong and B. Zheng, On norm-additive maps between the maximal groups of positive continuous functions, *Result Math.* **74** (2019), 7.
- (9) H. B. Cohen, A bound-two isomorphism between  $C(X)$  Banach spaces, *Proc. Amer. Math. Soc.* **50**(1) (1975), 215–217.
- (10) Y. Dong, L. Li, L. Molnár and N. -C. Wong, Transformations preserving the norm of means between positive cones of general and commutative  $C^*$ -algebras, *J. Operator Theory* **88**(2) (2022), 365–406.
- (11) Y. Dutrioux and N. Kalton, Perturbations of isometries between  $C(K)$ -spaces, *Studia Math.* **166** (2) (2005), 181–197.
- (12) R. Górak, Coarse version of the Banach-Stone theorem, *J. Math. Anal. Appl.* **377**(1) (2011), 406–413.
- (13) R. Górak, Perturbations of isometries between Banach spaces, *Studia Math.* **207** (1) 2011, 47–58.
- (14) E. M. Galego and A. L. Porto da Silva, An optimal nonlinear extension of Banach-Stone theorem, *J. Funct. Anal.* **271**(8) (2016), 2166–2176.
- (15) M. I. Garrido and J. A. Jaramillo, Variations on the Banach-Stone theorem, *Extracta Math.* **17**(3) (2002), 351–383.
- (16) M. Hosseini and J. J. Font, Real-linear isometries and jointly norm-additive maps on function algebras, *Mediterr. J. Math.* **13**(4) (2016), 1933–1948.

- (17) K. Jarosz, Nonlinear generalizations of the Banach-Stone theorem, *Studia Math.* **93** (2) 1989, 97–107.
- (18) S. Mazur and S. Ulam, Sur les transformations isométriques d'espaces vectoriels normés, *C.R. Acad. Sci. Paris* **194** (1932), 946–948.
- (19) L. Molnár, Spectral characterization of Jordan-Segal isomorphisms of quantum observables, *J. Operator Theory* **83**(1) (2020), 179–195.
- (20) M. Omladič and P. Šemrl, On non linear perturbations of isometries, *Math. Ann.* **303**(1) (1995), 617–628.
- (21) M. H. Stone, Applications of the theory of Boolean rings to general topology, *Trans. Amer. Math. Soc.* **41**(3) (1937), 375–481.
- (22) T. Tonev and R. Yates, Norm-linear and norm-additive operators between uniform algebras, *J. Math. Anal. Appl.* **357**(1) (2009), 45–53.
- (23) I. A. Vestfrid, Non-surjective coarse version of the Banach-Stone theorem, *Ann. Funct. Anal.* **11** (2020), 634–642.