



Ontogenetic and spatial variability in parasite communities of white shrimp *Penaeus setiferus* (Decapoda: Penaeidae)

Research Article

Cite this article: Zuidema SR, de Buron I, Kingsley-Smith PR, Hill-Spanik KM, Fanani N, Kendrick MR (2023). Ontogenetic and spatial variability in parasite communities of white shrimp *Penaeus setiferus* (Decapoda: Penaeidae). *Parasitology* **150**, 230–239. <https://doi.org/10.1017/S0031182022001597>


Received: 3 August 2022
Revised: 24 October 2022
Accepted: 26 October 2022
First published online: 12 December 2022

Keywords:

Apostomatida; parasite interactions; Rhabditida; shrimp health; shrimp life history; Trypanorhyncha

Author for correspondence:

Michael R. Kendrick,
E-mail: kendrickm@dnr.sc.gov

Sarah R. Zuidema^{1,2}, Isaure de Buron², Peter R. Kingsley-Smith¹,
Kristina M. Hill-Spanik², Natalia Fanani² and Michael R. Kendrick¹ 

¹South Carolina Department of Natural Resources, Marine Resources Research Institute, 217 Fort Johnson Road, Charleston, SC 29412, USA and ²Department of Biology, Grice Marine Laboratory, College of Charleston, 205 Fort Johnson Road, Charleston, SC 29412, USA

Abstract

Understanding the combined effects of multi-parasite infections on their hosts is necessary for documenting parasite impacts and is particularly important for developing effective management strategies for economically important organisms. The white shrimp *Penaeus setiferus* supports important recreational and commercial fisheries along the southeastern and Gulf coasts of the United States and occupies an important ecological niche in estuarine and offshore habitats throughout these regions. The goal of this study was to identify and assess ontogenetic and spatial variation in white shrimp parasite communities and their relation to shrimp health. We used a series of trawl surveys in tidal creek and open water habitats of an estuary in the southeastern USA to collect and identify parasites of white shrimp using morphological and DNA sequencing techniques. Parasite communities in white shrimp were composed of organisms belonging to 6 classes: Conoidasida (gregarines), Oligohymenophorea (apostome and sessilid ciliates), Microsporea (meiodihaplophasids), Chromadorea (rhabditids), Cestoda (cyclophyllideans, lecanoccephalideans and trypanorhynchids) and Trematoda (plagiorchiids). Parasite communities differed significantly among white shrimp life stages and localities. Furthermore, the health condition known as black gill occurred in some shrimp and was significantly related to parasite community structure. Infection metrics for the apostome ciliate *Hyalophysa lynni*, the trypanorhynch larvae *Prochristianella* sp. and the rhabditid larvae *Hysterothylacium* sp. were significantly different between shrimp exhibiting and not exhibiting black gill. These results highlight the importance of understanding parasite communities and the potential interactive effects of multiple parasite infections on shrimp health.

Introduction

Single-species parasite infections have received significant attention (e.g. Lefèvre *et al.*, 2008; Lafferty, 2017), but multi-parasite infections are increasingly understood as significant drivers of host health and disease outbreaks (Hellard *et al.*, 2015; Clay and Rudolf, 2019). Impacts of parasite infections may include decreased host immune function, increased parasite fitness (Pedersen and Fenton, 2007; Budischak *et al.*, 2015), induction of severe pathologies (Gleichsner *et al.*, 2018) and altered host response to future infections (Rodríguez *et al.*, 1999; Graham, 2008). Co-infections by multiple parasites can either relieve or exacerbate these impacts on host health at both the individual and population level (Reckardt and Kerth, 2009; Hellard *et al.*, 2015), such that a community-level understanding of parasite infections is important for quantifying parasite impacts on host health.

Commercially important finfish and crustaceans commonly serve as hosts for a diversity of parasites in freshwater and marine environments and wherever possible their presence and impacts should be incorporated into management approaches (Marcogliese, 2004; Byers, 2021). For example, over the course of their life cycle, white shrimp, *Penaeus setiferus* (L., 1767) (Decapoda: Penaeidae) serve as hosts to a variety of parasites, including but not limited to, platyhelminthes, nematodes, ciliates and microsporidians (e.g. Overstreet, 1978; Domínguez-Machín *et al.*, 2011; Del Río-Rodríguez *et al.*, 2013). Penaeid shrimp are one of the most important fisheries and aquaculture industries worldwide (FAO, 2018). Specifically, adult white shrimp support important commercial and recreational fisheries along the coasts of the southeastern USA (Gillet, 2008) and the Gulf of Mexico (Muncy, 1984; NMFS, 2020), including \$13.7 million annually from the Georgia Bight in the western Atlantic (Kendrick *et al.*, 2021). Additionally, white shrimp provide a food source for many recreationally and ecologically important finfish, including red drum, *Sciaenops ocellatus* (see Scharf and Schlicht, 2000) and spotted seatrout, *Cynoscion nebulosus* (see Fujiwara *et al.*, 2016). As such, white shrimp comprise an integral part of the estuarine ecosystem and economy and monitoring population health is necessary to ensure sustainable management.

© South Carolina Department of Natural Resources and the Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Commercial landings of white shrimp in parts of the Georgia Bight have declined during the last 2 decades (Kendrick *et al.*, 2021), coinciding with numerous biotic and abiotic changes in the estuarine waters of the southeastern USA (e.g. Hollebhone, 2006; Shearman and Lentz, 2010). Among these biotic changes is the increased prevalence of a condition known as black gill (Fowler *et al.*, 2018; Swinford and Anderson, 2021; Tuckey *et al.*, 2021). This condition refers to the melanization of gill tissues that occurs as part of an immune response against irritants or pathogens (Vaseeharan and Ramasamy, 2003; Johnson *et al.*, 2011; Burnett and Burnett, 2015; Karthikeyan *et al.*, 2015; Frischer *et al.*, 2022). Black gill in wild-caught penaeid shrimp has been associated with the presence of the apostome ciliate *Hyalophysa lynni* (see Landers *et al.*, 2020), but also occurs in response to heavy metals (Ghate, 1984), bacteria (Vaseeharan and Ramasamy, 2003) and fungi (Karthikeyan *et al.*, 2015).

Parasites of white shrimp in the Georgia Bight, including South Carolina (SC) have not been intensively surveyed, and their impacts on white shrimp health at the individual and population level, and consequently on commercial landings, warrant investigation. The objectives of this study were to: (1) identify parasite community composition of white shrimp in the Charleston Harbor, SC; (2) document ontogenetic and spatial variability in the parasite communities of white shrimp; and (3) assess the relationships between parasites and black gill in white shrimp.

Material and methods

Collection of post-larval white shrimp

Zooplankton sampling was conducted at the surface during nocturnal flood tides that coincided with neap tides to correspond with the timing of shrimp ingress (DeLancey *et al.*, 1994; Wenner *et al.*, 2005). Collections occurred between 6 June 2018

and 1 September 2018 and consisted of 18 sampling events. Two 0.75 m diameter plankton nets (500 μ m mesh) were deployed for 30–75 min (depending on changes in tidal flow or the timing of the flood tide with respect to darkness) from a floating dock in Charleston Harbor, SC, USA (Fig. 1). Plankton samples were washed with seawater using a 500 μ m mesh sieve and transported in an aerated container of seawater to the laboratory. Within 24 h, post-larval shrimp were isolated using a dissecting microscope and identified as penaeid following the key of Johnson and Allen (2005). These specimens were presumed to be white shrimp based on the timing of ingress, which largely does not overlap with any other penaeid species in SC (DeLancey *et al.*, 1994). Specimens were flattened between 2 slides and examined whole under a compound microscope to detect parasites.

Collection of juvenile, subadult and adult white shrimp

Juvenile, subadult and adult white shrimp were collected within the greater Charleston Harbor watershed at each of 3 localities (Fig. 1); tidal creek localities in the Ashley River watershed ($n=3$), tidal creek localities in the Wando River watershed ($n=3$) and open water localities in Charleston Harbor ($n=2$). Specimens were collected from tidal creek localities monthly from June to September 2018 ($n=15$ per site) and following their egress from tidal creeks in late-summer, from open water localities monthly ($n=23$ per site) from August to November 2018. Collections from tidal creek localities were made using a 3.05 m otter trawl (0.95 cm stretch mesh net) towed for 5 min at ~ 2 knots. Collections from open water localities were made using a 6.2 m otter trawl (2.5 cm stretch mesh net) for 15 min at ~ 2 knots. Bottom water temperature ($^{\circ}$ C) and salinity (psu) were recorded at each site on each collection date using a handheld YSI Pro2030. Shrimp were measured from the tip of the rostrum to the tip of the telson, and size classes were defined according to




Fig. 1. Map of sampling localities in the greater Charleston Harbor, South Carolina, USA: Wando River, Ashley River and Charleston Harbor. Credit: Gary Sundin, South Carolina Department of Natural Resources (SCDNR).

Whitaker and Kingsley-Smith (2014) as follows: juvenile (≤ 114 mm length), subadult (115–126 mm length) and adult (≥ 127 mm length).

Identification of parasite community composition

Specimens were brought back on ice to the laboratory and examined for parasites immediately, or stored at -20°C for later processing. To collect parasites, individual shrimp were submerged in a dish of deionized water and, using a dissecting microscope, the carapace was removed and the organs excised from the cephalothoracic cavity. The hepatopancreas and feeding palps were deconstructed to isolate parasites, while the nerve cord was resected from the full length of the shrimp, flattened between 2 slides and examined with a compound microscope under $100\times$ total magnification. The anterior caecum was resected from the digestive tract and opened along with the stomach, intestine and hindgut to examine their contents. A $\sim 1\text{--}2\text{ mm}^2$ squash of the abdominal muscle was performed between 2 slides and examined at $100\times$ total magnification. Smears were performed when gonads or muscle were infected and stained with Giemsa. Parasite specimens were fixed in 95% ethanol for molecular analysis.

As morphological identification of larval parasites can be challenging, morphotypes of parasite groups were first developed using morphological characteristics and infection sites (Overstreet, 1973, 1978; Deardorff and Overstreet, 1981; Palm, 2004; Sokolova *et al.*, 2015). These morphotypes were screened using genetic sequencing for further identification. Subsamples of each morphotype group (except for the sessilid ciliates, which were not preserved) were processed for molecular analyses. DNA extractions ($n = 26$) were performed using a DNeasy Blood and Tissue kit (Qiagen, Valencia, CA, USA) following manufacturer protocols, except that elution volumes were reduced to $40\ \mu\text{L}$. For each taxonomic group, primers were chosen that would amplify a gene region most likely to provide species-level identifications (Table 1).

For platyhelminthes, a $25\ \mu\text{L}$ total reaction contained $1\times$ Promega GoTaq[®] Flexi PCR Buffer (Madison, WI, USA), $0.1\times$ Invitrogen Rediload[™] loading buffer (Thermo Fisher Scientific, Waltham, MA, USA), $1.5\ \text{mM}$ MgCl_2 , $0.2\ \text{mM}$ dNTPs, each primer at $0.4\ \mu\text{M}$, $0.05\ \text{U}\ \mu\text{L}^{-1}$ Promega GoTaq[®] DNA polymerase and $1\ \mu\text{L}$ template DNA. Cycling was performed as described by Jensen and Bullard (2010), with one alteration in that cycling began with a 5 min denaturation at 95°C rather than at 94°C . For ciliates, PCR reagents and concentrations were the same as above except $0.5\ \mu\text{M}$ of each primer was used. Cycling was performed as described by Guo *et al.* (2012). For nematodes, a $25\ \mu\text{L}$ total reaction contained $1\times$ PCR Buffer (Thermo Fisher Scientific),

$0.1\times$ Invitrogen Rediload[™] loading buffer (Thermo Fisher Scientific), $2.5\ \text{mM}$ MgCl_2 , $0.2\ \text{mM}$ dNTPs, each primer at $0.2\ \mu\text{M}$, $0.2\ \mu\text{M}$ Invitrogen Platinum[™] Taq DNA polymerase and $1\ \mu\text{L}$ template DNA. Cycling was performed as follows: 5 min at 96°C then 40 cycles at 96°C for 40 s, 45°C for 40 s and 72°C for 40 s, then 72°C for 5 min. For microsporidians, PCR reagents and concentrations were the same as for the platyhelminthes PCR (described above), except that $1\ \mu\text{M}$ of each primer was used in a $20\ \mu\text{L}$ total reaction volume. Cycling performed was as follows: 4 min at 94°C then 35 cycles at 94°C for 30 s, 55°C for 30 s and 72°C for 2 min and then at 72°C for 5 min. For gregarines, several small subunit ribosomal RNA gene primers (Leander *et al.*, 2003; Rueckert and Leander, 2009; Diakin *et al.*, 2016) were tested but did not result in amplification.

All PCR products were electrophoresed on 1% agarose gels ($100\ \text{V}$, 30 min) stained with GelRed[®] (Biotium, Inc., Hayward, CA, USA), and visualized under UV light. Products were cleaned using ExoSAP-IT[™] (Affymetrix, Santa Clara, CA, USA) following manufacturer protocols. Cleaned products were sent to Eurofins MWG Operon LLC (Louisville, KY, USA) for direct, bi-directional sequencing using the primers shown in Table 1. Complementary sequences were compared to one another and to their chromatograms using Sequencher[®] version 5.4 (Gene Codes, Ann Arbor, MI, USA). Resulting sequences were compared against the NCBI GenBank database using the Basic Local Alignment Search Tool (BLAST; Altschul *et al.*, 1990). A 99% sequence similarity threshold was used for species-level identifications for all taxa; assignments to genus level were based on percent similarity and the BLAST taxonomy report.

Parasite prevalence, intensities and densities were defined according to Bush *et al.* (1997). Mean intensities could not be determined for the cyclophyllidean, the ciliates (apostome and sessilid), the gregarine or the microsporidian (Table 2). Prevalence only was recorded for the cyclophyllidean, gregarine and microsporidian. For the ciliates, relative abundance was calculated for the apostome and estimated for the sessilid (see below). Mean intensities and relative abundances are expressed as \pm standard error (S.E.).

Macroscopic assessment of black gill condition

A black gill score was determined macroscopically for individual shrimp by evaluating the darkest portion of gills in each specimen in the field using a colour standard scale comprised of increasingly pigmented paint swatches as follows based on the hexadecimal colour code: 1 = #DFD3C3, 2 = #DACAB2, 3 = #CCB79B, 4 = #C0A98B, 5 = #AB9579, 6 = #947F65, 7 = #6E543C, 8 = #5A4A3D, 9 = #564536 and 10 = #48423C. A black gill score from 1 (no melanization) to 10 (darkest melanization) was

Table 1. Primers used for amplification and sequencing of white shrimp parasites

Phylum	Gene region	Primer name	Primer orientation	Primer sequence (5'-3')	Primer reference
Ciliophora	SSU	18SF-ciliate-458	+	AGCAGGCGCGHAAATTRCCAATCY	Frischer <i>et al.</i> (2017)
		18SR-ciliate-1260	-	CCGTGTTGAGTCAAATTAAGCCG	Frischer <i>et al.</i> (2017)
Microsporidia	SSU	ss18f	+	CACCAGGTTGATTCTGCC	Ghosh and Weiss (2009)
		ss1482r	-	GGTTACCTTGTTACGACTT	Ghosh and Weiss (2009)
Nematoda	cox2	211F	+	TTTTCTAGTTATATAGATTGRTTTYAT	Nadler and Hudspeth (2000)
		210R	-	CACCAACTCTTAAAATTATC	Nadler and Hudspeth (2000)
Platyhelminthes	LSU	LSU5	+	TAGGTCGACCCGCTGAAYTTAAGCA	Jensen and Bullard (2010)
		1200R	-	GCATAGTTCCACATCTTTCGG	Jensen and Bullard (2010)

The gene region used was based on the most informative marker for species-level identification: large subunit ribosomal RNA (LSU rRNA) gene for platyhelminthes, partial small subunit (SSU) rRNA gene for ciliates and microsporidians, and mitochondrial cytochrome c oxidase II (cox2) for nematodes. For primer orientation, +, sense; -, antisense.

Table 2. Parasite taxa identifications, BLAST results and quantitative factors used in analyses. N/A = sequencing was unsuccessful or not done. P = prevalence. Only species with intensity (I) or relative abundance (RA) were included in NMDS and associated analyses.

Phylum	Class	Order	Specimen ID	Organism name	BLAST results		
					% ID	Accession No.	Quantitative factors
Apicomplexa	Conoidasida	Gregarinasina ^a	Gregarine	N/A	N/A	P	
Ciliophora	Oligohymenophorea	Apostomatida	<i>Hyalophysa lynni</i>	100	KX906567	P/RA	
Ciliophora	Oligohymenophorea	Sessilida	<i>Zoothamnium</i> sp.	N/A	N/A	P/RA	
Microsporidia	Microsporea	Meiodiaphlophasida	<i>Agmasoma penaei</i>	100	KF549987	P	
Nematoda	Chromadorea	Rhabditida	<i>Hysterathyliacium</i> sp.	96–97	AF179916	P/I	
Platyhelminthes	Cestoda	Cyclophyllidea	<i>Parorchites</i> sp.	96	KP893423	P	
Platyhelminthes	Cestoda	Lecanicephalidea	<i>Polypocephalus</i> sp.	100	GQ470200	P/I	
Platyhelminthes	Cestoda	Trypanorhyncha	<i>Kotorella pronosoma</i> ^c	100	MK558802	N/A	
Platyhelminthes	Cestoda	Trypanorhyncha	<i>Prochristianella</i> sp.	98	DQ642783	P/I	
Platyhelminthes	Trematoda	Plagiiorchiida	<i>Opecoeloides fimbriatus</i>	100	KJ0011211	P/I	

^aGregarinasina is a subclass of Apicomplexa, but specimens could not be identified past this level of classification, and are consequently referenced only as 'gregarines'.

^bAs of this publication, the GenBank record had not been updated to reflect the findings of Landers *et al.* (2020) who associated this GenBank accession number with *Hyalophysa lynni*. Specimen ID agrees with organism name where BLAST results returned 99% similarity or higher, otherwise only genus level was confirmed.

^cA single specimen was identified so it was included as part of the trypanorhynch for the purpose of analyses.

recorded for each specimen, with scores >5 categorizing individuals as shrimp with black gill and scores of ≤5 categorizing individuals as shrimp without black gill.

Assessment of infection by gill parasites

To examine gill parasites (i.e., the sessilid and apostome ciliates), the 4 most posterior gill filaments of the haphazardly chosen left or right side were removed from each juvenile, subadult and adult shrimp ($n = 543$) and preserved in 3–5% formalin or frozen in deionized water until later examination (Martin *et al.*, 2000). Wet mounts of the gill filaments were observed using light microscopy at 200× total magnification. Gill filaments were subdivided into 3 non-overlapping fields of view (FOVs) each of 0.785 mm². Smaller gills typically allowed for only 1 or 2 FOVs per filament. The number of trophonts of the apostome ciliate in each FOV was recorded and relative abundance calculated as the number of apostome ciliates per mm² of gill tissue. For sessilid ciliates, relative abundance was estimated according to the number of FOVs showing stalks of these parasites; this value was corrected for the number of FOVs available in an individual shrimp.

Statistical analyses

Parasite prevalence, intensities and abundances were used for statistical analyses, which were performed in R version 4.0.3 (R Core Team, 2019) (Table 2). As only 1 individual of the trypanorhynch *Kotorella pronosoma* (Stossich, 1901) was observed, it was combined with the other trypanorhynch (*Prochristianella* sp.) in the analyses. Due to low parasite infections in white shrimp post-larvae, statistical analyses were restricted to juvenile, subadult and adult life stages. When testing how individual parasites relate to the patterns of black gill prevalence, analyses were restricted to collections made from the open water localities during the black gill season (August through October).

To assess variability among parasite communities across ontogenetic and spatial factors (i.e., localities), parasite abundances were first averaged for each trawl and then standardized using the Wisconsin double standardization tool, which standardizes species by the maximum value and then standardizes localities by totals, using the 'vegan' package (Oksanen *et al.*, 2019). Bray–Curtis distance matrices of parasite abundances were then developed and used to create non-metric multidimensional scaling (NMDS) plots with 95% data ellipses to examine shrimp parasite communities among localities. Water temperature (°C), salinity (psu), shrimp length and black gill score were tested for correlation with multivariate distances using permutation tests with 999 iterations and visualized as vectors using the 'envfit' function in the 'vegan' package (Oksanen *et al.*, 2019). Parasite taxa names overlaid onto the ordination plot illustrate where taxa correspond to one another within the ordination space. Permutational analyses of variance (PERMANOVA) and corresponding pairwise *post hoc* tests using the 'RVAideMemoire' package (Hervé, 2021) were also conducted to examine specific differences among parasite communities of shrimp groups (i.e., juvenile vs subadult vs adult and Ashley River vs Wando River vs open water Charleston Harbor localities).

Indicator species analyses were performed using multi-level pattern analysis ('multipatt' function within the 'indicpecies' package; de Cáceres and Legendre, 2009) to test for parasite taxa that were significantly associated with white shrimp life stage or locality. Indicator values (IV), i.e., the proportion of parasite x occurring in group A multiplied by the proportion of individuals in group A that contain parasite x (Dufrene and Legendre, 1997), were also calculated to determine how strongly a parasite grouped with a particular locality or host life stage.

Species-specific mixed-effects logistic regression analyses ('glmer' function within the 'lme4' package; Bates *et al.*, 2015), with collection event included as a random effect to account for the lack of independence within sampling events, were used to assess how parasite presence related to white shrimp life stage and collection locality. Model significance was assessed using a likelihood ratio test followed by Tukey's *post hoc* analyses to identify pairwise differences performed using the 'multcomp' package (Hothorn *et al.*, 2008). The relationship between the apostome ciliates and black gill from open water localities in fall months was tested using a binomial test and a mixed-effects segmented regression with collection event included as a random effect. Differences in parasite prevalence and intensity/relative abundance between shrimp with black gill and those without black gill were examined using mixed-effects approaches in the 'lme4' package (Bates *et al.*, 2015) followed by Tukey's comparisons of estimated marginal means in the 'emmeans' package (Lenth, 2022). These models were restricted to collections in open water habitats in fall months and for quantifiable parasites found in the Harbor during these months (i.e., apostomes, rhabditids, lecanicephalideans, trypanorhynch and plagiorchiid). For these models, random factors of individual shrimp, collection month, sampling event, date and sampling event nested within date were included if they enhanced model performance by not contributing to model overfitting.

Results

Parasite community composition

Parasites belonging to 10 genera of 6 classes were found to infect white shrimp in the Charleston Harbor watershed (Table 2;

Fig. 2). Of the 597 white shrimp specimens examined (post-larvae to adult), 82% (491) were infected with at least 1 type of parasite. These parasites included 6 helminths [4 cestodes (1 lecanicephalidean, 1 cyclophyllidean and 2 trypanorhynch), 1 plagiorchiid digenean and 1 rhabditid nematode], 2 ciliates (1 sessilid and 1 apostome), 1 gregarine and 1 microsporidian. All worms were found in a larval stage (i.e., metacercariae for the plagiorchiid, plerocerci for trypanorhynch and juveniles for the rhabditid). Four of the parasite taxa were able to be identified to species-level using DNA sequencing (Table 2). Sequences from the nematode and 2 of the cestodes were only 96–97% similar to any sequence in GenBank and thus could only be identified to genus level (Table 2). Gregarine specimens could not be amplified or sequenced. Sequences from this study were deposited into GenBank under accession numbers OL456224–OL456226 and OL467311–OL467319 [platyhelminthes ($n=6$), ciliate ($n=1$), nematodes ($n=3$) and microsporidians ($n=2$)]. Parasites will hereafter be referenced as gregarines, apostomes, sessilids, microsporidians, rhabditids, cyclophyllids, lecanicephalideans, trypanorhynch and plagiorchiid (Table 2).

Ontogenetic and spatial variation in parasite communities

Prevalence and mean intensity or relative abundance of parasite infections varied across localities and white shrimp life stages (Table 3). Juvenile shrimp had significantly lower prevalence of infection by the rhabditid, lecanicephalidean and trypanorhynch than adult shrimp (Table 4). Subadult shrimp had significantly higher prevalence of infection by the plagiorchiid and rhabditid than juvenile shrimp, but significantly lower prevalence of infection by the sessilid and lecanicephalidean than adult shrimp




Fig. 2. Parasites of the white shrimp, *Penaeus setiferus* observed in the Charleston Harbor watershed, South Carolina, USA. (A) Excysted metacercaria of the plagiorchiid *Opecoeloides fimbriatus*, scale bar 150 μm . (B) Juvenile of the rhabditid *Hysterothylacium reliquens*, scale bar 200 μm . (C) Microsporidian meiodihaplophasid *Agmasoma penaei*, scale bar 25 μm . (D) Sessilid *Zoothamnium* sp., scale bar 75 μm . (E) Cyclophyllidean *Parorchites zederi*, scale bar 85 μm . (F) Lecanicephalidean *Polypocephalus* sp., scale bar 75 μm . (G) Scolex of plerocercus of the trypanorhynch *Prochristianella* sp., scale bar 100 μm . (H) Gregarine gametocysts, scale bar 100 μm . (I) Apostome *Hyalophysa lynni*, scale bar 40 μm .

Table 3. Prevalence (%)/mean intensity or relative abundance (\pm s.e.) of black gill and parasite infections by shrimp life stage and by locality

Parasite and black gill	Shrimp life stage				Locality		
	Post-larvae	Juveniles	Subadults	Adults	Wando River	Ashley River	Harbor
<i>N</i>	54	428	47	68	181	201	161
Gregarine	0/NA	59/NA	83/NA	82/NA	56/NA	57/NA	81/NA
Apostome ^a	0.0	43/0.6 (0.06)	64/2.8 (0.8)	59/1.0 (0.2)	34/0.4 (0.1)	45/0.4 (0.1)	62/1.8 (0.23)
Sessilid ^a	0.0	32/51.7 (2.6)	4/45.8 (7.9)	0/NA	31/47.8 (3.8)	42/54.6 (3.7)	1/16.7 (1.3)
Microsporidian	0/NA	<1/NA	0/NA	0/NA	0/NA	0/NA	1/NA
Rhabditid	0.0	3/1.6 (0.08)	43/2.3 (0.4)	69/2.5 (0.3)	0/NA	1/1.00 (0.1)	50/2.3 (0.12)
Cyclophyllidean	NA	0/NA	4/NA	3/NA	0/NA	0/NA	3/NA
Lecanicephalidean	4/0.67	2/1.1 (0.05)	4/1.0 (0.1)	25/2.4 (0.5)	2/1.00 (0.1)	2/1.3 (0.1)	13/2.1 (0.3)
Trypanorhynch	0.0	47/5.6 (0.5)	83/14.9 (2.4)	90/13.1 (2.0)	36/1.8 (0.1)	49/6.1 (0.7)	86/12.9 (1.2)
Plagiorchiid	2/1	58/7.7 (0.9)	98/23.5 (3.4)	100/22.2 (2.5)	57/3.8 (0.3)	52/4.0 (0.5)	96/23.9 (2.1)
Black gill	0/NA	8/NA	51/NA	41/NA	1/NA	5/NA	50/NA

^aSee *Materials and methods: shrimp gill examination and ciliate abundance* for differences in abundance calculations for these 2 ciliate parasites.

(Table 4). White shrimp collected at the open water Charleston Harbor localities had a higher prevalence of infection by the rhabditid, lecanicephalidean, trypanorhynch and plagiorchiid, but lower prevalence of infection by the sessilid than those collected in the Ashley River and Wando River watersheds (the rhabditid was not collected from the Wando River; Tables 3 and 4). Parasite prevalence did not differ significantly between the Ashley River and Wando River watersheds (Tables 3 and 4).

Parasite community structure varied significantly among both white shrimp life stages (PERMANOVA, $n = 65$, $R^2 = 0.24$, $P = 0.001$) and localities (PERMANOVA, $n = 52$, $R^2 = 0.31$, $P = 0.001$). The parasite communities of juveniles were significantly different from those of both adults and subadults ($P = 0.002$), which were not significantly different from each other ($P = 0.233$). Parasite community dissimilarities were also significantly correlated with black gill score ($R^2 = 0.29$, $P = 0.002$), shrimp length ($R^2 = 0.41$, $P = 0.001$), salinity ($R^2 = 0.16$, $P = 0.022$), but not water temperature ($R^2 = 0.11$, $P = 0.068$; Fig. 3). The apostome and the black gill score vectors were both located at higher y -axis values, but trypanorhynch and plagiorchiids were located at lower y -axis values, highlighting that these factors were associated with different communities of parasites (Fig. 3).

Significant indicators of white shrimp collected at open water Charleston Harbor localities included the rhabditid (IV = 0.998,

$P = 0.001$), plagiorchiid (IV = 0.907, $P = 0.001$), apostome (IV = 0.868, $P = 0.001$), trypanorhynch (IV = 0.855, $P = 0.001$) and lecanicephalidean (IV = 0.772, $P = 0.001$), while the lecanicephalidean (IV = 0.852, $P = 0.005$) and rhabditid (IV = 0.799, $P = 0.005$) were significant indicators of adult white shrimp. The sessilid was a significant indicator of juvenile white shrimp (IV = 0.355; $P = 0.005$) and of shrimp collected in the Ashley River watershed (IV = 0.767; $P = 0.001$).

Parasite communities and black gill

No post-larval white shrimp exhibited macroscopic evidence of black gill, which was almost exclusively found in specimens collected at open water Charleston Harbor localities, regardless of shrimp life stage. Black gill score was a significant vector associated with parasite community structure (Fig. 3). For shrimp collected in open water habitats in the fall, parasite communities between shrimp with and without black were significantly different (PERMANOVA, $n = 9$, $R^2 = 0.31$, $P = 0.008$). Indicator species analyses showed that only the apostome was a significant indicator of shrimp with black gill (IV = 0.93; $P = 0.005$). The apostome was observed across 3 localities (Table 3). Juvenile, subadult and adult white shrimp were all observed with both black gill and apostome infections (Table 3), but significantly more adult

Table 4. Pairwise P values from Tukey's *post hoc* tests of logistic regression models that examined parasite presence across shrimp life stages and localities ($n = 532$). Shrimp life stage and locality were significant factors ($P < 0.01$) in their respective models based on likelihood ratio tests unless indicated by 'n.s.' for not significant, in which case pairwise analyses were not conducted.

Parasite	Shrimp life stage			Locality		
	Juveniles vs subadults	Juveniles vs adults	Subadults vs adults	Wando River vs Ashley River	Wando River vs Harbor	Ashley River vs Harbor
Gregarine	0.173	0.235	0.939	0.995	0.102	0.113
Apostome	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Sessilid	<0.001	<0.001	<0.001	0.210	<0.001	<0.001
Meiodihaplophasid	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Rhabditid	0.011	<0.001	0.113	1	1	<0.001
Lecanicephalidean	0.724	<0.001	0.038	0.995	0.043	0.034
Trypanorhynch	0.145	0.008	0.574	0.138	<0.001	<0.001
Plagiorchiid	0.012	0.999	0.999	0.559	<0.001	<0.001

Significance at $P < 0.05$ of pairwise tests is indicated in bold font.




Fig. 3. Non-metric multi-dimensional scaling (NMDS) plot of white shrimp infected with parasites in the greater Charleston Harbor watershed, South Carolina, USA (stress = 0.11). Data are presented by collection localities: Ashley River (filled black circles), Wando River (open circles) and Charleston Harbor (filled grey circles). Vectors represent significant ($P \leq 0.01$) factors related to community structure; taxa names (e.g. Sessilid) highlight locations in ordinal space most associated with those taxa. Dashed ovals represent data ellipses for parasite communities from each locality.

white shrimp were infected with the apostome ciliate than exhibited black gill (binomial test $P < 0.001$). A positive slope existed between black gill scores ≥ 6 (modeled breakpoint = 5.68 and the relative abundance of the apostome on the gills of white shrimp collected during the black gill season (slope = 0.83, confidence interval = 0.35 – 1.3), but there was no significant relationship between these 2 variables for scores of ≤ 5 (slope = 0.025, confidence interval = -0.47 – 0.35, Fig. 4).

Generalized mixed-effects models on prevalence showed parasite ($\chi^2 = 275.9$, $P < 0.001$) and black gill ($\chi^2 = 24.8$, $P < 0.001$) as significant factors, with individual shrimp (s.d. = 0.341) and collection month (s.d. = 0.211) included as random effects to account for sampling design. *Post hoc* analysis showed prevalence of infection by the trypanorhynchs, and with marginal significance of the rhabditid, was higher in shrimp without black gill than those with this condition (Table 5). The opposite occurred for the apostome, for which prevalence was higher in shrimp with black gill (Table 5; Fig. 5A). Generalized mixed-effects models on intensity showed parasite ($\chi^2 = 20.8$, $P = 0.008$) and black gill ($\chi^2 = 22.1$, $P < 0.001$) as significant factors, with individual shrimp (s.d. = 0.212) and collection month (s.d. = 0.001) included as random effects to account for sampling design. *Post hoc* analysis showed

mean intensities of the rhabditid, and with marginal significance of the trypanorhynchs, were higher in shrimp without black gill than in those with this condition (Table 5; Fig. 5B). This pattern was again reversed for the apostome (Table 5; Fig. 5B). There was no significant association between the presence or intensity of the lecanicephalidean or plagiorchiid and black gill status.

Discussion

The analyses presented here demonstrate differences in the parasite communities of white shrimp that were associated with localities (i.e., tidal creek vs open water habitats), host life stages and the occurrence of black gill. Juvenile white shrimp generally live in tidal creeks until they reach a certain size or level of maturity at which time they egress towards more saline waters in the summer and fall (Lindner and Cook, 1967). While the habitat and life stage of white shrimp were both significantly related to the composition of the parasite community in this study, the confounding nature of these variables makes it difficult to tease apart the individual effects of these factors. Despite this limitation, this study allowed for a deeper understanding of the ontogenetic and spatial dynamics of white shrimp parasite community composition, as




Fig. 4. Black gill score related to apostome ciliate *Hyalophysa lynni* abundance for individual white shrimp (filled grey circles) and mean abundances (filled black circles \pm s.e.; $n = 306$). Segmented regression showed a significant relationship between black gill score (when ≥ 6) and apostome abundance on individual shrimp.

Table 5. Marginal means pairwise contrasts from mixed-effects models of parasite prevalence or intensity in shrimp with and without black gill

Parasite	Prevalence		Intensity	
	Estimate	P value	Estimate	P value
Apostome	-1.767	0.001	-0.618	0.001
Rhabditid	0.738	0.057	0.416	0.027
Lecanicephalidean	-0.087	0.870	0.152	0.418
Trypanorhynch	1.443	0.031	0.350	0.062
Plagiorchiid	15.055	0.992	0.224	0.233

Significance at $P < 0.05$ is indicated with bold font.




Fig. 5. Infection metrics of parasites in white shrimp with black gill (black) and without black gill (white). Prevalence presented as (A) the percentage of shrimp infected and (B) mean intensity (\pm s.e.) and mean relative abundance (\pm s.e.) (for the apostome). Asterisks indicate significant differences between groups for prevalence (logistic regression, Table 5) and mean intensity or mean relative abundance (general linear mixed-effects model, Table 5).

well as the potential physiological impacts of parasites on this economically and ecologically important shrimp species.

To our knowledge, the only previous helminth survey of white shrimp in the Georgia Bight was limited to a total of 29 specimens and reported infection by ‘cestodes, trematodes and nematodes’ (Tripp and Turner, 1983). All parasites encountered herein have previously been reported from penaeid shrimp, including white shrimp in the Gulf of Mexico (Overstreet, 1973; Couch, 1978; Deardorff and Overstreet, 1981; Fontaine, 1985; Carreon *et al.*, 2018), with the exception of the apostome *H. lynni*, which was only recently described from white shrimp from the South Atlantic Bight (Landers *et al.*, 2020). An unidentified apostome ciliate on shrimp gills was associated with black gill in the Gulf of Mexico, albeit prior to the description of *H. lynni* (Overstreet, 1973; Couch, 1978).

In host–parasite systems, the diversity of parasites often changes ontogenetically with habitat and diet (Muñoz and Zamora, 2011; Münster *et al.*, 2015). As such, higher prevalence and mean intensities of infection by the rhabditid, trypanorhynch and lecanicephalidean in subadult and adult white shrimp are potentially attributable to the nature of the life cycles of these parasites. These 3 parasites infect white shrimp *via* their ingestion of either free-living stages or infected planktonic hosts (Johnson, 1975; Carreon *et al.*, 2018). Thus, as shrimp age, opportunities for infection by these parasites increase and parasites can accumulate in their hosts.

The apostome ciliate was not found in post-larval white shrimp but it did infect all other life stages and was found across localities. Frischer *et al.* (2017) found correlations between the presence of this ciliate and black gill suggesting a linear response of the degree of gill melanization to the level of apostome infection. Results herein confirm such a relationship; however, it does not appear linear according to our analyses as many individuals infected with this ciliate did not exhibit black gill. Our data also indicate that a *H. lynni* abundance threshold may exist whereby only high abundances are associated with gill melanization (i.e., black gill). A delay between infection and melanization of the gill tissue, or the host response eliminating suitable parasite habitat leading to no infection at extreme levels of melanization (*sensu* Frischer *et al.*, 2017) could also contribute to the non-linearity of this relationship.

Black gill is indicative of an immune response (Lightner and Redman, 1977; Martin *et al.*, 2000; Cerenius *et al.*, 2010) that can generate suboptimal living conditions for endoparasites (Burnett and Burnett, 2015) given its negative impact on the physiology of the hosts. Hence, activated immune response in shrimp with black gill could lead to the observed reduced prevalence and/or intensities of trypanorhynch and rhabditids in white shrimp with black gill compared to those without black gill. Alternatively, given the role of the hepatopancreas in the crustacean immune system (Röszer, 2014; Cao *et al.*, 2021), damage caused by trypanorhynchs, which embed deeply in this organ and alter its function in portunid crabs (Gurney *et al.*, 2006), could lead to immunosuppression. This could explain significantly higher prevalence of infection of white shrimp without black gill and immunosuppression could facilitate infection by other parasites, including the rhabditid that was also found in higher intensities in shrimp without black gill. Thus, while the apostome ciliate *H. lynni* remains a significant factor associated with black gill outbreaks and the degree of gill melanization in individual white shrimp, other parasites such as the trypanorhynch and rhabditids may also influence the patterns of black gill. To further support the role of parasitic co-infection in black gill occurrence, analyses of fisheries-independent data show no significant relationship between black gill prevalence and white shrimp abundance (Kendrick *et al.*, 2021). Environmental factors, such as salinity and temperature, also have the potential to influence both patterns of black gill (Fowler *et al.*, 2018; Swinford and Anderson, 2021) as well as parasite communities (Koprivnikar *et al.*, 2014; Strepparava *et al.*, 2018), such that future studies should investigate the potential interrelatedness of biotic and abiotic factors in determining white shrimp health metrics.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182022001597>

Data availability. The data that support the findings of this study are available from the corresponding author, MRK, upon reasonable request.

Acknowledgements. The authors thank Dr Robert D. Podolsky for his comments on an earlier draft of this manuscript, as well as Stephen Czwartacki, Kristin Hamilton, Elizabeth Gooding and Elizabeth Underwood for their field assistance. The authors also thank Pam Corwin (SCDNR) for providing the shrimp artwork for the graphical abstract. This publication represents SCDNR Marine Resources Research Institute Contribution Number 860.

Author's contributions. S. R. Z. collected the data and performed the analyses. I. d. B. co-conceived and designed the analysis, collected the data and contributed to analyses. P. R. K.-S. contributed to the design of the analyses. K. M. H.-S. collected the data and contributed to data analysis. N. F. assisted with the processing of specimens in the laboratory. M. R. K. co-conceived and designed the analysis, collected the data, contributed to, and performed, the analyses. All authors contributed to the writing and editing of the article.

Financial support. This study was supported by the South Carolina Saltwater Recreational Fishing License Fund and Award# NA14OAR4170088 from South Carolina Sea Grant Consortium.

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Bates D, Maechler M, Bolker B and Walker S (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1–48.
- Budischak SA, Sakamoto K, Megow LC, Cummings KR, Urban Jr JF and Ezenwa VO (2015) Resource limitation alters the consequences of co-infection for both hosts and parasites. *International Journal of Parasitology* **45**, 455–463.

- Burnett KG and Burnett LE** (2015) Respiratory and metabolic impacts of crustacean immunity: are there implications for the insects? *Integrative and Comparative Biology* **55**, 856–868.
- Bush AO, Lafferty K, Lotz J and Shostak AW** (1997) Parasitology meets ecology on its own terms: Margolis, et al. Revisited. *Journal of Parasitology* **83**, 575–583.
- Byers JE** (2021) Marine parasites and disease in the era of global climate change. *Annual Review of Marine Science* **13**, 397–420.
- Cao X-T, Pan X-Y, Sun M, Liu Y and Lan J-F** (2021) Hepatopancreas-specific lectin participates in the antibacterial immune response by regulating the expression of antibacterial proteins. *Frontiers in Immunology* **12**, 6796767.
- Carreon N, Faulkes Z and Fredensborg BL** (2018) *Polypocephalus* sp. infects the nervous system and increases activity of commercially harvested white shrimp (*Litopenaeus setiferus*). *Journal of Parasitology* **97**, 755–759.
- Cerenius L, Kawabata SI, Lee BL, Nonaka M and Soderhall K** (2010) Proteolytic cascades and their involvement in invertebrate immunity. *Trends in Biochemical Science* **35**, 575–583.
- Clay PA and Rudolf VH** (2019) How parasite interaction strategies alter virulence evolution in multi-parasite communities. *Evolution* **73**, 2189–2203.
- Couch JA** (1978) Diseases, parasites, and toxic responses of commercial penaeid shrimps of the Gulf of Mexico and South Atlantic coasts of North America. *Fishery Bulletin* **76**, 1–44.
- Deardorff TL and Overstreet RM** (1981) Larval *Hysterothylacium* (= *Thynnascaris*) (Nematoda: Anisakidae) from fishes and invertebrates in the Gulf of Mexico. *Proceedings of the Helminthological Society of Washington* **48**, 113–126.
- De Cáceres M and Legendre P** (2009) Associations between species and groups of sites: indices and statistical inference. *Ecology* **90**, 3566–3574.
- DeLancey LB, Jenkins JE and Whitaker JD** (1994) Results of long-term, seasonal sampling for *Penaeus* postlarvae at Breach Inlet, South Carolina. *Fishery Bulletin* **92**, 633–640.
- Del Rio-Rodríguez RE, Pech D, Soto-Rodríguez SA, Gomez-Solano MI and Sosa-Lopez A** (2013) A ten-month diseases survey on wild *Litopenaeus setiferus* (Decapoda: Penaeidae) from southern Gulf of Mexico. *Revista de Biología Tropical* **61**, 1175–1188.
- Diakin A, Paskerova GG, Simdyanov TG, Aleoshin VV and Valigurová A** (2016) Morphology and molecular phylogeny of coelomic gregarines (Apicomplexa) with different types of motility: *Urospora ovalis* and *U. trivisiae* from the polychaete *Travisia forbesii*. *Protist* **167**, 279–301.
- Domínguez-Machín ME, Hernández-Vergara MP, Jiménez-García I, Simá-Álvarez R and Rodríguez-Canul R** (2011) Survey of protozoan, helminth and viral infections in shrimp *Litopenaeus setiferus* and prawn *Macrobrachium acanthurus* native to the Jamapa River region, Mexico. *Diseases of Aquatic Organisms* **96**, 97–103.
- Dufrene M and Legendre P** (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* **67**, 345–367.
- Fontaine CT** (1985) A survey of potential disease-causing organisms in bait shrimp from West Galveston Bay, Texas. NOAA Technical Memorandum NMFS-SEC **169**, 1–63.
- Food and Agriculture Organization of the United Nations (FAO)** (2018) *The State of World Fisheries and Aquaculture 2018 – Meeting the Sustainable Development Goals*. Rome: License, CC BY-NC-SA 3.0 IGO.
- Fowler AE, Leffler JW, Johnson SP, DeLancey LB and Sanger DM** (2018) Relationships between meteorological and water quality variables and fisheries-independent white shrimp (*Litopenaeus setiferus*) catch in the ACE basin NERR, South Carolina. *Estuaries and Coasts* **41**, 79–88.
- Frischer ME, Lee R, Price AR, Walters TL, Bassette MA, Verdiyev R, Torris MC, Bulski K, Geer PJ, Powell SA, Walker AN and Landers SC** (2017) Causes, diagnostics, and distribution of an ongoing penaeid shrimp black gill epidemic in the U.S. South Atlantic Bight. *Journal of Shellfish Research* **36**, 487–500.
- Frischer ME, Landers SC, Walker AN, Powell SA and Lee RF** (2022) Black gill in marine decapod crustaceans: a review. *Reviews in Fisheries Science & Aquaculture* **30**, 1–22.
- Fujiwara M, Zhou C, Acres C and Martinez-Andrade F** (2016) Interaction between penaeid shrimp and fish populations in the Gulf of Mexico: importance of shrimp as forage species. *PLoS ONE*. doi: doi.org/10.1371/journal.pone.0166479
- Ghate H** (1984) Gill melanization and heavy metals in freshwater prawns. *Indian Journal of Fisheries* **31**, 389–393.
- Ghosh K and Weiss LM** (2009) Molecular diagnostic tests for Microsporidia. *Interdisciplinary Perspectives on Infectious Diseases*. doi: doi.org/10.1155/2009/926521
- Gillet R** (2008) Global study of shrimp fisheries. Part II. Shrimp fisheries in selected countries. Shrimp fishing in the United States of America. FAO Fisheries Technical Paper 475, pp. 289–312. FAO Fiji.
- Gleichsner AM, Reinhart K and Minchella DJ** (2018) Of mice and worms: are co-infections with unrelated parasite strains more damaging to definitive hosts? *International Journal of Parasitology* **48**, 881–885.
- Graham AL** (2008) Ecological rules governing helminth-microparasite coinfection. *Proceedings of the National Academy of Sciences of the USA* **105**, 566–570.
- Guo ZL, Liu S, Hu SM, Li T, Huang YS, Liu GX, Zhang H and Lin SJ** (2012) Prevalent ciliate symbiosis on copepods: high genetic diversity and wide distribution detected using small subunit ribosomal RNA gene. *PLoS ONE*. doi: doi.org/10.1371/journal.pone.0044847
- Gurney RH, Johnston DJ and Nowak BF** (2006) The effect of parasitism by trypanorhynch plerocercoids (Cestoda, Trypanorhyncha) on the digestive enzyme activity of *Carcinus maenas* (Linnaeus, 1758) (Decapoda, Portunidae). *Crustaceana* **79**, 663–675.
- Hellard E, Fouchet D, Vavre F and Pontier D** (2015) Parasite-parasite interactions in the wild: how to detect them? *Trends in Parasitology* **31**, 640–652.
- Hervé M** (2021) RVAideMemoire: testing and plotting procedures for biostatistics. R package version 0.9–79. Available at <https://CRAN.R-project.org/package=RVAideMemoire>.
- Hollebone AL** (2006) *An invasive crab in the South Atlantic Bight: friend or foe?* (Thesis), 114 pp. Georgia Institute of Technology.
- Hothorn T, Bretz F and Westfall P** (2008) Simultaneous inference in general parametric models. *Biometrical Journal* **50**, 346–363.
- Jensen K and Bullard SA** (2010) Characterization of a diversity of tetraphylloidean and rhinebothriidean cestode larval types, with comments on host associations and life-cycles. *International Journal of Parasitology* **40**, 889–910.
- Johnson SK** (1975) Handbook of shrimp diseases. *Aquaculture* **601**, 1–27.
- Johnson WS and Allen DM** (2005) *Zooplankton of the Atlantic and Gulf Coasts: A Guide to Their Identification and Ecology*. Baltimore, MD: Johns Hopkins University Press.
- Johnson NG, Burnett LE and Burnett KG** (2011) Characterization of bacteria that trigger hemocytopenia in the Atlantic blue crab, *Callinectes sapidus*. *Biological Bulletin* **221**, 164–175.
- Karthikeyan V, Selvakumar A and Gopalakrishnan A** (2015) A novel report of fungal pathogen *Aspergillus awamori* causing black gill infection on *Litopenaeus vannamei* (Pacific white shrimp). *Aquaculture* **444**, 36–40.
- Kendrick MR, Brunson JF, Frischer ME and Kingsley-Smith PR** (2021) Climate indices predict black gill prevalence in white shrimp *Penaeus setiferus* (Linnaeus, 1767) in South Carolina and Georgia, USA. *Journal of Shellfish Research* **40**, 145–151.
- Koprivnikar J, Ellis D, Shim KC and Forbes MR** (2014) Effects of temperature and salinity on emergence of *Gynaecotyla adunca* cercariae from the intertidal gastropod *Ilyanassa obsoleta*. *Journal of Parasitology* **100**, 242–245.
- Lafferty KD** (2017) Marine infectious disease ecology. *Annual Review of Ecology, Evolution, and Systematics* **48**, 473–496.
- Landers SC, Lee RF, Walters TL, Walker AN, Powell SA, Patel M and Frischer ME** (2020) *Hyalophysa lynni* n. sp. (Ciliophora, Apostomatida), a new pathogenic ciliate and causative agent of shrimp black gill in penaeid shrimp. *European Journal of Protistology* **73**, 1–10.
- Leander BS, Clopton RE and Keeling PJ** (2003) Phylogeny of gregarines (Apicomplexa) as inferred from small-subunit rDNA and β -tubulin. *International Journal of Systematic and Evolutionary Microbiology* **53**, 345–354.
- Lefèvre T, Lebarbenchon C, Gauthier-Clerc M, Missé D, Poulin R and Thomas F** (2008) The ecological significance of manipulative parasites. *Trends in Ecology and Evolution* **24**, 41–48.
- Lenth RV** (2022) emmeans: Estimated marginal means, aka least-squares means. R package version 1.7.2. Available at <https://CRAN.R-project.org/package=emmeans>.
- Lightner DV and Redman R** (1977) Histochemical demonstration of melanin in cellular inflammatory processes of penaeid shrimp. *Journal of Invertebrate Pathology* **30**, 298–302.
- Lindner MJ and Cook HL** (1967) Synopsis of biological data on the white shrimp *Penaeus setiferus*. Food and Agriculture Organization of the United Nations (FAO) Fisheries Reports **57**, 1439–1469.
- Marcogliese DJ** (2004) Parasites: small players with crucial roles in the ecological theater. *EcoHealth* **1**, 151–164.

- Martin GG, Quintero M, Quigley M and Khosrovian H** (2000) Elimination of sequestered material from the gills of decapod crustaceans. *Journal of Crustacean Biology* **20**, 209–217.
- Muncy RJ** (1984) Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico). *U.S. Fish and Wildlife Service Biological Report* 82/11.20.
- Muñoz G and Zamora L** (2011) Ontogenetic variation in parasite infracommunities of the clingfish *Sicyases sanguineus* (Pisces: Gobiesocidae). *Journal of Parasitology* **97**, 14–19.
- Münster J, Klimpel S, Fock HO, MacKenzie K and Kuhn T** (2015) Parasites as biological tags to track an ontogenetic shift in the feeding behavior of *Gadus morhua* off West and East Greenland. *Parasitology Research* **114**, 2723–2733.
- Nadler SA and Hudspeth DSS** (2000) Phylogeny of the Ascaridoidea (Nematoda: Ascaridida) based on three genes and morphology: hypotheses of structural and sequence evolution. *Journal of Parasitology* **86**, 380–393.
- National Marine Fisheries Service (NMFS)** (2020) Fisheries of the United States, 2018. U.S. Department of Commerce, NOAA Current Fishery Statistics No. 2018 Available at <https://www.fisheries.noaa.gov/national/commercial-fishing/fisheries-united-states-2018>.
- Oksanen J, Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens M, Henry H, Szoecs E and Wagner H** (2019) Vegan: community ecology package. R package version 2.5–5. Available at <https://CRAN.R-project.org/package=vegan>.
- Overstreet RM** (1973) Parasites of some penaeid shrimps with emphasis on reared hosts. *Aquaculture* **2**, 105–140.
- Overstreet RM** (1978) *Marine Maladies? Worms, Germs, and Other Symbionts From the Northern Gulf of Mexico*. Ocean Springs, Mississippi: Blossman Printing.
- Palm HW** (2004) *The Trypanorhyncha Diesing, 1863*. Bogor, Indonesia: PKSP-IPB Press.
- Pedersen AB and Fenton A** (2007) Emphasizing the ecology in parasite community ecology. *Trends in Ecology and Evolution* **22**, 133–139.
- R Core Team** (2019) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Reckardt K and Kerth G** (2009) Does the mode of transmission between hosts affect the host choice strategies of parasites? Implications from a field study on bat fly and wing mite infestation of Bechstein's bats. *Oikos* **118**, 183–190.
- Rodríguez ML, Terrazas I, Márquez R and Bojalil R** (1999) Susceptibility to *Trypanosoma cruzi* is modified by a previous non-related infection. *Parasite Immunology* **21**, 177–185.
- Röszer T** (2014) The invertebrate midintestinal gland ('hepatopancreas') is an evolutionary forerunner in the integration of immunity and metabolism. *Cell Tissue Research* **358**, 685–695.
- Rueckart S and Leander BS** (2009) Molecular phylogeny and surface morphology of marine archigregarines (Apicomplexa), *Selenidium* spp., *Filipodium phascosomae* n. sp., and *Platyproteum* n. g. and comb. from North-Eastern Pacific peanut worms (Sipuncula). *Journal of Eukaryotic Microbiology* **56**, 428–439.
- Scharf FS and Schlicht KK** (2000) Feeding habits of red drum (*Sciaenops ocellatus*) in Galveston Bay, Texas: seasonal diet variation and predator-prey size relationships. *Estuaries* **23**, 128–139.
- Shearman RK and Lentz SJ** (2010) Long-term sea surface temperature variability along the U.S. east coast. *Journal of Physical Oceanography* **40**, 1004–1017.
- Sokolova Y, Pelin A, Hawke J and Corradi N** (2015) Morphology and phylogeny of *Agmasoma penaei* (Microsporidia) from the type host, *Litopenaeus setiferus*, and the type locality, Louisiana, USA. *International Journal of Parasitology* **45**, 1–16.
- Strepparava N, Segner H, Ros A, Hartikainen H, Schmidt-Posthaus H and Wahli T** (2018) Temperature-related parasite infection dynamics: the case of proliferative kidney disease of brown trout. *Parasitology* **145**, 281–291.
- Swinford JL and Anderson JD** (2021) Prevalence of black gill (*Hyalophysa lynni*) in white shrimp *Litopenaeus setiferus* and brown shrimp *Farfantepenaeus aztecus* along the Texas Gulf Coast. *Marine and Coastal Fisheries* **13**, 263–274.
- Tripp MR and Turner RM** (1983) Helminth infections of some invertebrates of the Georgia Bight. *Journal of Invertebrate Pathology* **41**, 57–67.
- Tuckey TD, Swinford JL, Fabrizio MC, Small HJ and Shields JD** (2021) Penaeid shrimp in Chesapeake Bay: population growth and black gill disease syndrome. *Marine and Coastal Fisheries* **13**, 159–173.
- Vaseeharan B and Ramasamy P** (2003) Abundance of potentially pathogenic micro-organisms in *Penaeus monodon* larvae rearing systems in India. *Microbiological Research* **158**, 299–308.
- Wenner EL, Knott DM, Barans CA, Wilde S, Blanton JO and Amft J** (2005) Key factors influencing transport of white shrimp (*Litopenaeus setiferus*) post-larvae into the Ossabaw Sound system, Georgia, USA. *Fisheries Oceanography* **14**, 175–194.
- Whitaker JD and Kingsley-Smith PR** (2014) Sea science: an information/education series from the Marine Resources Division. Shrimp life cycle. Available at <https://www.dnr.sc.gov/marine/pub/seascience/shrimpcycle.html> (Accessed January 2020).