

MAXIMAL IMMEDIATE EXTENSIONS ARE NOT NECESSARILY MAXIMALLY COMPLETE

HANS HEINRICH BRUNGS and GÜNTER TÖRNER

(Received 4 May 1989)

Communicated by R. Lidl

Abstract

An extension R_1 of a right chain ring R is called immediate if R_1 has the same residue division ring and the same lattice of principal right ideals as R . Properties of such immediate extensions are studied. It is proved that for every R , maximal immediate extensions exist, but that in contrast to the commutative case maximal right chain rings are not necessarily linearly compact.

1980 *Mathematics subject classification* (*Amer. Math. Soc.*) (1985 Revision): primary 16 A 20, 16 A 14; secondary 13 A18.

Keywords and phrases: maximal valuation ring, completion, right chain ring.

1. Introduction

A commutative valuation ring V that has no proper extension with the same group of values and the same residue field is called maximal (Krull [7]). Such valuation rings can also be characterized by the fact that every pseudo convergent sequence in V has a limit in V or that V is linearly compact (Kaplansky [5]), that is, V is maximally complete.

Here, we consider right chain rings R , that is, rings (not necessarily commutative) with identity for which $aR \subseteq bR$ or $bR \subset aR$ hold for any $a, b \in R$, and show that the above-mentioned notions carry over to this case.

After collecting some basic notations and results in Section 2 we define in Section 3 immediate extensions of right chain rings (see Definition 3.1)

The first author was supported in part by NSERC.

© 1990 Australian Mathematical Society 0263-6115/90 \$A2.00 + 0.00

discuss properties of such extensions and show that maximal immediate extensions exist (Theorem 3.6). In Section 4 we discuss pseudo convergent sequences and linear compactness, and show in particular that every element $z \in R_1 \setminus R$ with R_1 an immediate extension of R , is a limit of a pseudo convergent sequence in R (Theorem 4.12). Finally, in Section 5, we use some of the earlier results to show by an example that there exist right chain rings R without proper immediate extensions, that is, maximal rings R , which are not maximally complete, that is, which contain pseudo convergent sequences that do not have a limit in R .

2. Preliminaries

All rings are assumed to be associative, with identity element, but not necessarily commutative. The Jacobson radical of a ring R is denoted by $J(R) = J$ and the group of units of R by $U(R) = U$. A ring is said to be a *right chain ring* if its lattice of right ideals is linearly ordered by inclusion, that is, for each $a, b \in R$ we have $aR \subseteq bR$ or $bR \subset aR$. Left chain rings are defined similarly and a left right chain ring is called a *chain ring*.

A domain is a ring without zero-divisors. Chain domains are the valuation rings studied in Mathiak [8]. A ring is *ring invariant* if all its right ideals are two-sided and *invariant* if it is right and left invariant. The valuation rings in Schilling [10] are exactly the invariant chain domains; these are the rings corresponding to valuations for skew fields into ordered groups.

A two-sided ideal P of a ring R is called *prime* if $aRb \subseteq P$ implies $a \in P$ or $b \in P$ in P , for $a, b \in R$, and *completely prime (c.prime)* if this implication follows from $ab \in P$.

In case Q is a prime but not c.prime ideal of a right chain ring R , a c.prime ideal P exists in R with $P \supset Q$ such that there is no prime ideal between P and Q and $P^2 = P$ (Bessenrodt et al. [1]). The question whether such prime ideals Q can exist in chain rings is interesting and Dubrovin [4] constructed some examples; however it is difficult to follow his arguments at some places.

3. Immediate extensions

Krull defined immediate extensions of commutative valuation domains in [7] and showed that maximal immediate extensions exist. For right chain rings the following definition generalizes Krull's original definition.

DEFINITION 3.1. Let R be a right chain ring. A ring extension $R_1 \supseteq R$ is called an *immediate extension* of R if the following conditions hold.

- (i) For every element $a_1 \in R_1$ there exist $a \in R, u_1 \in U(R_1)$ with $a_1 = au_1$.
- (ii) $\phi(r + J(R)) = r + J(R_1)$ defines an isomorphism from the skew field $R/J(R)$ onto $R_1/J(R_1)$.

Before we list properties of immediate extensions we make a general observation about extensions of right chain rings and characterize immediate extensions by somewhat different conditions.

LEMMA 3.2. *The following conditions are equivalent for right chain rings $R \subseteq R_1$:*

- (a) $U(R_1) \cap R = U(R)$;
- (b) $J(R_1) \cap R = J(R)$.

PROOF. We observe that $U(R) = R \setminus J(R), U(R) \subseteq U(R_1) \cap R$ and $J(R_1) \cap R \subseteq J(R)$. Assume (a) and it follows that the element $j \in J(R)$ is also in $J(R_1)$, since j would otherwise be in $U(R_1) \cap R = U(R)$. Similarly, under the assumption of (b), an element $v \in U(R_1) \cap R$ cannot be in $J(R)$, since it would then be in $J(R_1)$.

LEMMA 3.3. *Let R be a right chain ring. An extension $R_1 \supseteq R$ is an immediate extension if and only if $U(R_1) \cap R = U(R)$ and for every element $a_1 \in R_1$ there exist $a \in R$ and $j_1 \in J(R_1)$ with $a_1 = a(1 + j_1)$.*

PROOF. We assume that R_1 satisfies the conditions of the lemma and (ii) of Definition 3.1 follows immediately, since $1 + j_1 \in U(R_1)$. To prove (b) define $\phi(a_1) = a + J(R) \in R/J(R)$ if $a_1 = a(1 + j_1) \in R_1$, with $a \in R, j_1 \in J(R_1)$. This mapping defines a homomorphism from R_1 onto $R/J(R)$ with kernel $J(R_1)$ and it follows that the mapping ϕ in (b) of Definition 3.1 exists.

To prove the converse we observe first that an immediate extension of a right chain ring is again a right chain ring. It follows from (b) in Definition 3.1 that $J(R) = R \cap J(R_1)$ and hence $U(R) = U(R_1) \cap R$ by the above lemma. Finally, if $a_1 = a'u_1$ for $a_1 \in R_1, a' \in R$ and $u_1 \in U(R_1)$ then $u_1 = u + j'_1$ for $u \in U(R), j'_1 \in J(R_1)$ and $a_1 = a'u(1 + u^{-1}j'_1)$ shows that the conditions of the lemma are satisfied.

PROPOSITION 3.4. *Let R_1 be an immediate extension of a right chain ring R . Then the following are satisfied.*

- (i) R_1 is a right chain ring.

- (ii) $J(R_1) \cap R = J(R)$.
 (iii) $U(R_1) \cap R = U(R)$.
 (iv) $\chi(aR) = aR_1$ defines a one-to-one mapping from the lattice $\mathscr{W}(R)$ of principal right ideals onto $\mathscr{W}(R_1)$.
 (v) R_2 is an immediate extension of R_1 if and only if R_2 is an immediate extension of R containing R_1 .
 (vi) Let I be a right ideal in R . Then $I_1 = \{ir \mid i \in I, r \in R_1\} = IR_1$ is a right ideal in R_1 and $IR_1 \cap R = I$.
 (vii) Let I_1 be a right ideal in R_1 . Then the contracted right ideal $I = I_1 \cap R$ satisfies $I_1 = IR_1$.
 (viii) The contraction of a two-sided ideal in R_1 is two-sided in R ; the contraction of a c.prime ideal in R_1 is c.prime in R .
 (ix) The contraction $I = I_1 \cap R$ is idempotent in R if I_1 is an idempotent ideal in R_1 .

PROOF. The statements (i)–(iii) follow from Lemma 3.3. That the mapping χ is onto follows from (i) in Definition 3.1 and if $a = bc$ in R and $aR_1 = bR_1$, then $a = bc = bu_1$ for $u_1 \in U(R_1)$. Hence, $b(c - u_1) = 0$ implies $b = 0$ if $c \in J(R) \subseteq J(R_1)$.

(v) For an element $a_2 \in R_2$ we have $a_2 = a_1u_2 = au_1u_2$ for $a_1 \in R_1$, $u_2 \in U(R_2)$, $u_1 \in U(R_1)$, $a \in R$. The skew fields of residues of R and R_2 are also identical.

To prove (vi) let $\sum_{k=1}^n i_k r_k$ be an element in IR_1 with $i_j \in I$, $r_j \in R_1$, and we can assume $i_j = i_1 a_j$ for $a_j \in R$ and all j . We have $\sum i_k r_k = i_1 \sum a_k r_k = i_1 r$ for some $r \in R_1$ which shows that $I_1 = IR_1$ is indeed a right ideal in R_1 . An element in $IR_1 \cap R$ has the form iru_1 for $i \in I$, $r \in R$, $u_1 \in U(R_1)$. If such an element is not contained in I , there exists $q \in J(R)$ with $iru_1 q = i$ and $i = 0$ follows, a contradiction.

(vii) The inclusion $IR_1 \subseteq I_1$ is obvious. Conversely, let $iu \in I_1$ with $i \in R$, $u \in U(R_1)$ and $i \in I$, $iu \in IR_1$ follows.

(viii) is obvious.

(ix) We can assume that I_1 and I are nonzero. Nonzero idempotent ideals are c.prime and it follows that I_1 and hence I are c.prime. If $I^2 \subset I$ there exists an element $p \in I \setminus I^2$ and $p = p_1 u_1 p_2 u_2 \in I_2 = I_1^2$ for $p_1, p_2 \in I$, $u_1, u_2 \in U(R_1)$. We have $u_1 p_2 = p'_2 u'_1$ with $u'_1 \in U(R_1)$, $p'_2 \in R$ and $p'_2 \in I$ follows. Hence, $pR_1 = p_1 p'_2 R_1$ and $pR = p_1 p'_2 R$ by (iv). This implies $p \in I^2$, a contradiction.

The next result is the crucial step in the proof that maximal immediate extensions exist for right chain rings. The original proof of this result (see

Krull [7]) For commutative valuation domains cannot be as easily adapted to our situation as the arguments in Rayner [9].

LEMMA 3.5. *Let R be a right chain ring, $R/J(R) = D$, $\mathscr{W} = \{aR \mid 0 \neq a \in R\}$ and R_1 an immediate extension of R . Then $|R_1| \leq |D|^{\mathscr{W}}$.*

PROOF. Let $\mathscr{C} = \{c_i \mid c_i \in R, i \in T\}$ be a set of representatives of D in R and let W' be a subset of R containing exactly one generator for each nonzero principal right ideal of R , that is $\mathscr{W} = \{aR \mid a \in W'\}$ and $aR \neq a'R$ for $a \neq a'$ in W' . Let $a \in W'$ and aR_1 be the corresponding principal right ideal in R_1 . We have $R_1 = \bigcup_{i \in T} (c_i + J(R_1))$, and $aR_1 = \bigcup_{i \in T} (ac_i + aJ(R_1))$. Observe that $a(c_i - c_j) \neq aJ(R_1)$ since $c_i - c_j \in U(R)$ for $i \neq j$. For aR_1 , considered as a subgroup of the additive group $(R_1, +)$, choose coset representatives $S = \{a_j \mid j \in L\}$, and $R_1 = \bigcup_{j \in L} (a_j + aR_1) = \bigcup_{i \in T, j \in L} (a_j + ac_i + aJ(R_1))$ follows. Let r be any element in R_1 and consider in $(R_1, +)$ the coset of $aJ(R_1)$ that contains r , that is, $r + aJ(R_1) = a_j + ac_i + aJ(R_1)$ for a certain $j \in L, i \in T$. We define a mapping f from R_1 into $\mathscr{C}^{\mathscr{W}}$ by defining $f(r) = \bar{r} \in \mathscr{C}^{\mathscr{W}}$ with $\bar{r}(w) = c_i$ if $w = aR$ in \mathscr{W} . By construction, f is well defined and we show that f is one-to-one.

Let $r \neq t$ be elements in R_1 and $(r - t)R_1 = aR_1$ follows a certain $a \in W'$. We have $r + aR_1 = t + aR_1$, but $r + aJ(R_1) \neq t + aJ(R_1)$, since $r - t = au, u \in U(R_1)$. Hence, $r + aJ(R_1) = a_j + ac_1 + aJ(R_1)$ and $t + aJ(R_1) = a_j + ac_2 + aJ(R_1)$ where $a_j \in S$ and $c_1 \neq c_2 \in \mathscr{C}$. This implies $\bar{r}(w) = c_1 \neq c_2 = \bar{t}(w)$ for $w = aR$ and proves the lemma.

As a corollary we obtain

THEOREM 3.6. *Any right chain ring R has maximal immediate extensions.*

PROOF. We consider (using Lemma 3.5) the set \mathscr{E} of immediate extensions. Let $R \subseteq R_1 \subseteq R_2 \subseteq \dots$ be an ascending chain of immediate extensions of R and $\hat{R} = \bigcup_i R_i$ their union. We must show that \hat{R} is in \mathscr{E} .

If $a, b \in \hat{R}$, then $a, b \in R_i$ some i and $ac = b$ or $a = bc$ for some c in R_i . Hence, \hat{R} is a right chain ring. We have $J(\hat{R}) \cap R \subseteq J(R)$ and if $j \in J(R)$ is not in $J(\hat{R})$, it is a unit in some R_i , a contradiction, since $J(R_i) \cap R = J(R)$ (Lemmas 3.2, 3.3). Finally, every $\hat{a} \in \hat{R}$ is in some R_i and $\hat{a} = a(1 + j_i), j_i \in J(R_i)$. But $J(\hat{R}) = \bigcup_i J(R_i)$, since $U(\hat{R}) = \bigcup_i U(R_i)$ and $\hat{R} \in \mathscr{E}$ follows from Lemma 3.3.

We consider properties of a right chain ring R that are inherited by immediate extensions.

PROPOSITION 3.7. *Let R_1 be an immediate extension of the right chain ring R . Then the following conditions are satisfied.*

(i) IR_1 is a prime ideal in R_1 if I is a prime ideal in R and IR_1 is a two-sided ideal in R_1 .

(ii) R_1 is prime if R is prime.

(iii) Let I be a two-sided ideal in R with $I_1 = IR_1$ a two-sided ideal in R_1 . Then I is c.prime if and only if I_1 is c.prime.

(iv) R is a domain if and only if R_1 is a domain.

(v) R is right noetherian if and only if R_1 is right noetherian.

PROOF. To prove (i) let $x_1 = xu_1$, $y_1 = yv_1$ be elements in R with $x, y \in R$, $u_1, v_1 \in U(R_1)$ and $xu_1R_1yv_1 \subseteq IR_1$. Then $IR_1 \supseteq xu_1R_1y = xR_1y \supseteq xRy$ and $xRy \subseteq IR_1 \cap R = I$ follows (Proposition 3.4(vii)). We have $x \in I$ or $y \in I$ which shows $x_1 \in I_1$ or $y_1 \in I_1$.

(ii) is obvious using (i) and $I = (0)$.

(iii) Suppose I is completely prime and $xu_1yv_1 \in IR_1$ with $x_1 = xu_1$, $y_1 = yv_1 \in R_1$, $x, y \in R$, $u_1, v_1 \in U(R_1)$. If $yv_1 \in IR_1$ we are done, otherwise $u_1y = y'u'_1 \notin IR_1$ for $y' \in R$, $u'_1 \in U(R_1)$. Hence, $y' \in R \setminus I$, but $xu_1yv_1 = xy'u'_1v_1 \in IR_1$ implies $xy' \in I$ and $x \in I$ follows. The converse is obvious and (iv) follows immediately.

To prove (v) we observe that a right chain ring is right noetherian if and only if the maximum condition holds for principal right ideals. By Proposition 3.4(iv) this condition holds for R if and only if it holds for R_1 .

Right noetherian right chain rings are right invariant. We will show by an example that this is no longer true for discrete right chain rings where we use the following definition.

DEFINITION 3.8. A right chain ring R is called *discrete* if R satisfies a.c.c. for c.prime ideals and $P \neq P^2$ for every proper c.prime ideal P .

LEMMA 3.9. *Let R be a discrete right chain ring. Then every prime ideal of R is completely prime and R_P/QR_P is a right noetherian right chain ring with Q the maximal prime ideal properly contained in P . The ring R_P is a right noetherian right chain ring provided P is the minimal prime ideal of R .*

Here, R_P denotes the localization of R at a prime ideal P .

PROOF. To prove the first part of the lemma we recall the result about prime ideals mentioned in Section 2. Since a prime ideal which is not completely prime is paired with a c.prime ideal $P = P^2 \neq (0)$ all prime ideals in a discrete right chain ring are c.prime.

If P is a c.prime ideal in a right chain ring and $S = R \setminus P$, then $N_r(S) = \{a \in R \mid sa = 0 \text{ some } s \in S\}$ and $N_l(S) = \{a \in R \mid as = 0 \text{ some } s \in S\}$ are ideals in R and the set \bar{S} consisting of the images of the elements of S in $R/I = \bar{R}$, $I = N_l(S) \cup N_r(S)$ is a right Ore set in \bar{R} and $\overline{RS^{-1}} = \{\bar{r}\bar{s}^{-1} \mid \bar{r} \in \bar{R}, \bar{s} \in \bar{S}\}$ exists and is denoted by R_p .

If Q is the maximal c.prime ideal in R properly contained in P , then $I \subseteq Q = \bigcap_{n \in \mathbb{N}} P^n$ and QR_p is the maximal c.prime ideal in R_p contained in $J(R_p)$ (as is common for commutative localizations, we write Q for \bar{Q} if there is no ambiguity).

Since PR_p is two-sided, we have $(PR_p)^2 = P^2R_p \neq PR_p$. To prove this last statement choose $m \in P \setminus P^2$ in R and assume $\bar{m} \in P^2R_p$. This means $\bar{m} = \bar{p}_1\bar{p}_2\bar{s}^{-1}$ or $ms = p_1p_2 + n$ in R with $n \in I$ and $n = p_1n_1$ follows. Here, $n_1 \in P$, since $\bar{n} = 0 \neq \bar{p}_1$ shows that $n_1 \in S$ is not possible. If $p_1 = mt$ we obtain $m(s - t(p_2 + n_1)) = 0$ and $\bar{m} = 0$, a contradiction. If $p_1t = m$, then $t \in S$ and $p_1(ts - (p_2 + n_1)) = 0$ and $\bar{p}_1 = 0 = \bar{m}$, again a contradiction.

This shows that $\bar{m} \neq P^2R_p$, $PR_p = mR_p$ is a principal right ideal and $\bigcap (PR_p)^i = \bigcap m^iR_p = QR_p$ and the only right ideals in R_p/QR_p are of the form $m^iR_p/QR_p, (0)$.

If P is the minimal prime ideal of R , it could happen that $I = P$. Then R_p is a division ring. Otherwise we show as before that $PR_p \neq P^2R_p$ and R_p is right noetherian. (We choose $m \in P \setminus (P^2 \cup I)$ for the argument as above.)

A partial converse of the last lemma is

LEMMA 3.10. *Let R be a right chain ring whose prime ideals are c.prime and satisfy the a.c.c. Further, assume that the ring R_p/QR_p is a right noetherian right chain ring for all pairs of neighbouring prime ideals $P \supset Q$. If R is not a domain, assume R_p also to be right noetherian for the minimal prime ideal P of R . Then R is discrete.*

PROOF. By Definition 3.8 it suffices to prove $P^2 \neq P$ for all prime ideals $P \neq (0)$. We consider the situation $P \supset Q \neq (0)$, and denote by $m + Q_p$ the generator of the maximal (right) ideal in the right noetherian ring R_p/QR_p . If we assume $P^2 = P$, we obtain $m = m_1m_2 \in P \setminus Q$ for some $m_1, m_2 \in P$. On the other hand, $(m + Q_p)(rt^{-1} + Q_p) = m_1 + Q_p$ holds for some $rt^{-1} + Q_p$, thus $mrt^{-1} - m_1 = m_1m_2rt^{-1} \in Q_p$. The element $1 - m_2rt^{-1}$ is a unit in the local ring R_p/QR_p , thus $M_1 \in Q_p$, a contradiction and $P^2 \neq P$ follows.

Now, consider the case where P is the minimal completely prime ideal in R and R_P is right noetherian. If the kernel $I = N_l(S) \cup N_r(S)$ is strictly contained in P we use similar arguments as above and we are done.

Therefore, we are left with the case $P = I + N_l(S) \cup N_r(S)$. First let $I = N_l(S)$ and $a \in P$, then $as = 0$ holds for some $s \in S$ which implies $aP = 0$, thus $(0) = P^2 \neq P$.

Now we consider the case $N_r(S) = I$ and assume $P^2 = P$. If the segment $(0) \subset P$ is simple and if X, Y are two-sided ideals with $X, Y \not\subseteq (0)$, we must have $P \subseteq X, Y$, and hence $(0) \subset P = P^2 \subseteq XY$. This shows that (0) is prime, by assumption even c.prime, and hence R a domain. However, this leads to $N_r(S) = (0) \neq P$, a contradiction.

Therefore, it remains to discuss the case where $(0) \subset P$ is not simple and $P^2 = P = N_r(S)$ is assumed. With the arguments as above we know that for each two-sided ideal L with $(0) \subset L \subset P$ the segment $L \subset P$ is not simple. For every $a \in P$ we can find a two-sided ideal L satisfying $aR \subseteq L \subset P$; to prove this, take the union I of all two-sided ideals contained in aR . The union I is again a two-sided ideal, thus, by the previous arguments there exists a two-sided ideal L satisfying $aR \subseteq L \subset P$. The intersection $\bigcap L^n$ of a two-sided ideal which is not nilpotent is c.prime, therefore we conclude $L^n = (0)$ for some $n \in \mathbb{N}$ as P is the minimal prime ideal and finally $a^n = 0$ holds for some $n \in \mathbb{N}$, a arbitrarily chosen in P . Now we want to show that even $P^3 = 0$ which would lead to $P = (0)$, a contradiction. Take $x, y, z \in P$. As $I = N_r(S)$ holds we have $sx = 0$ for some $s \in S$. Let $sr = x$ for some $r \in P$ and $xyz = sryz$ follows. If $ry = yr_1$ holds for a suitable $r_1 \in R$, we are done. Otherwise $ryr_1 = y$ follows for some $r_1 \in R$. If $r_1R \supseteq zR$ then $xyz = sryz = sryr_1r_2 = syr_2 = 0$ with $r_1r_2 = z$. However, $r_1 = zr_2$ leads to $ryzr_2 = y$ and thus $r^n y(zr_2)^n = y$, a contradiction as $zr_2 \in P$ implies $(zr_2)^n = 0$ for a sufficiently large n .

Discrete right chain rings need not to be right invariant as the following example shows.

If one denotes by V_1 and V_2 the two extensions of $\mathbb{Z}_{(5)}$, the localization of \mathbb{Z} at (5) in $\mathbb{Z}[i]$ and with σ conjugation in $\mathbb{Q}(i)$, then

$$R = \left\{ \sum_{k=0}^{\infty} t^k a_k \mid a_k \in \mathbb{Q}(i), a_0 \in V_1 \right\} \subseteq \mathbb{Q}(i)[[t, \sigma]]$$

is a non right invariant discrete valuation ring; here $\mathbb{Q}(i)[[t, \sigma]]$ is the skew power series ring in one variable t with coefficients in $\mathbb{Q}(i)$ and $at = ta^\sigma$ defines the multiplication. If we assume that $V_1 = \mathbb{Z}[i]_{(2-i)}$ then $\frac{1}{2+i}t \notin tR$

since $t^{-1} \frac{1}{2+i} t = \frac{1}{2-i} \notin R$. However, the two prime ideals $P_1 = (2 - i)R = J(R)$ and P_2 generated by $\{t(2 - i)^{-n} | n = 1, 2, \dots\}$ are complete and satisfy $P_i^2 \neq P_i - R$ is a discrete right chain ring that is not right invariant.

LEMMA 3.11. *If R_1 is an immediate extension of a discrete right chain ring R then R_1 is discrete.*

PROOF. R_1 cannot contain a prime ideal Q that is not completely prime, since in that case a c.prime ideal P would exist in R_1 and $P^2 = P$. The intersection $P \cap R$ would be an idempotent c.prime ideal in R (Proposition 3.4(ix))—a contradiction to the assumption. Applying this last argument again, we conclude that all complete prime ideals $\neq (0)$, R in R_1 are not idempotent. If P_1, P_2 are distinct c.prime ideals in R_1 then $R \cap P_2, R \cap P_1$ are distinct c.prime ideals in R and hence a.c. for c.prime ideals follows for R_1 .

We conclude this section with an open problem.

Open problem. If R_1 is an immediate extension of the right chain ring R and I a two-sided ideal of R is then IR_1 a two-sided ideal of R_1 ?

4. Pseudo convergent sequences and linear compactness

The elements in an immediate extension R_1 of a right chain ring R that are not contained in R can be described as limits of pseudo convergent sequences in R . If every such sequence in R has already a limit in R , then R is called maximally complete. Some of the definitions extend to uniserial modules.

Let M_R be a uniserial R -module, R a ring. We define a mapping v from $M_R \setminus \{0\}$ onto the set $\mathscr{M} = \{aR | 0 \neq a \in M\}$ by $v(a) = aR$ and set $v(a) \geq v(b)$ if and only if $aR \subseteq bR$. It follows that $v(a - b) \geq \min\{v(a), v(b)\}$ with equality if $v(a) \neq v(b)$.

The definitions of pseudo convergent sequences and limits as considered by Ostrowski, Kaplansky, Schilling and others extend, as do the basic propositions, to uniserial modules.

DEFINITION 4.1. Let M_R be a uniserial R -module. A sequence $(a_\rho)_{\rho \in \Lambda}$, $a_\rho \in M_R$ and Λ well-ordered with no last element is called *pseudo convergent* (p.c.) if $v(a_\tau - a_\sigma) > v(a_\sigma - a_\rho)$ for $\rho < \sigma < \tau \in \Lambda$.

LEMMA 4.2. *If $(a_\rho)_{\rho \in \Lambda}$ is a p.c. sequence in M_R , then either*

- (i) $v(a_\rho) < v(a_\sigma)$ for $\rho < \sigma$ or
- (ii) $v(a_\rho) = v(a_\sigma)$ for $\rho, \sigma \geq \lambda$ and some ordinal λ .

PROOF. Suppose (i) does not hold, that is $v(a_\rho) \geq v(a_\sigma)$ for some $\rho < \sigma$. Then $v(a_\tau) = v(a_\sigma)$ for all $\tau > \sigma$, since otherwise

$$v(a_\tau - a_\sigma) = \min\{v(a_\tau), v(a_\sigma)\} \leq v(a_\sigma),$$

but $v(a_\sigma - a_\rho) \geq v(a_\sigma)$ —a contradiction

LEMMA 4.3. Let $(a_\rho)_{\rho \in \Lambda}$, be a p.c. sequence in M_R . Then $v(a_\sigma - a_\rho) = v(a_{\rho+1} - a_\rho)$ for all $\sigma > \rho$.

PROOF. We can assume that $\sigma > \rho + 1$ and obtain

$$v(a_\sigma - a_{\rho+1}) > v(a_{\rho+1} - a_\rho)$$

and $a_\sigma - a_\rho = (a_\sigma - a_{\rho+1}) + (a_{\rho+1} - a_\rho)$. It follows that

$$v(a_\sigma - a_\rho) = \min\{v(a_\sigma - a_{\rho+1}), v(a_{\rho+1} - a_\rho)\} = v(a_{\rho+1} - a_\rho).$$

Let $(a_\rho)_{\rho \in \Lambda}$ be a p.c. sequence. We write $v(a_\sigma - a_\rho) = \gamma_\rho$ where $\gamma_\rho \in \mathcal{W}_M$ is a constant for all $\sigma > \rho$ by the above lemma.

DEFINITION 4.4. An element $x \in M_R$ is called a *limit* of the p.c. sequences $(a_\rho)_{\rho \in \Lambda}$, if $v(x - a_\rho) = \gamma_\rho$ for all ρ .

DEFINITION 4.5. The submodule B of M consisting of elements $b \in M_R$ with $v(b) > \gamma_\rho$ for all ρ is called the *breadth* of the p.c. sequence $(a_\rho)_{\rho \in \Lambda}$.

Let $(a_\rho)_{\rho \in \Lambda}$ be a p.c. sequence of type (i) as described in Lemma 4.2. Then an element $b \in M$ is a limit of (a_ρ) if and only b is in the breadth of (a_ρ) : we observe that $v(a_\sigma - a_\rho) = v(a_\rho) = \gamma_\rho$ for $\rho < \sigma$ then $v(b - a_\rho) = \gamma_\rho$ and b is a limit. Conversely, if $v(x - a_\rho) = \gamma_\rho = v(a_\rho)$ then $v(x) \geq \gamma_\rho$ for all ρ and hence $v(x) \geq \gamma_{\rho+1} > \gamma_\rho$, that is, x is in the breadth of (a_ρ) .

With similar arguments one proves the following result.

LEMMA 4.6. Let a be a limit of a p.c. sequence (a_ρ) in M_R . Then $c \in M_R$ is a limit of (a_ρ) in M if and only if $c = a + b$ for $b \in B$, the breadth of (a_ρ) .

DEFINITION 4.7. We say a uniserial right R -module M is *maximally complete* if every p.c. sequence in M has a limit in M .

Let M_R be an uniserial R -module, $I_\alpha \subseteq M_R$, $\alpha \in \Lambda$, a family of submodules of M where Λ is a well-ordered set of indices and $I_\alpha \supset I_\beta$ for $\alpha < \beta$. Further, let $(x_\alpha)_{\alpha \in \Lambda}$ be a sequence of elements in M_R and $\mathcal{F} = (x_\alpha + I_\alpha)_{\alpha \in \Lambda}$ the corresponding sequence of cosets in M .

DEFINITION 4.8. With $M_R, I_\alpha, \Lambda, (x_\alpha)$ as above we have the following definitions.

- (i) \mathcal{F} has the *finite intersection property (FIP)* if $\bigcap_{\alpha \in \mathcal{F}} (x_\alpha + I_\alpha) \neq \emptyset$ for all finite subsets $\mathcal{F} \subseteq \Lambda$.
- (ii) \mathcal{F} is *pairwise solvable* if $(x_\alpha + I_\alpha) \cap (x_\beta + I_\beta) \neq \emptyset$ for all $\alpha, \beta \in \Lambda$; we also say that \mathcal{F} is *compatible*.
- (iii) An element $a \in \bigcap_{\alpha \in \Lambda} (x_\alpha + I_\alpha)$ is called a *solution* of $x \equiv x_\alpha \pmod{I_\alpha}$ for all $\alpha \in \Lambda$; we also say that a is a solution for \mathcal{F} .
- (iv) The intersection $\bigcap I_\alpha = B$ is called the *breadth* of \mathcal{F} .
- (v) Let B be a submodule of M_R . Then M_R is called *linearly B-compact* if all sequences \mathcal{F} with breadth B have a solution.
- (vi) M is called (almost) *linearly compact* if M is linearly B compact for all submodules $B \subseteq M (B \neq (0))$.
- (vii) A ring is called (almost) *right linearly compact* if R_R is (almost) right linearly compact.

REMARK. If B is a submodule of the uniserial module M_R which has an immediate upper neighbour $B' \supset B$ in the lattice of submodules of M then any sequence $\mathcal{F} = (x_\alpha + I_\alpha)_{\alpha \in \Lambda}$, with breadth $\mathcal{F} = \bigcap I_\alpha = B$, satisfies $I_\alpha = B$ for some α , that is, Λ has a last element.

LEMMA 4.9. Let $M_R, \Lambda, (x_\alpha), (I_\alpha)_{\alpha \in \Lambda}$, and \mathcal{F} be as above. Then the following conditions are satisfied.

- (i) The sequence \mathcal{F} is pairwise solvable if and only if $x_\alpha - x_\beta \in I_\alpha \cup I_\beta$ for all $\alpha, \beta \in \Lambda$.
- (ii) \mathcal{F} is pairwise solvable if and only if \mathcal{F} has FIP.
- (iii) The sequence of cosets $(x_\alpha + I_\alpha)_{\alpha \in \Lambda}$ is linearly ordered by inclusion if \mathcal{F} is pairwise solvable.
- (iv) If all compatible sequences \mathcal{F} with breadth I and cyclic R -modules I_α are solvable, then all compatible sequences \mathcal{F} with breadth I are solvable.
- (v) If $x, x' \in M_R$ are solutions for \mathcal{F} , then $x - x' \in I$ where I is the breadth of \mathcal{F} .

PROOF. (i) If T is pairwise solvable there exists an element $x \in M$ with $x - x_\alpha \in I_\alpha, x - x_\beta \in I_\beta$ and $x_\alpha - x_\beta \in I_\alpha \cup I_\beta = I_\alpha$ for $\alpha \leq \beta$. On the other hand, if $x_\alpha - x_\beta \in I_\alpha \supseteq I_\beta$ then $x_\beta - x_\alpha \in I_\alpha$ and $x_\beta \in (x_\beta + I_\beta) \cap (x_\alpha + I_\alpha)$.

(ii) Let $I_{\alpha_1} \supset I_{\alpha_2} \supset \dots \supset I_{\alpha_n}$ for $\alpha_1 < \alpha_2 < \dots < \alpha_n$. Then $x_{\alpha_n} - x_{\alpha_i} \in I_{\alpha_i}$ for $i = 1, \dots, n$, if we assume pairwise solvability.

(iii) Suppose $x_\alpha + I_\alpha, x_\beta + I_\beta$ are given with $I_\alpha \supset I_\beta$ and $x_\beta - x_\alpha \in I_\alpha$. Then $x_\beta + I_\beta \supseteq x_\alpha + I_\alpha$ follows immediately

(iv) Let $\mathcal{F} = (x_\alpha + I_\alpha)_{\alpha \in \Lambda}$, be a compatible sequence. The statement is trivial if Λ has a last element. Otherwise consider $x_\alpha - x_{\alpha+1} = a'_\alpha \in I_\alpha$ for every $\alpha \in \Lambda$, and choose $a_\alpha \in M$ with $I_{\alpha+1} \cup a'_\alpha R \supseteq a_\alpha R \supseteq I_\alpha$ and $y_\alpha = x_{\alpha+1}$. Then $\bigcap a_\alpha R = \bigcap I_\alpha$ and $\bigcap (y_\alpha + a_\alpha R) = \bigcap (x_\alpha + I_\alpha)$ where $\mathcal{F}' = (y_\alpha + a_\alpha R)$ is a compatible sequence.

LEMMA 4.10. *Let M_R be a uniserial R -module which is linearly I -compact. Then the R -module M/N is linearly I/N -compact for every submodule N of M with $N \subseteq I$.*

PROOF. Let $(x_\alpha + N_\alpha/N)_{\alpha \in \Lambda}$ be a compatible sequence. Then, obviously $(x_\alpha + N_\alpha)_{\alpha \in \Lambda}$ is compatible as $N \subseteq N_\alpha$ holds for all α . The image of a solution $(x_\alpha + N_\alpha)_{\alpha \in \Lambda}$ yields a solution of the given sequence.

We state the next theorem without proof since the original proof for commutative valuation domains can be directly adapted to our situation (see [10]).

THEOREM 4.11. *Let M_R be a uniserial module and $(a_\rho)_{\rho \in \Lambda}$ a p.c. sequence in M_R with breadth B . Then there exists a compatible sequence \mathcal{F} of cosets in M_R with the same B and the set of limits of (a_ρ) is equal to the set of solutions for \mathcal{F} . Conversely, for any compatible sequence $\mathcal{F} = (x_\rho + I_\rho)_{\rho \in \Lambda}$ with breadth B there exists a subset $\Lambda_0 \subseteq \Lambda$ such that $(x_\rho)_{\rho \in \Lambda_0}$ is a p.c. sequence with breadth B and $\bigcap_{\rho \in \Lambda} (x_\rho + I_\rho)$ is the set of limits of $(x_\rho)_{\rho \in \Lambda_0}$ or $x_\tau \in \bigcap (x_\rho + I_\rho)$ for some $\tau \in \Lambda$.*

This results shows that a uniserial module M is maximally complete if and only if it is linearly compact or more specifically, every p.c. sequence in M with breadth I has a limit in M if and only if M is linearly I -compact.

The next result shows that maximally complete right chain rings are maximal; it generalizes a theorem by Kaplansky; see [5].

THEOREM 4.12. *Let R be a right chain ring and R_1 an immediate extension of R . Then every element $z \in R_1 \setminus R$ is a limit of a p.c. sequence in R that has no limit in R .*

PROOF. Consider the subset $S = \{v(z - a) | a \in R\} \subseteq \mathcal{W}$. This is well-defined since $z - a \neq 0$ for all $a \in R$. Further, S cannot have a largest element, w say. Otherwise $v(z - a) = w$ and $z - a = bu$, for $b \in R$, $u \in U(R_1)$ and some $a \in R$. However $u = d + j_1$ where d is a representative of $u + J(R_1)$ in R and $j_1 \in J(R_1)$. It follows that $z - a - bd = bj_1$ and $v(z - a - bd) > v(z - a)$.

We can therefore select a well-ordered cofinal sequence in S and the corresponding sequence $(a_\rho)_{\rho \in \Lambda}$ and $a_\rho \in R$ and Λ well-ordered. We have $v(z - a_\sigma) < v(z - a_\rho)$ by definition, $v(\alpha_\sigma - a_\rho) = v(z - a_\rho)$ follows and no last element exists in $\{z - a_\rho \mid \rho \in \Lambda\}$ and $(a_\rho)_{\rho \in \Lambda}$ is a p.c. sequence in R with z one of its limits in R_1 . If we assume that z_1 is a limit of $(a_\rho)_{\rho \in \Lambda}$ in R then $v(z - z_1) = v((z - a_\rho) - (z_1 - a_\rho)) \geq \min\{v(z - a_\rho), v(z_1 - a_\rho)\} = \gamma_\rho = v(z - a_\rho)$ for all ρ . This contradicts the fact that $\{v(z - a_\rho) \mid \rho \in \Lambda\}$ is cofinal in S .

We have the following partial converse of Theorem 4.12.

THEOREM 4.13. *Let $R \subseteq R_1$ be right chain rings with $J(R) = J(R_1) \cap R$ such that for every $x \in R_1 \setminus R$ is a limit of a p.c. sequence in R which has no limit in R . Then R_1 is an immediate extension of R .*

PROOF. We observe that $aR \neq bR$, $a, b \in R$, implies $aR_1 \neq bR_1$ since $J(R) - J(R_1) \cap R$. Otherwise $au = b$, $u \in U(R_1)$ and $aj = b$, $j \in J(R) \subseteq J(R_1)$ and $a(u - j) = 0$, $u - j \in U(R_1)$, $a = 0$. A p.c. sequence (a_ρ) in R is also a p.c. sequence in R_1 . We use Lemma 3.3 and it is enough to show that for $x \in R_1 \setminus R$ we have $xR_1 = aR_1$ for some $a \in R$, since then $x = au_1 = a(u + j_1) = au(1 + u^{-1}j_1)$ for $u_1 \in U(R_1)$ and $u - u_1 \in J(R_1)$ for some $u \in R \cap U(R_1) = U(R)$, since u_1 is in R or the limit of a p.c. sequence in R . The element x is the limit in R_1 of a c.c. sequence (a_ρ) of R which has no limit in R . Hence, $v(a_\rho) = v(a_\sigma)$ for any $\sigma > \rho \geq \lambda$ for some index λ using Lemma 4.2 and the remark after Definition 4.5. If $x = a_\rho j_1$ for $j_1 \in J(R_1)$ then also $x = a_\sigma j_2$ for some $j_2 \in J(R_1)$ and $v(x - a_\rho) = v(a_\rho) = v(a_\sigma) = v(x - a_\sigma)$, a contradiction. Conversely, if $xj_1 = a_\rho$, $j_1 \in J(R_1)$, then also $xj_2 = a_\sigma$ for some $j_2 \in J(R_1)$ and $v(x - a_\rho) = v(x) = v(x - a_\sigma)$, again a contradiction. This leaves $xR_1 = a_\sigma R_1$ as the only possibility and proves the lemma.

PROPOSITION 4.14. *Let $R \subset R_1$ be a proper immediate extension and $x \in R \setminus R_1$ a limit of a p.c. sequence $(a_\rho)_{\rho \in \Lambda}$ in R with breadth B with no limit in R . Then $R/I \subset R_1/IR_1$ for every right ideal $I \subseteq B$.*

PROOF. The p.c. sequence (a_ρ) in R is a p.c. sequence in R_1 with breadth BR_1 if B is its breadth in R . If z is an element in R with $z + BR_1 = x + BR_1$ then $z = x + b_1$, $b_1 \in BR_1$, is a limit of (a_ρ) in R (Lemma 4.6), a contradiction; $R/I \subset R_1/IR_1$ for every right ideal $I \subset B$ follows.

We conclude this section with a result that provides some information on the set of right ideals I' of a right chain ring R for which R is linearly

I' -compact provided R is linearly I -compact. Right ideals A, B of R are called *related* if $s^{-1}A = t^{-1}B$ for some elements $s, t \in R$ with $s \notin A, t \notin B$, where $s^{-1}A = \{x \in R \mid sx \in A\}$. It is proved that R is linearly B -compact provided R is A -compact and B is related to A . However the general problem seems to be open.

Open problem. Assume that R is A -compact for all $A \in \mathcal{A}$ with \mathcal{A} a subset of the set of right ideals of R . Describe all right ideals B for which R must be B -compact too.

THEOREM 4.15. *Let R be a right chain ring and A, B right ideals satisfying $s^{-1}A = t^{-1}B$ for some $s \notin A, t \notin B$. Then the following conditions are equivalent:*

- (i) R is linearly A -compact.
- (ii) R is linearly B -compact.

PROOF. It is sufficient to prove that A -compactness is equivalent to $s^{-1}A$ -compactness, $s \notin A$ and we assume first that R is linearly A -compact and $\mathcal{F} = (x_\alpha + X_\alpha)_{\alpha \in \Lambda}$ a compatible sequence of cosets with $\bigcap X_\alpha = s^{-1}A$. Then $(sx_\alpha + sX_\alpha)_{\alpha \in \Lambda}$ is a compatible sequence of cosets with breadth A and hence a solution a exists.

If $a = sx$ for some $x \in R$, then $s(x - x_\alpha) \in sX_\alpha, x - x_\alpha - n \in X_\alpha$ for $n \in R$ with $sn = 0$. Hence, $n \in s^{-1}A \subseteq X_\alpha$ and x is a solution for \mathcal{F} .

If $aq = s$ for some $q \in J(R)$, then $a - aqx_\alpha = a(1 - qx_\alpha) \in sX_\alpha$ and $a \in sX_\alpha$ follows for all $\alpha \in \Lambda$. We have $a \in sX_\alpha, s \in sX_\alpha$ and $s \in \bigcap sX_\alpha = A$, a contradiction.

If conversely every compatible system with breadth $s^{-1}A$ is solvable let $\mathcal{F}_1 = (a_\alpha + A_\alpha)_{\alpha \in \Lambda}$ be a compatible sequence with $\bigcap A_\alpha = A$. There exists an index α_0 with $A_{\alpha_0} \subset sR$ and we define $\Lambda' = \{\alpha \mid \alpha \geq \alpha_0\}$. It is sufficient to show that $(a_\alpha + A_\alpha)_{\alpha \in \Lambda'}$ has a solution. Since $a_\alpha - a_{\alpha_0} \in A_{\alpha_0}, \alpha \in V'$, there exists $t_\alpha \in R$ with $a_\alpha - a_{\alpha_0} = st_\alpha$. The system $\mathcal{F} = (t_\alpha + s^{-1}A_\alpha)_{\alpha \in \Lambda'}$ has $\bigcap s^{-1}A_\alpha = s^{-1} \bigcap_{\alpha \in \Lambda'} A_\alpha$ as its breadth and is compatible, since $s(t_\alpha - t_\beta) = (a_\alpha - a_{\alpha_0}) \in A_\alpha$ for $\alpha \leq \beta$. Hence, a solution t exists for \mathcal{F} and $a_{\alpha_0} + st$ is a solution for \mathcal{F}_1 .

5. A counterexample

We saw that every element $x \in R_1/R$, with R_1 maximal immediate extension of a right chain ring R , is the limit of a p.c. sequence in R . We

show now by an example that a right chain ring R can be maximal without being maximally complete.

Consider $R = \mathbb{Q}(t_1, t_2, \dots)[x]_{(x)}[[y, \sigma]]$ with $xy = yt_1$, $t_i y = yt_{i+1}$ where $\sum_{n=0}^{\infty} y^n f_n(t_i, x)$ are the elements of R . $\mathbb{Q}(t_1, t_2, \dots)$ is a function field in the variables t_i , $i = 1, \dots$, over the rational numbers \mathbb{Q} , and $\mathbb{Q}(t_1, t_2, \dots)[x]_{(x)}$ is the localization at (x) of the polynomial ring in one variable x over $\mathbb{Q}(t_1, t_2, \dots)$ containing the coefficients $f_n(t_i, x)$. The ring R is then a skew power series ring over this ring in the variable y with $\sigma(x) = t_1$, $\sigma(t_i) = t_{i+1}$ and is a right invariant right chain domain with its right ideals $\neq (0)$ of the form $y^n x^m R$, $n, m \geq 0$.

We show that there cannot exist a proper immediate extension R_1 of R . Otherwise, R_1 is again right noetherian, and hence right invariant with right ideals $y^n x^m R_1$ (by Proposition 3.7(v), Lemma 3.4(iv)). By Theorem 4.12, R_1 contains an element f , $f \notin R$, which is a pseudo limit of a p.c. sequence $(a_\rho)_{\rho \in \Lambda}$ of elements a_ρ in R with breadth B and which has no limit in R . Since B is a right ideal without an upper neighbour in the lattice of right ideals, it follows that B is either (0) or $y^i R$, $i = 1, \dots$. However, R is linearly (0) -compact by construction and if R is not $y^n R$ -compact for some n , then R is not yR -compact, from Theorem 4.15. Finally, using Proposition 4.14, we can assume that R/yR , which is a ring, is contained in but is not equal to R_1/yR_1 , where we use $yR_1 \cap R = yR$ by Proposition 3.4(vi), that is, we can assume that the image of the element f in R_1/yR_1 is not contained in R/yR .

Next, let g be any element in $R_1 \setminus yR_1$. Then g can be written as $g = x^k(c_0 + xh)$ where $k \geq 0$ is an integer, $c_0 \in \mathbb{Q}(t_1, t_2, \dots) \simeq R/J(R) \simeq R_1/J(R_1)$, $c_0 \neq 0$, $h \in R_1$ and $c_0 + xh = u$ is a unit in R_1 . We consider $gy = x^k(c_0 + xh)y = x^k uy - yt_1^k u' = yg'$ with $u' \in U(R_1)$, $g' \in R_1$ and $g' = d_0 + xh_1 \in U(R_1)$ follows for some $h_1 \in R_1$ with $0 \neq d_0 \in \mathbb{Q}(t_1, \dots)$. The element d_0 is uniquely determined by g and we define a mapping Ψ from R_1 to $\mathbb{Q}(t_1, \dots)$ by $\Psi(g) = d_0$ as defined above if $g \in R_1 \setminus yR_1$ and $\Psi(g) = 0$ otherwise. It follows that $\Psi(g_1) + \Psi(g_2) = \Psi(g_1 + g_2)$ for $g_1, g_2 \in R_1$ and we prove $\Psi(g_1, g_2) = \Psi(g_1)\Psi(g_2)$. Let $g_1 y = y(d_0 + xh_1)$, $g_2 y = y(d'_0 + xh_2)$ for $d_0, d'_0 \in \mathbb{Q}(t_1, \dots)$, $h_1, h_2 \in R_1$, then $g_2 g_1 y = y(d'_0 + xh_2)(d_0 + xh_1) = y(d'_0 d_0 + xw)$ for a certain element $w \in R_1$, that is, $\Psi(g_2 g_1) = d'_0 d_0 = \Psi(g_2)\Psi(g_1)$ and Ψ is a ring homomorphism from R_1 to $\mathbb{Q}(t_1, \dots)$ with kernel yR_1 .

Consider $f \in R_1 \setminus R$, $f \notin yR_1$ as chosen above. Then

$$\Psi(f) \in \mathbb{Q}(t_1, \dots) \setminus \mathbb{Q}(t_2, \dots)[t_1]_{(t_1)}$$

since $\Psi(\mathbb{Q}(t_1, \dots)[x]_{(x)}) = \mathbb{Q}(t_2, \dots)[t_1]_{(t_1)}$ and $R/yR = \mathbb{Q}(t_1, \dots)[x]_{(x)} \subset$

R_1/y_1R . This implies $\Psi(f) = t_1^{-k} h_1(t_i)/g_1(t_i)$ with $k \geq 1$, $h_1(t_i), g_1(t_i) \in \mathbb{Q}[t_1, \dots]$ and $(t_1, h_1(t_i)g_1(t_i)) = 1$. There exists g in R with

$$\Psi(g) = g_1(t_i)/h_1(t_i) \in \mathbb{Q}(t_2, \dots)[t_1]_{(t_1)}$$

and it follows that

$$\Psi(x^k g f - 1) = \Psi(x^k) \Psi(g) \Psi(f) - 1 = t_1^k \cdot \frac{g_1(t_i)}{h_1(t_i)} \cdot t_1^{-k} \frac{h_1(t_i)}{g_1(t_i)} - 1 = 0$$

and $x^k g f = 1 + yr'$ for some $r' \in R_1$. This shows that x^k is a unit for some $k > 0$, a contradiction.

References

- [1] C. Bessenrodt, H. H. Brungs and G. Törner, 'Prime ideals in right chain rings,' *Mitt. Math. Sem. Giessen* **163** (1984), 141–167.
- [2] C. Bessenrodt, H. H. Brungs and G. Törner, 'Right chain rings.' Part 1, FB Mathematik, Preprint No. 74, Universität Duisburg, 1985.
- [3] W. Brandal, 'Almost maximal integral domains and finitely generated modules,' *Trans. Amer. Math. Soc.* **183** (1973), 203–222.
- [4] N. I. Dubrovin, 'An example of a nearly simple chain ring with nilpotent elements,' (Russian) *Mat. Sb. (N.S.)* **120** (1983), 441–447.
- [5] I. Kaplansky, 'Maximal fields with valuations,' *Duke Math. J.* **9** (1942), 303–321.
- [6] I. Kaplansky, 'Modules over Dedekind rings and valuation rings,' *Trans. Amer. Math. Soc.* **72** (1952), 327–340.
- [7] W. Krull, 'Allgemeine Bewertungstheorie,' *J. Reine Angew. Math.* **167** (1932), 160–196.
- [8] K. Mathiak, *Valuations of skew fields and projective Hjelmslev spaces*, (Lecture Notes in Mathematics 1175, Springer, Berlin, 1986).
- [9] F. J. Rayner, 'Gravett's proof of a theorem of Krull', *Quart. J. Math.* **24** (1973), 409–410.
- [10] O. Schilling, *The theory of valuation*, (Math. Surveys, Amer. Math. Soc., Providence, R. I., 1950).

University of Alberta
Edmonton, Alberta
Canada T6G 2G1

Fachbereich Mathematik
Universität Duisburg
D 4100 Duisburg
Federal Republic of Germany