

Corrigendum

Conference on 'Malnutrition matters'

Symposium 4: Hot topics in parenteral nutrition Rationale for using new lipid emulsions in parenteral nutrition and a review of the trials performed in adults – Corrigendum

Philip C. Calder

doi:10.1017/S0029665109001268, Published by Cambridge University Press 11 May 2009

In the subsection 'Fish oil in parenteral nutrition' on page 255 (column 2, lines 28–29) incorrect information about the EPA plus DHA content of Lipoplus is given.⁽¹⁾

The information provided (0.6 g per 100 ml) was calculated based upon the EPA and DHA contents of a standard fish oil. However Lipoplus contains more concentrated marine *n*-3 fatty acids as reported by Driscoll *et al.*⁽²⁾ Therefore the statement should read:

Each 100 ml of Lipoplus will typically contain about 1.2 g EPA+DHA.

References

1. Calder PC (2009) Rationale for using new lipid emulsions in parenteral nutrition and a review of the trials performed in adults. *Proc Nutr Soc* **68**, 252–260.
2. Driscoll DF, Ling PR, Bistran BR (2009) Pharmacopeial compliance of fish oil-containing parenteral lipid emulsion mixtures: Globule size distribution (GSD) and fatty acid analyses. *Int J Pharmaceut* **379**, 125–130.