

Dietary supplement use and mortality in a cohort of Swedish men

Maria Messerer, Niclas Håkansson, Alicja Wolk and Agneta Åkesson*

Institute of Environmental Medicine, Karolinska Institutet, Box 210, SE-171 77 Stockholm, Sweden

(Received 18 December 2006 – Revised 5 July 2007 – Accepted 6 July 2007)

The use of dietary supplements has increased substantially in most industrialized countries. The aim of this study was to prospectively examine the association between use of dietary supplements and all-cause mortality, cancer mortality and CVD mortality in men. We used the population-based prospective cohort of 38 994 men from central Sweden, 45–79 years of age, with no cancer or CVD at baseline and who completed a self-administered FFQ including questions on dietary supplement use and life-style factors in 1997. During average 7.7 years of follow-up, 3403 deaths were ascertained; among them, 771 due to cancer and 930 due to CVD (during 5.9 years of follow-up). In multivariate adjusted models including all men there was no association observed between use of any dietary supplement or of multivitamins, vitamin C, vitamin E or fish oil specifically and all-cause mortality, cancer or CVD mortality. Among current smokers, regular use of any supplement was associated with statistically significant increased risk of cancer mortality: relative risk (RR) 1.46 (95% CI 1.06, 1.99). Among men reporting an inadequate diet at baseline (assessed by Recommended Food Score), there was a statistically significant inverse association between use of any dietary supplement and CVD mortality (RR 0.72; 95% CI 0.57, 0.91), no associations were observed among men with adequate diets. In conclusion, we cannot exclude that the use of dietary supplements is harmful for smokers. On the other hand, among men with an insufficient diet, the use of supplements might be beneficial in reducing CVD mortality.

Dietary supplements: Vitamins: All-cause mortality: Cancer mortality: Cardiovascular mortality: Prospective studies

Over the last decades the use of dietary supplements has increased substantially in most industrialized countries^{1,2}. The users are paying significant amounts of money in expectation to decrease their morbidity and mortality, but there is little evidence of such effects.

A few randomized controlled supplementation trials have shown decreased cancer incidence and all-cause mortality in European³ and Chinese populations⁴, but baseline micronutrients status was considered suboptimal in the study participants. In general, the conducted randomized trials show little support of a preventive effect of dietary supplement use on cancer and CVD development^{5–8} and even increased mortality has been reported in certain high-risk groups such as smokers and exposed workers^{9–11}. Randomized controlled trials may, however, be too short to detect any effects, particularly regarding the incidence of cancer. In addition, with few exceptions, trials tend to use relatively high dosages of a few agents and focus on persons with high risk for disease or those with existing disease and therefore are not necessarily valid for the effects of long-term, low doses of several agents in the general population¹².

Observational studies offer the possibility to investigate the effect on mortality of the long-term use of low doses of several different micronutrients. The results from prospective cohort studies are, however, inconsistent. Absence of any association between use of specific dietary supplements and

mortality was observed in several cohort studies^{13–16}, while others suggested an inverse association between use of single or combinations of vitamins and/or multivitamins and all-cause mortality^{17,18}, CVD mortality^{18–21} and cancer mortality^{22–24}. Inconsistent results may in part be explained by the fact that the use of dietary supplements is often associated with other lifestyle factors that may predict the risk of disease^{25–28}. In addition, there is a need for observational studies exploring the association between supplement use and mortality taking into account diet adequacy/deficiency.

In the present study we explored the association between use of dietary supplements and mortality in relation to diet quality in a prospective population-based cohort of Swedish middle-aged and elderly men, taking lifestyle factors such as self-perceived health status into consideration.

Methods

A population-based prospective cohort of Swedish men (COSM), living in central Sweden (Västmanland and Örebro Counties), was established during the autumn of 1997. All men 45–79 years of age, living in the area (identified in the population register; *n* 100 303) received a mailed FFQ, and 48 850 responded. The COSM study well represents the Swedish male population with regard to age distribution, relative weight and educational level as compared with representative

Abbreviations: RR, relative risk.

* **Corresponding author:** Dr Agneta Åkesson, fax +46 8 30 45 71, email Agneta.Akesson@ki.se

data from Statistics Sweden²⁹. Men with a previous cancer diagnosis other than non-melanoma skin cancer (*n* 2592) identified by linkage to the National Swedish Cancer Registry, and a history of CVD according to the International Classification of Diseases (*n* 5069), identified by linkage to the National Hospital Discharge Registry, were excluded. We also excluded men with missing or incorrect national identification numbers, participants returning their questionnaires empty or with incomplete data regarding dietary supplement use, leaving 38,994 men at baseline. The regional Ethical Review Board at Karolinska Institutet approved the study.

The use of dietary supplements was self-reported to be either regular, occasional or no use³⁰. Regular and occasional users further specified what kind of predefined supplement (multivitamins, vitamin C, vitamin E and fish oil) they used. The estimated mean content of a multivitamin preparation used in the study population was 60 mg vitamin C, 9 mg vitamin E, 1.2 mg thiamine, 1.4 mg riboflavin, 1.8 mg vitamin B₆, 3 µg vitamin B₁₂ and 400 mg folic acid. For supplements containing only one specific nutrient the estimated content was: 1000 mg vitamin C, 100 mg vitamin E. The mean content of fish oil was 500 mg (18% EPA and 12% docosapentaenoic acid, respectively). Sensitivity and specificity of self-reported use of any dietary supplement in the FFQ was 78 and 93%, respectively³¹.

The FFQ contained questions on food frequencies including alcoholic beverages (ninety-six items), sociodemographic data, height, weight, smoking status, physical activity (time spent on walking/bicycling and exercising), self-perceived health status, hypertension, high cholesterol levels and diabetes. The validity of the FFQ in comparison to fourteen 24 h recalls in 248 men varied between (Spearman correlation coefficient; *r*) 0.38 and 0.81 (mean 0.62) for total micronutrients (including supplements)^{30,32}. The FFQ was used to define diet according to recommended food scores which is a simple way to define the quality by separating 'healthy' from 'less healthy' foods, based on current knowledge and dietary guidelines³³. Such a score has been shown to be highly predictive of mortality^{33,34}. The recommended food scores included fruits, vegetables, legumes, nuts, low-fat dairy products, whole-grain products and fish. Each food with a consumed frequency of at least one to three times/month, was assigned a food score of one. This added up to a maximum of 25 for the recommended food score. We considered those who scored in the lowest quintile (scores <16) of the recommended food score as having an inadequate diet, while the other (scores 16–25) were considered to have an adequate diet.

Date of death was ascertained through linkage of the cohort to the Swedish Death Registry at Statistics Sweden. The follow-up period for all-cause mortality extended from 1998 to 2005 (8 years), while specified causes of death extended to 2003 (6 years), ascertained through linkage to the Swedish Registry of Causes of Death at the National Board of Health and Welfare. This registry is considered almost 100% complete. We classified cause of death according to the International Classification of Diseases 10 (CVD codes I00–I79 and cancers codes C00–C97).

Relative risk (RR) and 95% CI were estimated using the Cox proportional hazards models. Follow-up was censored at date of death or end of follow-up period, whichever occurred first. RR values were adjusted for age (continuous),

BMI (<20, 20–25, 25–30 or >30), smoking status (never smoker, former smoker, current smoker), education (≤9 years, 10–12 years, ≥12 years), marital status (single, married, divorced, widowed), physical activity (≤30, 30–68 and ≥68 min/d), self-perceived health status (five categories), recommended food score (tertiles), hypertension (yes/no), high cholesterol (yes/no) and diabetes (yes/no).

Results

Among the 38 994 men in the cohort, 13 295 (34%) reported any use of dietary supplements. Out of the users, 6049 (15%) reported regular use and 7246 (19%) reported occasional use of one or more dietary supplements. Among the different supplements, multivitamins were used by 5530 (14%), vitamin C by 4328 (11%), vitamin E by 1593 (4%) and fish oil by 1869 (5%) men in the cohort. Compared to men who did not use supplements, users tended to be older, more educated, more physically active and more seldom current smokers (Table 1).

During 300 725 person-years (mean 7.7 years) of follow-up from 1998 to 2005 for all-cause mortality, 3403 deaths were ascertained. During the cause-specific mortality follow-up from 1998 to 2003 of 228 270 person-years (mean 5.9

Table 1. Age-standardized baseline characteristics of a cohort of 38 994 Swedish men, aged 45–79 years, stratified by use of any dietary supplements, presented as mean or percentage (*n* 38 994)

Characteristics	Non-users	Users
No. of subjects	25 699	13 295
% of total subjects	66	34
Age (years)		
Mean	58.5	60.5
SD	9.2	9.7
BMI (kg/m ²)		
Mean	25.9	25.4
SD	3.3	3.2
Smoking status (%)		
Never smokers	36.6	38.8
Former smokers	37.4	39.3
Current smokers	26	21.9
Education (%)		
Less than high school	70.1	62.4
High school	14.0	16.0
University	15.9	21.6
Married (%)	83.5	81.9
Physical activity (min/d)*		
Mean	55.2	58
SD	44.0	43.4
Self-perceived health status (%)		
Very good	23.7	22.50
Good	55.5	55.0
Average	19.0	20.3
Bad	1.6	2.0
Very bad	0.3	0.3
Recommended food score†		
Mean	18.3	19.0
SD	4.4	4.4
Hypertension (%)	20.0	20.2
High cholesterol levels (%)	11.0	11.8
Diabetes mellitus (%)	5.6	5.2

* Time spent on walking/bicycling and exercising.

† Each point refers to the consumption of fruits, vegetables, legumes, nuts, low-fat dairy products, whole-grain products and fish at least one to three times per month (maximum 25).

years), 771 deaths occurred due to cancer and 930 due to CVD.

There was no association between the overall use of dietary supplements and risk of all-cause mortality, CVD or cancer mortality in age-adjusted models or after adjustment for relevant covariates (Table 2). The risk estimates remained the same after further adjustment for alcohol consumption (g ethanol/d). In order to minimize a possible impact on the associations by subjects starting to take supplements because of a not yet diagnosed morbidity, we stratified the analyses by time of follow-up. There were no significant associations between regular users of any supplement and all-cause mortality, cancer mortality and CVD mortality after exclusion of the first 3 years of follow-up (Table 2). There were no associations between the use of any specific supplement (multivitamins, vitamin C, vitamin E or fish oil) and risk of all-cause mortality, CVD or cancer mortality (data not shown).

In analyses stratified by smoking status we observed, among current smokers, a statistically significant positive association corresponding to 46% increased risk (95% CI, 6, 99) between regular use of any dietary supplement and cancer mortality in comparison to no use (Table 3). This association persisted even after exclusion of the first 3 years of follow-up (RR 1.48; 95% CI 1.01, 2.16). There were, however, no associations between supplement use and CVD mortality or all-cause mortality. Among never smokers, there was a tendency of a statistically significant inverse association between regular use of any supplement and cancer mortality (RR 0.68; 95% CI, 0.46, 1.01).

To assess whether a deficient diet may affect the association between use of supplements and risk of mortality, we stratified the diet according to inadequate (the lowest quintile of recommended food score) and adequate intake (quintiles 2–5). Men who reported an inadequate diet and used dietary supplements had a 28% statistically significant lower risk of CVD mortality (95% CI 9, 43) compared to never users (Table 4). This association was not observed among those with an adequate diet. After exclusion of the first 3 years of follow-up, the RR of CVD mortality was 0.76 (95% CI 0.56, 1.01). There was a non-significant inverse association

between supplement use and cancer mortality (RR 0.88; 95% CI 0.67, 1.16).

Discussion

In this large population-based prospective cohort of middle-aged and elderly Swedish men, we found no association between use of any dietary supplement or of multivitamins, vitamin C, vitamin E or fish oil specifically and all-cause mortality, cancer or CVD mortality in all men. We observed, however, that among current smokers, regular use of any supplement was associated with 46% increased risk of death due to cancer. In addition, among men with an inadequate diet there was a statistically significant inverse association, corresponding to 28% decreased risk, between combined occasional and regular supplement use and CVD mortality.

The present results of an inverse association between the use of any supplement and decreased risk of CVD mortality taking into account quality of diet are in agreement with those observed in randomized trials from the Linxian Province in China, a rural area with a chronically low intake of several nutrients. Men, but not women, had after a 6-year multivitamin/mineral supplement trial a 58% decreased CVD mortality (95% CI 7, 81)³⁵. Another trial performed in the same area showed a similar reduction in CVD mortality among men³⁶. A lower cancer incidence restricted to men in the SU.VI.MAX randomized trial in France was suggested to be caused by a lower baseline antioxidant status in men than in women³. These results are in agreement with the present findings for men with inadequate diet. Among observational studies that have not considered adequacy of diet, some have reported inverse associations between the use of single or combinations of vitamins and/or multivitamins and all-cause mortality^{17,18}, CVD mortality^{18–21} and cancer mortality^{22–24}.

In analyses stratified by smoking status, we found a statistically significant increased risk between regular use of any supplement and cancer mortality among current smokers. This agrees with results from a large prospective cohort study, showing an increased cancer mortality rate among smokers who used multivitamin supplements¹⁷ and a suggested

Table 2. Relative risk (RR) of all-cause mortality, cancer mortality and CVD mortality according to non-specific supplement use in a cohort of 38 994 Swedish men

Use of supplements	No. of deaths	RR*	95% CI	RR†	95% CI	No. of deaths‡	RR†	95% CI
All-cause mortality								
No use	2102	1§		1§		1591	1§	
Occasional	631	0.93	0.85, 1.02	0.94	0.85, 1.02	480	0.94	0.85, 1.04
Regular	670	1.00	0.91, 1.09	1.04	0.95, 1.13	487	1.00	0.90, 1.11
Cancer mortality								
No use	479	1§		1§		319	1§	
Occasional	140	0.92	0.76, 1.11	0.93	0.77, 1.12	85	0.85	0.67, 1.09
Regular	152	1.02	0.85, 1.23	1.07	0.89, 1.29	103	1.10	0.88, 1.38
CVD mortality								
No use	582	1§		1§		355	1§	
Occasional	163	0.85	0.71, 1.01	0.86	0.72, 1.02	98	0.84	0.67, 1.06
Regular	185	0.95	0.80, 1.11	1.00	0.85, 1.19	104	0.93	0.74, 1.16

* Adjusted for age.

† Fully adjusted model (adjusted for age, BMI, smoking status, education, marital status, physical activity, self-perceived health and recommended food score, hypertension, high cholesterol levels and diabetes).

‡ The first 3 years of follow-up excluded (follow-up comprises 2001–2005 for all-cause and 2001–2003 for cause-specific mortality).

§ Reference group.

Table 3. Relative risk (RR) of all-cause mortality, cancer mortality and CVD mortality in relation to use of non-specific supplements, stratified by smoking status in a cohort of Swedish men*

Use of supplements	Never smokers			Former smokers			Current smokers		
	No. of deaths	RR	95 % CI	No. of deaths	RR	95 % CI	No. of deaths	RR	95 % CI
All-cause mortality									
No	601	1.00†		731	1.00†		730	1.00†	
Occasional	192	0.91	0.77, 1.07	227	0.99	0.85, 1.15	197	0.63	0.79, 1.09
Regular	188	0.91	0.77, 1.08	267	1.10	0.96, 1.27	202	1.14	0.97, 1.34
Cancer mortality									
No	134	1.00†		172	1.00†		164	1.00†	
Occasional	47	0.98	0.70, 1.37	45	0.85	0.61, 1.18	47	1.01	0.73, 1.40
Regular	31	0.68	0.46, 1.01	64	1.14	0.85, 1.52	55	1.46	1.06, 1.99
CVD mortality									
No	156	1.00†		213	1.00†		205	1.00†	
Occasional	44	0.79	0.56, 1.11	65	0.97	0.73, 1.28	51	0.83	0.61, 1.13
Regular	55	1.00	0.73, 1.36	77	1.09	0.83, 1.41	49	0.90	0.65, 1.25

* Each stratified model is adjusted for age, BMI, education, marital status, physical activity, self-perceived health, recommended food score, hypertension, high cholesterol levels and diabetes.

† Reference group.

increased mortality rate among elderly smokers from eleven European countries (the SENECA study) who used supplements³⁷. Increased mortality was also reported in two intervention trials among high-risk individuals (smokers and workers exposed to asbestos) taking high dosage of β -carotene^{9,10}. It was suggested that β -carotene affected the growth of preclinical tumours rather than the induction of tumours³⁸. The free radical-rich environment produced by cigarette smoke and the resultant inflammatory response in the lung may combine to induce oxidation of some antioxidants, resulting in a pro-oxidant effect³⁹⁻⁴¹.

Strengths of the present study are the population-based and prospective design and the practically complete follow-up of the study population through linkage with computerized registers. We were also able to perform separate analyses among smokers and among men considered to have a deficient diet while controlling for several potential confounders. As a more frequent use of dietary supplements has been associated with low self-perceived health^{27,28}, there is a risk of reverse causality in cohort studies, especially

during a short follow-up. In the present study, we were able to control for the responders' self-perceived health status and exclusions of possible not yet diagnosed diseases during the first 3 years of follow-up did not alter our conclusion.

On the other hand, if supplement use is assessed only at baseline, a longer follow-up will increase the risk of exposure misclassification attenuating the observed estimates, given that the use of these preparations varies over time. Although the sensitivity and the specificity of the self-reported supplement use in the present study were satisfactory³¹, the main limitation associated with present study may be that the use of supplements was assessed only at baseline. Thus, there is a potential risk that the absence of statistically significant association between supplement use and mortality in some of the analyses in the data could be due to measurement error.

In conclusion, although present study results need to be confirmed by other studies, they suggest that supplement use may have a harmful effect among smokers and a beneficial effect when the diet is inadequate.

Table 4. Relative risk (RR) of all-cause mortality, cancer mortality and CVD mortality in relation to any use (occasional or regular) of non-specific supplements, stratified by diet quality in a cohort of Swedish men*

Use of supplement	Inadequate diet (21 %)†			Adequate diet (79 %)		
	No. of deaths	Any use		No. of deaths	Any use	
		RR	95 % CI		RR	95 % CI
All-cause mortality						
No	740	1.00‡		1360	1.00‡	
Yes	399	0.94	0.83, 1.06	901	1.01	0.92, 1.10
Cancer mortality						
No	157	1.00‡		322	1.00‡	
Yes	77	0.88	0.67, 1.16	215	1.04	0.87, 1.24
CVD mortality						
No	235	1.00‡		346	1.00‡	
Yes	103	0.72	0.57, 0.91	245	1.07	0.91, 1.27

* Adjusted for age, BMI, smoking status, education, marital status, physical activity, self-perceived health, recommended food score, hypertension, high cholesterol levels and diabetes.

† Classified as the lowest quintile of recommended food score (scores < 16).

‡ Reference group.

Acknowledgements

The authors acknowledge the Swedish Research Council/Longitudinal Studies, the Swedish Research Council/Medicine and the Centre of Health Care Sciences, Karolinska Institutet. M. Messerer wrote the first draft of the paper and contributed to the statistical analysis. A. Åkesson was the principal investigator of this study, contributed substantially to the writing of the paper and made all revisions. N. Håkansson performed the statistical analysis. A. Wolk, principal investigator of the cohort of Swedish men, supervised the study and contributed to the writing of the paper. All the authors contributed to manuscript editing. There are no conflicts of interest for any of the authors.

References

- Messerer M, Johansson SE & Wolk A (2001) Use of dietary supplements and natural remedies increased dramatically during the 1990s. *J Intern Med* **250**, 160–166.
- Millen AE, Dodd KW & Subar AF (2004) Use of vitamin, mineral, nonvitamin, and nonmineral supplements in the United States: the 1987, 1992, and 2000 National Health Interview Survey results. *J Am Diet Assoc* **104**, 942–950.
- Hercberg S, Galan P, Preziosi P, *et al.* (2004) The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* **164**, 2335–2342.
- Blot WJ, Li JY, Taylor PR, *et al.* (1993) Nutrition intervention trials in Linxian, China: supplementation with specific vitamin/mineral combinations, cancer incidence, and disease-specific mortality in the general population. *J Natl Cancer Inst* **85**, 1483–1492.
- Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG & Gluud C (2007) Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* **297**, 842–857.
- Morris CD & Carson S (2003) Routine vitamin supplementation to prevent cardiovascular disease: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* **139**, 56–70.
- Bjelakovic G, Nikolova D, Simonetti RG & Gluud C (2004) Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet* **364**, 1219–1228.
- Miller ER 3rd, Pastor-Barriuso R, Dalal D, Riemersma RA, Appel LJ & Guallar E (2005) Meta-analysis: high-dosage vitamin E supplementation may increase all-cause mortality. *Ann Intern Med* **142**, 37–46.
- The Alpha-Tocopherol Beta-Carotene Cancer Prevention Study Group (1994) The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med* **330**, 1029–1035.
- Omenn GS, Goodman GE, Thornquist MD, *et al.* (1996) Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *N Engl J Med* **334**, 1150–1155.
- Virtamo J, Pietinen P, Huttunen JK, *et al.* (2003) Incidence of cancer and mortality following alpha-tocopherol and beta-carotene supplementation: a postintervention follow-up. *JAMA* **290**, 476–485.
- Byers T (1999) What can randomized controlled trials tell us about nutrition and cancer prevention? *CA Cancer J Clin* **49**, 353–361.
- Kushi LH, Folsom AR, Prineas RJ, Mink PJ, Wu Y & Bostick RM (1996) Dietary antioxidant vitamins and death from coronary heart disease in postmenopausal women. *N Engl J Med* **334**, 1156–1162.
- Knekt P, Reunanen A, Jarvinen R, Seppanen R, Heliovaara M & Aromaa A (1994) Antioxidant vitamin intake and coronary mortality in a longitudinal population study. *Am J Epidemiol* **139**, 1180–1189.
- Yochum LA, Folsom AR & Kushi LH (2000) Intake of antioxidant vitamins and risk of death from stroke in postmenopausal women. *Am J Clin Nutr* **72**, 476–483.
- Muntwyler J, Hennekens CH, Manson JE, Buring JE & Gaziano JM (2002) Vitamin supplement use in a low-risk population of US male physicians and subsequent cardiovascular mortality. *Arch Intern Med* **162**, 1472–1476.
- Watkins ML, Erickson JD, Thun MJ, Mulinaire J & Heath CW Jr (2000) Multivitamin use and mortality in a large prospective study. *Am J Epidemiol* **152**, 149–162.
- Losonczy KG, Harris TB & Havlik RJ (1996) Vitamin E and vitamin C supplement use and risk of all-cause and coronary heart disease mortality in older persons: the Established Populations for Epidemiologic Studies of the Elderly. *Am J Clin Nutr* **64**, 190–196.
- Enstrom JE, Kanim LE & Klein MA (1992) Vitamin C intake and mortality among a sample of the United States population. *Epidemiology* **3**, 194–202.
- Sahyoun NR, Jacques PF & Russell RM (1996) Carotenoids, vitamins C and E, and mortality in an elderly population. *Am J Epidemiol* **144**, 501–511.
- Knekt P, Ritz J, Pereira MA, *et al.* (2004) Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *Am J Clin Nutr* **80**, 1508–1520.
- Jacobs EJ, Henion AK, Briggs PJ, *et al.* (2002) Vitamin C and vitamin E supplement use and bladder cancer mortality in a large cohort of US men and women. *Am J Epidemiol* **156**, 1002–1010.
- Jacobs EJ, Connell CJ, Patel AV, *et al.* (2001) Vitamin C and vitamin E supplement use and colorectal cancer mortality in a large American Cancer Society cohort. *Cancer Epidemiol Biomarkers Prev* **10**, 17–23.
- Chan JM, Stampfer MJ, Ma J, Rimm EB, Willett WC & Giovannucci EL (1999) Supplemental vitamin E intake and prostate cancer risk in a large cohort of men in the United States. *Cancer Epidemiol Biomarkers Prev* **8**, 893–899.
- Block G, Sinha R & Gridley G (1994) Collection of dietary-supplement data and implications for analysis. *Am J Clin Nutr* **59**, Suppl. 1, 232S–239S.
- Foote JA, Murphy SP, Wilkens LR, Hankin JH, Henderson BE & Kolonel LN (2003) Factors associated with dietary supplement use among healthy adults of five ethnicities: the Multiethnic Cohort Study. *Am J Epidemiol* **157**, 888–897.
- Messerer M, Johansson SE & Wolk A (2001) Sociodemographic and health behaviour factors among dietary supplement and natural remedy users. *Eur J Clin Nutr* **55**, 1104–1110.
- Brustad M, Braaten T & Lund E (2004) Predictors for cod-liver oil supplement use - the Norwegian Women and Cancer Study. *Eur J Clin Nutr* **58**, 128–136.
- Norman A, Bellocco R, Vaida F & Wolk A (2002) Total physical activity in relation to age, body mass, health and other factors in a cohort of Swedish men. *Int J Obes Relat Metab Disord* **26**, 670–675.
- Messerer M, Johansson SE & Wolk A (2004) The validity of questionnaire-based micronutrient intake estimates is increased by including dietary supplement use in Swedish men. *J Nutr* **134**, 1800–1805.
- Messerer M & Wolk A (2004) Sensitivity and specificity of self-reported use of dietary supplements. *Eur J Clin Nutr* **58**, 1669–1671.

32. Messerer M (2004) *Dietary Supplements – Trends, Demographics and Mortality Among Users*. Stockholm: Institute of Environmental Medicine, Karolinska Institutet.
33. Michels KB & Wolk A (2002) A prospective study of variety of healthy foods and mortality in women. *Int J Epidemiol* **31**, 847–854.
34. Kant AK, Schatzkin A, Graubard BI & Schairer C (2000) A prospective study of diet quality and mortality in women. *JAMA* **283**, 2109–2115.
35. Mark SD, Wang W, Fraumeni JF Jr, *et al.* (1996) Lowered risks of hypertension and cerebrovascular disease after vitamin/mineral supplementation: the Linxian Nutrition Intervention Trial. *Am J Epidemiol* **143**, 658–664.
36. Blot WJ, Li JY, Taylor PR, Guo W, Dawsey SM & Li B (1995) The Linxian trials: mortality rates by vitamin-mineral intervention group. *Am J Clin Nutr* **62**, Suppl. 6, 1424S–1426S.
37. Brzozowska A, Kaluza J, de Groot C, Knoop K & Amarin Cruz J (2004) Supplementation practice and mortality in Seneca population [abstract]. *J Nutr Health Aging* **8**, 462.
38. Goodman GE, Thornquist MD, Balmes J, *et al.* (2004) The Beta-Carotene and Retinol Efficacy Trial: incidence of lung cancer and cardiovascular disease mortality during 6-year follow-up after stopping beta-carotene and retinol supplements. *J Natl Cancer Inst* **96**, 1743–1750.
39. Duffield-Lillico AJ & Begg CB (2004) Reflections on the landmark studies of beta-carotene supplementation. *J Natl Cancer Inst* **96**, 1729–1731.
40. Mayne ST, Handelman GJ & Beecher G (1996) Beta-carotene and lung cancer promotion in heavy smokers - a plausible relationship? *J Natl Cancer Inst* **88**, 1513–1515.
41. Lee KW, Lee HJ, Surh YJ & Lee CY (2003) Vitamin C and cancer chemoprevention: reappraisal. *Am J Clin Nutr* **78**, 1074–1078.