

referrals. In March 2021, authors delivered teaching on appropriate use of the MHA to local Police (Gardaí) and out-of-hours General Practitioners (GPs).

Results. Of 78 referrals for involuntary admission in 2019 and 115 in 2021, 19% and 44% respectively were unsuccessful (AO not signed). In 60% of unsuccessful referrals in 2019, the person presented with no mental disorder meeting criteria for involuntary detention. The same figure in 2021 was 27%. Individuals presented with substance misuse disorder in 73% and 27% of unsuccessful referrals in 2019 and 2021 respectively. In 2019, 93% of unsuccessful referrals came from Gardaí, 100% came from medical practitioners other than the patient's own GP and 73% came outside of normal working hours.

Conclusion. Unsuccessful referrals for involuntary admission are not uncommon. Audit cycle 1 highlighted that those subject to unsuccessful referrals frequently present out-of-hours, with substance misuse disorder and/or no mental disorder meeting criteria for involuntary admission. In 2021 a smaller proportion of unsuccessful referrals presented with no mental disorder and substance misuse disorder. This suggests that Gardaí/GP education offered benefit. We observed an increase in both the overall number of referrals and the percentage of unsuccessful referrals in 2021. This possibly reflects an impact of the COVID-19 pandemic. There is a need for ongoing education, discussion and feedback with Gardaí and GPs on the process of making referrals for involuntary admission. A limitation is that separate registrars completed each audit cycle.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Consideration of Clozapine in Patients With Treatment Resistant Schizophrenia

Dr Niema Moazzami* and Dr Aimee Devereux
Bradford District Care Trust, Bradford, United Kingdom
*Presenting author.

doi: 10.1192/bjo.2024.603

Aims. The aim of the audit is to measure the performance of Bradford District Care Trust (BDCT) against the NICE guideline's quality standard: 'Adults with schizophrenia that has not responded adequately to treatment with at least 2 antipsychotic drugs are offered clozapine.'

Methods. We identified the 273 patients prescribed clozapine in BDCT as of January 2023. 202 had been prescribed clozapine for more than 5 years and these were excluded. Of the remaining 71, 34 were excluded for reasons such as their diagnosis i.e. Parkinson's or personality disorder or inadequate information within the clinical record. The final sample consisted of 37 patients prescribed clozapine within the last 5 years with a diagnosis of treatment resistant schizophrenia (TRS).

To define an 'adequate trial' of an antipsychotic, an adequate duration and adequate dose was determined. It is widely recommended in literature that an adequate trial of antipsychotic should last at least 6 weeks and this time constraint was utilised for the audit. The Maudsley Guidelines minimum effective dose table was utilised for establishing adequate doses. For antipsychotics not included in this list the British National Formulary (BNF) maintenance doses were used. A data collection tool was then developed that allowed for retrospective collection of key information relating to the objectives outlined above.

Results. Clozapine was offered at the appropriate time [this includes where clozapine was considered but was not felt to be suitable/was contraindicated/declined by the patient] in 13 cases (35%). In 24 cases (65%) clozapine was not offered at the appropriate time. For 21 patients there was a delay in offering clozapine after 2 adequate trials of antipsychotic medication had been given. Delays ranged between 9 days and 15 years, with the average [mean] delay being 3.7 years. There were 3 patients who were initiated on clozapine without the completion of 2 adequate trials of other antipsychotic medication.

Conclusion. In summary, this audit measured BDCT's adherence to the NICE guideline on clozapine for TRS. Of the 37 patients in the final sample, 35% received clozapine at the appropriate time, while 57% experienced delays with an average delay of 3.7 years. Notably, 8% commenced clozapine without completing the recommended antipsychotic trials. These results emphasize the importance of improving adherence to the guideline, as delays in offering clozapine pose potential risks for patients with TRS. Ensuring timely access to this treatment is crucial, as it has the potential to significantly enhance patient outcomes.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Are We Monitoring Our Lithium Patients as Well as We Should be? A Two-Cycle Audit Evaluating Lithium Clinic Documentation in a District General Hospital in Wales

Dr Huda Mohammed*, Dr Rahatul Islam, Dr Cressida Sparrow and Dr Timothy Chan

Princess of Wales Hospital, Bridgend, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.604

Aims. Lithium is a mood-stabiliser with a narrow therapeutic index. Patients are known to be at risk of lithium toxicity if they are unaware of how to recognise its signs. NICE guidelines outline the information that must be relayed to these patients. Furthermore, *GMC Good Medical Practice* highlights the importance of clear and contemporaneous patient records that contain relevant clinical information.

The aim of this two-cycle audit was to assess the quality of documentation for patients reviewed in lithium clinic and to analyse the consistency of the notes recorded to ensure high quality care provision and communication within the department, in line with the NICE guidelines.

Methods. The inclusion criteria were patients over the age of 65, prescribed lithium and were actively reviewed in the monthly lithium clinic at the Older Person's Mental Health Service (OPMHS) at Princess of Wales Hospital in Wales.

A data collection form was created to ensure all the relevant data in line with NICE guidelines was captured including serum lithium level, lithium dose, other psychotropics, side effects, renal function, patient mood, safety netting advice provided including signs of toxicity and awareness of lithium card. A standard of 100% was set for this data to be captured for each patient.

Results. Cycle 1 was completed in November 2023 where a total of 18 patient records were selected (N = 18). Lithium dose, lithium level and renal function were recorded in over 83.3% (n = 15) of the files audited. Details on psychotropics were recorded in 61.1% (n = 11), side effects in 50% (n = 9) and patient mood in 77.8% (n = 14). Safety netting advice was recorded in