

## Editorial

## Supporting mental health disclosure decisions: the Honest, Open, Proud programme

Katrina Scior, Nicolas Rüsçh, Chris White and Patrick W. Corrigan

**Summary**

The stigma associated with mental health problems leaves many feeling they have to 'hide' their difficulties. Supporting them in making disclosure decisions can potentially improve well-being, reduce self-stigma and support recovery processes. In this editorial we discuss the case for interventions designed for this purpose and present one prominent programme: Honest, Open, Proud.

**Declaration of interest**

None.

**Keywords**

Stigma and discrimination; disclosure; concealment; decision-making; peer support.

**Copyright and usage**

© The Authors 2019.

**Katrina Scior** (pictured) is an associate professor in clinical psychology in the Research Department of Clinical Educational and Health Psychology at University College London. **Nicolas Rüsçh** is a professor for public mental health in the Department of Psychiatry II at the University of Ulm and BKH Günzburg. **Chris White** is a policy and information officer at the Mental Health Foundation in Scotland. **Patrick W. Corrigan** is Distinguished Professor of Psychology at Illinois Institute of Technology and the principal investigator in the National Consortium on Stigma and Empowerment, USA.

**Why focus on disclosure**

Individuals experiencing mental health problems (and their families) face two primary challenges as a result of their mental health problems: the symptoms themselves and the stigma associated with these symptoms and diagnostic labels. Although treatment innovations and services mainly address the former, stigma and its consequences often remain unaddressed, despite abundant evidence that stigma creates immense distress and inhibits recovery. Because of common negative stereotypes, many experiencing mental health difficulties not only experience discrimination, but also experience shame, think less of themselves and feel they are somehow to blame for their difficulties. Stigma also leads many of those affected to feel that they have to 'hide' their difficulties or diagnosis from others, to avoid being viewed in a negative light and becoming the target of discrimination. Although concealment may have real or perceived benefits for the person, a fear of 'being found out' often creates additional stress and distress for the person. Secrecy can also act as a barrier to accessing informal and formal support, both of which are central to recovery from mental health problems. Accordingly, supporting individuals to reach decisions around disclosure and act on these in ways that are personally meaningful and safe has multiple potential benefits, and ultimately supports recovery processes through generating hope, reducing shame and enhancing self-esteem.

**Supporting disclosure decisions**

The most prominent approaches to supporting individuals in reaching careful disclosure decisions are Honest, Open, Proud (HOP)<sup>1</sup> and Conceal or Reveal (CORAL).<sup>2</sup> They have some key differences.

HOP (originally called 'Coming Out Proud') is a group programme, usually delivered over three sessions and considers disclosure across settings. A community-based participatory model and empowerment are central to the ethos of HOP, and as such it is delivered as a peer-facilitated (or co-facilitated) group programme. In contrast, CORAL is a decision aid focused specifically on disclosure within an employment context, and is usually delivered in one session by an employment adviser with a primary aim of reducing decisional conflict.<sup>2</sup>

The HOP programme in particular has proven very popular with patient groups and its dissemination and adaptation have developed rapidly. HOP, in its original form or an adapted version, is now being delivered in the USA, as well as in Australia, Belgium, Canada, Chile, China, Germany, Israel, Italy, Switzerland and the UK. Hence in this editorial, we review some of the central premises of HOP, initial evidence on its impact, and consider the way forward for HOP and other approaches designed to support decisions regarding disclosure of mental health problems and other stigmatised conditions.

**The HOP programme**

The key premises of HOP are that disclosure of mental health problems is an individual and personal decision, that disclosure decision-making is mostly an ongoing process (unless someone has very publicly 'come out'), and that the potential benefits and costs of disclosure vary depending on the context and need to be carefully balanced. Accordingly, HOP supports individuals in reaching careful decisions relating to disclosure of current or past mental health problems and planning actions informed by such decisions within a peer-support setting. HOP does not shy away from fully recognising that disclosure can bring benefits, but also poses risks. Hence it helps participants to weigh the pros and cons of disclosure and does not have the goal of persuading them to disclose, but rather supports them in reaching a decision that feels right for them at a given moment in time. In the process, participants have the opportunity to construct a coherent narrative of their mental health problems, before deciding what, if anything, of this they want to share with others.

The programme is delivered using a manual and accompanying workbook, available on the programme website (<http://www.honestopenproud.org>). It is organised into three lessons plus a

booster lesson, with a clear suggestion that participants draw on the materials and lessons learnt in revisiting disclosure decisions over time. In lesson one, participants are encouraged to reflect on beliefs they hold about themselves and their mental health problems and to challenge hurtful, self-stigmatising beliefs. They are guided in carefully considering the potential benefits and costs of disclosing their mental health problems in different contexts, and weighing these up in reaching a decision about disclosure. In lesson two, participants are introduced to different ways of managing disclosure decisions, such as social withdrawal to avoid being found out at one end of the decision continuum, disclosure to selected individuals and actively sharing one's experiences with mental health problems with others. Programme participants are guided in anticipating how others might respond. Lesson three supports participants in telling their personal story in a safe space by inviting them to write an account of their history of mental health problems, and in deciding what, if anything, they want to share, and with whom, how and when. Those who decide to disclose are supported in considering carefully how to share their experiences. Finally, a booster session, in which participants reflect on disclosure experiences and review their decision, takes place around a month after lesson three. Those who decide against disclosure at the present time are encouraged to reflect on the benefits and costs of non-disclosure and to revisit their decision at future intervals.

In addition to the original peer group version of HOP, various adaptations have been developed to tailor HOP to specific groups and cultural contexts, including versions for adolescents and college students, survivors of attempted suicide, parents of children with mental health problems, active soldiers and military veterans, individuals with Tourette syndrome, those with a diagnosis of dementia and mental health professionals.

### HOP's theory of change

The central premise of HOP is that supporting disclosure decisions can reduce stress associated with stigma and a fear of being found out, reduce self-stigma and empower individuals by increasing their self-efficacy in coping with stigma, thus ultimately supporting the journey toward increased well-being and recovery.<sup>1</sup> This theory of change is partly supported by randomised controlled trials of HOP published to date, conducted with adolescents in Switzerland ( $N=100$ ),<sup>3</sup> and with adults in the USA ( $N=126$ )<sup>4</sup> and Germany ( $N=98$ ).<sup>5</sup> Large reductions in stigma stress were observed in two of these trials ( $\eta^2p=0.15^3$  and  $d=0.92^5$ ) and mixed effects on the two dimensions of stigma stress (perception of stigma as harmful to oneself and perceived resources to cope with such harm) was observed in the third.<sup>4</sup> Self-stigma, a key target of the HOP programme, was reduced in two of the three studies.<sup>4,5</sup> In one of these,<sup>5</sup> initial small effects on self-stigma ( $d=0.36$ ) increased to medium effects at follow-up ( $d=0.63$ ). HOP's beneficial effects on the personally detrimental aspects of self-stigma (applying negative stereotypes to oneself and experiencing harm as a result of self-stigma) were maintained at 1-month follow-up ( $P<0.05$ ) in the other study.<sup>4</sup> The effect of HOP on self-stigma was negligible in the third study ( $\eta^2p=0.002$ ).<sup>3</sup> HOP showed large positive effects on disclosure-related distress and perceived need for secrecy at follow-up in the two studies that assessed these outcomes (both around  $\eta^2p=0.06$  in the Swiss study<sup>3</sup> and  $d=0.7$  in the German study<sup>5</sup>). Finally, help-seeking intentions increased at follow-up in the one study that examined them ( $d=0.34$  for family/friends, and  $d=0.64$  for professionals).<sup>5</sup> Thus evidence collected to date suggests that completing the HOP programme appears to leave individuals less concerned about having to conceal their experiences, less likely to apply negative

stereotypes to themselves and more likely to perceive themselves as able to deal with stigma. Of note, the outcomes of HOP vary somewhat across the studies conducted to date, and should be viewed as tentative because of the small samples involved. The impact of HOP on access to both informal and formal support requires further evaluation, as does the suggestion in one study<sup>5</sup> that HOP improves quality of life.

### Future directions

The HOP programme has proven popular with people with mental health problems and its dissemination and adaptation have developed rapidly. There are now HOP adaptations for different target groups. So far, randomised controlled trials have only been published about two of them, adults with mental health problems in community settings and adolescents with mental health problems, and most of them have been in in-patient settings. Several other trials are underway. None of the studies published to date are representative of all people with mental health problems struggling with public and self-stigma. For example, it is unknown how HOP affects individuals with mental health problems from diverse ethnic backgrounds.

Another consideration is that the assessed outcomes and associated measures in trials of HOP have varied, making comparisons across studies and samples more difficult. Hence a priority for the future implementation of HOP should be careful evaluation of outcomes and change processes. Attention will also need to be paid to ensuring that any new HOP sites and adaptations use a community-based participatory approach, which has been central to HOP's philosophy and practice from the outset.

In addition to a general need for larger, well-designed outcome and process evaluations of HOP, ideally with representative samples, a number of questions should be the focus of further research. To date, we do not know to what extent decisions in favour of disclosure or, conversely, non-disclosure mediate HOP outcomes. Feedback from HOP participants suggests that being given the opportunity to carefully consider whether, when, how and what one might want to disclose can lead to positive effects, regardless of whether the person ultimately decides to disclose. So those who decide to disclose while completing the HOP programme may experience reduced stigma stress and self-stigma, because of a reduced need for continued secrecy and concealment, whereas for those who decide not to disclose, HOP may still bring benefits through empowered non-disclosure. Future research should test these hypotheses.

In view of evidence that direct contact with persons with mental health problems is the most effective strategy to reduce stigma, one would predict that increased disclosure as a result of engaging with programmes such as HOP would ultimately contribute to a reduction in public stigma. Assessing the impact of disclosure at an interpersonal and community level may render useful evidence to support efforts to reduce stigma. Therefore, future research should ask what effect witnessing a disclosure had on people's attitudes to those experiencing mental health problems.


An important question is whether support for disclosure decision-making should be available to anyone in a stigmatised group (for example, as standard post-diagnosis), or whether they should be targeted. In this context, a stepped-care model may be worth considering, whereby anyone given a stigmatising diagnosis is offered some support to help them consider whether, how and with whom to share this, and those who are particularly concerned with secrecy and the risks of being found out have access to programmes such as HOP.

In the HOP programme, family and friends are primarily characterised as people one needs to decide whether or not to disclose to.

Going forward it may be helpful to make their potential role as allies to whom one may look for support in making and implementing disclosure decisions more explicit. However, what role they play will vary from person to person, and a key aim of HOP is to promote self-determination in deciding who is safe and helpful to disclose to. In this and other aspects, HOP's ethos of supported decision-making is very different from interventions such as CORAL, which are about shared decision-making between a professional and a stigmatised individual.

### Conclusions

Programmes such as HOP address concerns about stigma and empower participants in taking control over disclosing their difficulties. By providing a non-judgemental forum for making disclosure decisions, programmes such as HOP help affected individuals problem-solve difficult decisions and act upon these in ways that are personally meaningful.

**Katrina Scior**, Associate Professor in Clinical Psychology, Research Department of Clinical Educational & Health Psychology, University College London, UK; **Nicolas Rüsch**, Professor for Public Mental Health, Department of Psychiatry II, University of Ulm and BKH Günzburg, Germany; **Chris White**, Policy and Information Officer, Mental Health Foundation, Scotland; **Patrick W. Corrigan** , Distinguished Professor of Psychology, Lewis College of Human Sciences, Illinois Institute of Technology; and Principal Investigator, National Consortium on Stigma and Empowerment, USA

**Correspondence:** Dr Katrina Scior. Email: [k.scior@ucl.ac.uk](mailto:k.scior@ucl.ac.uk)

First received 25 Feb 2019, final revision 24 Oct 2019, accepted 2 Nov 2019

### Author contributions

All four authors made a substantial contribution to the formulation of the ideas presented in this editorial, were involved in writing the editorial, gave their final approval of the submitted version, and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### References

- 1 Corrigan PW, Kosyluk KA, Rüsch N. Reducing stigma by coming out proud. *Am J Public Health* 2013; **103**: 794–800.
- 2 Henderson C, Brohan E, Clement S, Williams P, Lassman F, Schauman O, et al. Decision aid on disclosure of mental health status to an employer: feasibility and outcomes of a randomised controlled trial. *Br J Psychiatry* 2013; **203**: 350–7.
- 3 Rüsch N, Abbruzzese E, Hagedorn E, Hartenhauer D, Kaufmann I, Curschellas J, et al. Efficacy of coming out proud to reduce stigma's impact among people with mental illness: pilot randomised controlled trial. *Br J Psychiatry* 2014; **204**: 391–7.
- 4 Corrigan PW, Larson J, Michaels PJ, Buchholz BA, del Rossi R, Fontecchio MJ, et al. Diminishing the self-stigma of mental illness by coming out proud. *Psychiatry Res* 2015; **229**: 148–54.
- 5 Mulfinger N, Müller S, Böge I, Saka V, Corrigan PW, Evans-Lacko S, et al. Honest, Open, Proud for adolescents with mental illness: pilot randomized controlled trial. *J Child Psychol Psychiatry* 2018; **59**: 684–91.