

Selecting a dietary assessment method for a national nutrition survey: a review and evaluation of online 24-hour recall tools

Berit Follong^{1,2}, Sally Mackay², Cliona Ni Mhurchu²

¹National Institute for Health Innovation, The University of Auckland, Auckland, New Zealand

²Epidemiology and Biostatistics, School of Population Health, The University of Auckland, Auckland, New Zealand

Corresponding author: Berit Follong, Email: berit.follong@auckland.ac.nz, Address: The University of Auckland, 85 Park Road, Grafton, Auckland 1023, New Zealand.

Short title: Review and evaluation of online 24-hour recall tools

Acknowledgements: Thank you to Jacqui Grey and Maria Maiquez for organizing the meetings with the expert advisory groups and the Ministries. We would also like to thank our wider investigator group for their guidance, as well as the experts who provided critical advice to help select a 24-hour dietary recall tool.



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980024002507

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Abstract

Objective: Online 24-hour dietary recall tools are commonly used in nationwide nutrition surveys to assess population diets. With a steep rise in the development of new and more advanced 24-hour recall tools, the decision of which tool to use for a national nutrition survey becomes increasingly challenging. Therefore, this short communication outlines the process of selecting a 24-hour dietary recall tool for a national nutrition survey in New Zealand.

Design: To identify suitable 24-hour dietary recall tools, a review of peer-reviewed and grey literature was conducted (2019-2022). Data on functionalities, validation, usability, and adaptability were extracted for 18 pre-specified tools, which were used in the subsequent evaluation process.

Results: Six of the 18 tools had new relevant publications since 2019. The 14 new publications described six validation studies and eight usability studies. Based on pre-selection criteria (e.g., availability, adaptability, previous use in national surveys), three tools were shortlisted: ASA24, Intake24, and MyFood24. These tools were further evaluated, and expert advice was sought to determine the most suitable tool for use in the New Zealand context.

Conclusions: A comprehensive yet time- and cost-efficient approach was undertaken to identify the potential use of online 24-hour dietary recall tools for a national nutrition survey. The selection process included key evaluation criteria to determine the tools' suitability for adaptation within the New Zealand context and ultimately to select a preferred tool. A similar approach may be useful for other countries when having to select 24-hour dietary recall tools for use in national nutrition surveys.

Introduction

National nutrition surveys collect population-based diet-related data that are essential in assessing food and nutrient intakes, monitoring nutritional status, and informing and evaluating public health nutrition programmes and policies. Given the significance of the data obtained through these national surveys, it is important that the methodologies used are robust and valid⁽¹⁾.

The most common dietary assessment method used in national nutrition surveys is the 24-hour dietary recall^(2, 3), due to its standardised data collection process, and its ability to provide reasonably accurate nutrient intake data and collect data from large population groups whilst minimising participant burden. Many traditional 24-hour dietary recall tools have undergone technological advancements in recent years to enhance their cost-efficiency, data quality, user experience, and scalability^(4, 5). Today, a wide range of online 24-hour dietary recall tools are available, each varying in their functionalities, ease of use, validity, and adaptability^(2, 6). Selecting a 24-hour dietary recall tool to collect population dietary intake data can therefore be challenging.

The methods for a future national nutrition survey in New Zealand were developed recently. Since the 24-hour dietary recall tool used in the previous national nutrition surveys was no longer available, a new tool had to be selected. This short communication describes the process and results of a review and evaluation of online 24-hour dietary recall tools for adaptation and use in a New Zealand national nutrition survey.

Methods

Updated review of existing tools

To identify existing online 24-hour dietary recall tools that could potentially be adapted for use in a New Zealand nutrition survey, a systematic review published in 2021 by Gazan and colleagues was updated⁽²⁾. The original review aimed to provide an overview of online 24-hour dietary recall tools that had the potential to collect dietary intake data for national nutrition surveys and therefore provided an ideal starting point. The authors identified 18 online 24-hour dietary recall tools in publications between 2000 and 2019. We decided to update Gazan's review for the 24-hour dietary recall tools already identified and therefore limited the updated searches to the same 18 tools. Any tools developed after the initial search

was performed would have been in an early stage of development and it was therefore less likely that validation and usability studies would have been conducted. Selection of a well-established tool was preferred over a new tool that required more testing. Furthermore, the contractor (New Zealand Ministry of Health) advised that they would prefer a 24-hour dietary recall tool that had already been used in nutrition surveys in other countries, and tool selection needed to be completed within a short timeframe to meet contract deliverables.

Search strategy, study selection, and data extraction

The search for relevant online 24-hour dietary recall tools was updated (from 2019 to January 2022) using the same methods (i.e., search strategy and eligibility criteria) described in Gazan et al.⁽²⁾. The original search terms were combined with the names of the eighteen 24-hour dietary recall tools to identify relevant publications in PubMed and Google (Scholar).

Study selection was based on publications or (scientific) literature that described the functionality, validity, user usability, or flexibility of relevant online self-administered 24-hour dietary recall tools⁽²⁾. Relevant studies were only included if published in English. A standard screening process for rapid scoping reviews was performed⁽⁷⁾ to determine eligibility. Search results were initially screened for inclusion based on title and abstract by one researcher (BF) followed by assessment by a second independent reviewer (SM) of a random sample (~20%) of excluded records only. Full-text articles were obtained and screened following the same process. Records marked as 'unsure' or where decisions were in conflict were resolved through discussion by both researchers or with the wider research team.

Data extraction was undertaken in line with the previous review and included updated information on tools' characteristics, collection methods and functionalities (e.g., dietary recall steps), and any validation or usability studies⁽²⁾. One researcher (BF) extracted and tabulated the data which were subsequently checked by a second researcher (SM). Authors and/or owners of the recall tools were contacted with requests to provide missing data or verify information where needed.

Evaluation

Shortlisting, in-depth evaluation, and consultation

Evaluation of the online 24-hour dietary recall tools was divided into three steps including

creation of a shortlist, in-depth evaluation, and expert consultation. Tools were scored against a set of pre-defined criteria to shortlist those with the highest potential for use in a New Zealand nutrition survey. These pre-selection criteria were based on criteria used for the United Kingdom (UK) National Diet and Nutrition Survey (NDNS)⁽⁸⁾ with additional criteria related to specific requirements for a New Zealand survey (Table 1). Using a stepwise approach, the first five criteria were scored (0, 1 or 2 points each) and subtotaled. Any tools that did not meet criteria 3 or 4, or scored ≤ 4 points for criteria 1 to 5 were eliminated. Remaining tools were scored for criteria 6 to 8 and shortlisted if total scores were >7 points. This cut-off score was based on advice from an expert advisory group (including 18 members with expertise in nutrition and public health) and aimed to ensure essential criteria were met before advancing to the next evaluation stage. The scoring was conducted by one researcher (BF) and checked by a second (SM). Both have a background in public health nutrition and dietary assessment methods.

[Insert Table 1]

Shortlisted tools were then assessed against a larger set of evaluation criteria from the UK NDNS selection process^(8, 9). Several criteria were added to determine the suitability of the tool for New Zealand by aligning with specific requirements for a national survey and priority ethnic groups. These added criteria were developed in consultation with expert advisors, and key representatives of the Ministries of Health and Primary Industries. The final evaluation criteria were grouped into four categories: 1) Organisational, logistical, and financial aspects (e.g., ‘How many respondents can use the tool at the same time?’); 2) Applicability (e.g., ‘Will the tool be suitable for the population groups of interest?’); 3) Respondent, interviewer, and research usability (e.g., ‘Can the tool be used offline?’); and 4) Accuracy and precision (e.g., ‘Does the tool prompt for foods commonly consumed together?’). The full set of criteria can be provided upon request.

To evaluate the shortlisted 24-hour dietary recall tools, the relevant details related to the criteria above were extracted from the updated scoping review. Additionally, relevant websites were searched, and tool developers or owners, or corresponding authors of studies were contacted. The key strengths and limitations of each tool were identified.

In-person consultation with our nutrition and public health expert advisors was undertaken to guide the decision on which tool to select and adapt for use in a nutrition survey. Where needed, further information about the tools was collected to facilitate the discussions.

Results

Updated review of existing tools

Search and screening

After removing 53 duplicates, 644 records were screened for inclusion. Forty-six records were deemed eligible for full-text screening, from which 14 new publications were included in the updated review. Six of the 18 tools originally identified by Gazan et al. had new relevant publications since 2019. These 14 publications described six studies on the Automated Self-Administered 24-hour (ASA24) dietary assessment tool (n=2 validation^(10, 11), n=3 usability^(12, 13), n=1 both validation and usability⁽¹⁴⁾), two studies on Intake24 (n=2 usability^(15, 16)), two studies on Myfood24 (n=1 validation⁽¹⁷⁾, n=1 usability⁽¹⁸⁾), two studies on R24W (n=2 validation^(19, 20)), one study on NutriNet-Salud (n=1 development⁽²¹⁾), and one study on Foodbook24 (n=1 usability⁽²²⁾). Data extracted on the development, validation, and usability of each tool is not included in this short communication but can be provided upon request.

Evaluation

Selection of shortlist

Table 2 shows the scores for each of the 24-hour dietary recall tools. Based on the scoring of the pre-selection criteria, three of the 18 tools were shortlisted: ASA24 (9/10), Intake24 (10/10), and Myfood24 (9/10).

[Insert Table 2]

Evaluation and consultation

The three shortlisted 24-hour dietary recall tools were assessed against the evaluation criteria described above. Findings and key strengths and limitations were summarised (Table 3) and presented to expert advisory groups. Across the three tools, only a few differences were observed, including the tools' offline use, use with children, and costs associated with modifications, maintenance, and research use. Expert advisors highlighted the need for offline use of the tool to allow for data collection in rural or remote communities in New Zealand where access to Wi-Fi and mobile data may be limited. Myfood24 has the option to collect data offline but only using an Android application, limiting the use in a national nutrition survey in which laptops are the most likely device used for data collection. Intake24

allows local installation of the tool on researchers' laptops to enable offline data collection with the data synced to a server at a later stage. Contact with ASA24 confirmed that it is a web-based programme only and therefore not downloadable to be completed offline, with no future plans to develop this offline functionality.

Given the nutrition survey will include participants from the age of two years, advisory group members preferred a tool that had been designed for and tested with a wide range of age groups, including younger children. All three shortlisted recall tools were created for use with people aged 10 years and older. However, evidence could only be found for adaptations to Intake24 to make it suitable for use with younger ages (1.5 years and over).

Modifications needed to create a tool suited for use in the New Zealand population would require substantial efforts and close collaboration for each shortlisted tool. Costs associated with this process, but also the maintenance and use of the dietary recall tools, however, varied significantly. Both ASA24 and Intake24 charge no licence fee, while Myfood24 has an annual licence fee and an additional fee per anticipated use of the system (e.g., based on the number of participants, recalls, and time points). Furthermore, new foods and prompts can only be added by Myfood24 and additional charges apply. Adaptation of Myfood24 was therefore more restricted and costly compared to Intake24 and ASA24.

[Insert Table 3]

Feedback from the experts was used to recommend a preferred 24-hour dietary recall tool. Representatives of the Ministries of Health and Primary Industries then reviewed the full evaluation report, including expert advice, and made a final decision on the tool to be used in a national nutrition survey. Intake24 was considered best suitable for use in a national nutrition survey for three main reasons: 1) It can be used offline; 2) It has been used or is being adapted for use in national surveys by countries similar to New Zealand, including Australia and the UK, thus providing opportunities to establish collaborations, learn from other user experiences, and compare survey findings; 3) Intake24 has been developed for and tested in both child and adult populations, which was a key requirement for a New Zealand national survey.

Conclusion

The rigorous evaluation process outlined in this short communication enabled the selection of a robust 24-hour dietary recall tool for use in a New Zealand nutrition survey. It facilitated efficient and detailed data collection to determine the key strengths and limitations of online tools including their potential to be modified for the unique population demographics in New Zealand. This approach could be used as a guide for other countries when selecting a new 24-hour dietary recall tool for use in a national nutrition survey.

List of abbreviations

NDNS National Diet and Nutrition Survey

UK United Kingdom

Financial Support: The New Zealand Ministries of Health and Primary Industries funded this research. These funders played a role in the development of the evaluation criteria and the final decision to select Intake24 as the 24-hour dietary recall tool for use in a New Zealand nutrition survey.

Conflict of Interest: None

Authorship: BF, SM, and CNM conceptualized the research design. BF conducted the database search; BF and SM completed the study selection, data extraction, and evaluation; BF wrote the manuscript; SM and CNM edited and approved the final manuscript.

Ethical Standards Disclosure: NA

References

1. Moshfegh AJ, Rhodes DG, Martin CL (2022) National Food Intake Assessment: Technologies to Advance Traditional Methods. *Annual review of nutrition* 42, 401-422.
2. Gazan R, Vieux F, Mora S *et al.* (2021) Potential of existing online 24-h dietary recall tools for national dietary surveys. *Public Health Nutrition* 24, 5361-5386.
3. Alkhaldy AA, Aljaadi AM, Jalil AMM *et al.* (2024) Cross-continental national nutrition surveys: a narrative review. *BMC Nutrition* 10, 63.
4. Ngo J, Engelen A, Molag M *et al.* (2009) A review of the use of information and communication technologies for dietary assessment. *The British journal of nutrition* 101, S102-112.
5. Murai U, Tajima R, Matsumoto M *et al.* (2023) Validation of Dietary Intake Estimated by Web-Based Dietary Assessment Methods and Usability Using Dietary Records or 24-h Dietary Recalls: A Scoping Review. *Nutrients* 15.
6. Timon CM, van den Barg R, Blain RJ *et al.* (2016) A review of the design and validation of web-and computer-based 24-h dietary recall tools. *Nutrition Research Reviews* 29, 268-280.
7. Garritty C, Gartlehner G, Nussbaumer-Streit B *et al.* (2021) Cochrane Rapid Reviews Methods Group offers evidence-informed guidance to conduct rapid reviews. *Journal of clinical epidemiology* 130, 13-22.
8. Steer T, Roberts C, Amoutzopoulos B *et al.* (2021) Selection of an automated dietary assessment tool for use in the UK National Diet and Nutrition Survey (NDNS) Rolling Programme (RP). *Proceedings of the International Conference on Diet & Activity Methods (eICDAM), Online*, 8-12.
9. Illner A, Freisling H, Boeing H *et al.* (2012) Review and evaluation of innovative technologies for measuring diet in nutritional epidemiology. *International journal of epidemiology* 41, 1187-1203.
10. Raffoul A, Hobin EP, Sacco JE *et al.* (2019) School-age children can recall some foods and beverages consumed the Prior Day Using the Automated Self-Administered 24-Hour Dietary Assessment Tool (ASA24) without Assistance. *The Journal of nutrition* 149, 1019-1026.
11. Subar AF, Potischman N, Dodd KW *et al.* (2020) Performance and Feasibility of Recalls Completed Using the Automated Self-Administered 24-Hour Dietary Assessment Tool in Relation to Other Self-Report Tools and Biomarkers in the Interactive Diet and Activity Tracking in AARP (IDATA) Study. *Journal of the Academy of Nutrition and Dietetics* 120, 1805-1820.

12. Bekelman TA, Martin CK, Johnson SL *et al.* (2022) A comparison of the remote food photography method and the automated self-administered 24-h dietary assessment tool for measuring full-day dietary intake among school-age children. *British Journal of Nutrition* 127, 1269-1278.
13. Bekelman TA, Johnson SL, Steinberg RI *et al.* (2021) A Qualitative Analysis of the Remote Food Photography Method and the Automated Self-Administered 24-hour Dietary Assessment Tool for Assessing Children's Food Intake Reported by Parent Proxy. *Journal of the Academy of Nutrition and Dietetics*.
14. Silveira SL, Jeng B, Gower BA, Motl RW (2021) Feasibility, Acceptability, and Preliminary Validity of Self-Report Dietary Assessment in Adults with Multiple Sclerosis: Comparison with Doubly Labeled Water Measured Total Energy Expenditure. *Nutrients* 13, 1198.
15. Osadchiy T, Poliakov I, Olivier P *et al.* (2020) Progressive 24-Hour Recall: Usability Study of Short Retention Intervals in Web-Based Dietary Assessment Surveys. *Journal of medical Internet research* 22, e13266.
16. Rowland M, Rose J, McLean J *et al.* (2020) Pilot of Intake24 in the Scottish Health Survey: Scotland: ScotCen Social Research.
17. Koch SA, Conrad J, Cade JE *et al.* (2021) Validation of the web-based self-administered 24-h dietary recall myfood24-Germany: comparison with a weighed dietary record and biomarkers. *European Journal of Nutrition*, 1-14.
18. Ward HA, McLellan H, Udeh-Momoh C *et al.* (2019) Use of Online Dietary Recalls among Older UK Adults: A Feasibility Study of an Online Dietary Assessment Tool. *Nutrients* 11, 1451.
19. Brassard D, Laramée C, Robitaille J *et al.* (2020) Differences in population-based dietary intake estimates obtained from an interviewer-administered and a self-administered web-based 24-h recall. *Frontiers in nutrition* 7, 137.
20. Paradis F (2021) Validation d'un outil d'évaluation alimentaire à l'aide de biomarqueurs de l'apport alimentaire.
21. Meléndez J, Hercberg S, Galán P *et al.* (2019) NutriNet-Salud Mexico. Prospective study online: 2018-2028. Relationship between nutrition and health and determinants of dietary habits and nutritional status. *Cirugía y Cirujanos (English Edition)* 86.
22. Timon CM, Walton J, Flynn A, Gibney ER (2021) Respondent Characteristics and Dietary Intake Data Collected Using Web-Based and Traditional Nutrition Surveillance Approaches: Comparison and Usability Study. *JMIR public health and surveillance* 7, e22759.

Table 1 Pre-selection criteria used to shortlist 24-hour dietary recall tools.

#	Criteria	Scoring
1	Evidence of validation of 24-hour dietary recall tool	0 points; no validation studies 1 point; 1-2 validation studies 2 points; 3 or more validation studies
2	Evidence of use in previous (national) surveys or large studies	0 points; only used in small or non-national studies/surveys 1 point; used in large studies only 2 points; used in national surveys
3	The tool is available for use	0 points; no
4	The tool is in English or can easily be adapted to English	1 point; yes
5	The tool can use images to portray portion size	
6	The tool can record (or can be easily adapted to record) additional details required to accurately code dietary data (e.g., cooking methods, brands, different nutrient profiles of foods)	
7	The tool can be updated to align with the expected requirements for a New Zealand national nutrition survey (e.g., adding or changing language, portion size images, national food lists and nutritional composition data)	
8	The tool can automatically link food/drink consumption to food composition data	

Table 2 24-hour dietary recall tools scored against the pre-selection criteria.

	Criteria #									
	1	2	3	4	5	Subtotal	6	7	8	Total
<i>Max score</i>	2	2	1	1	1	7	1	1	1	10
ASA24	2	1	1	1	1	6	1	1	1	9
CANAA-W	0	0	0	1	1	2				
CAPIS	0	0	0	0	1	1				
Clin Share	0	0	1	0	0	1				
Compleat™	1	1	1	0	0	3				
Crème Diet/Food book24	2	1	0	1	1	5				
Diet Advice	1	0	0	1	1	3				
Diet Day	1	0	0	1	1	3				
FBQ	1	1	0	1	1	4				
FoRC	1	0	0	1	1	3				
Intake24	2	2	1	1	1	7	1	1	1	10
Myfood24	2	1	1	1	1	6	1	1	1	9
NutriNet- Sante/NutriNet- Salud	1	2	1	0	1	5				
PAC24	1	0	0	0	1	2				
R24W	2	1	0	1	1	5				
RiksmatenFlex	1	2	0	0	1	4				
SACANA	1	2	0	1	1	5				
Web-SPAN	1	1	0	1	1	4				

The first five criteria were scored and subtalled. Tools that did not meet criterion 3 or 4, or scored ≤ 4 for criteria 1 to 5 were eliminated. The remaining tools were scored for criteria 6 to 8.

Criteria: 1) Evidence of validation of 24-hour dietary recall tool; 2) Evidence of use in previous (national) surveys or large studies; 3) The tool is available for use; 4) The tool is in English or can easily be adapted to English; 5) The tool can use images to portray portion size; 6) The tool can record (or can be easily adapted to record) additional details required to accurately code dietary data; 7) The tool can be updated to align with the expected requirements for a New Zealand national nutrition survey; 8) The tool can automatically link food/drink consumption to food composition data.

ASA24: Automated Self-Administered 24-hour; CANAA-W: Children's and Adolescents' Nutrition Assessment and Advice on the Web; CAPIS: Computer-Assisted Personal Interview System; FBQ: Web-based Food Behaviour Questionnaire; FoRC: Food Record Checklist; Myfood24: Measure Your Food On One Day; PAC24: Portuguese self-administered computerised 24 hour dietary recall; R24W: Web-based 24-h dietary recall; SACANA: Self-Administered Children, Adolescents, and Adult Nutrition Assessment; Web-SPAN: Web-Survey of Physical Activity and Nutrition.

Table 3 Overview of strengths and limitations for the shortlisted 24-hour dietary recall tools.

Tool	Strengths	Limitations
ASA24	<ul style="list-style-type: none"> • Uses a researcher website that lets researchers manage the survey and data which has security/confidentiality measures in place. • Tool is in English and language can be further adapted. • Tool is easy to use for participants with some issues reported that are not specifically related to ASA24 but more common to 24-hour dietary recall tools in general. • Tool is compatible with common browsers and can be used on desktops, laptops, tablets and smartphones. • Takes on average 24 minutes to complete. • Tool used in large studies to track dietary intake over time. • Well-validated tool that uses a multiple-pass format including the option to add important prompts and checks for long time gaps between meals. • No licence fee. • Can adapt food list and nutrient database. • Tool can record dietary supplement intake. • Tool can enter and save recipes. • Tool can include a follow-up survey URL. • Australian ASA24 available (older version compared to US). 	<ul style="list-style-type: none"> • Mainly designed for and tested with people from the age of 10 years old and upwards. • High costs to adapt the food list and nutrient database, and incorporate New Zealand based respondent feedback reports. • Tool cannot be used offline and use as interviewer-led tool might be limited. • Common user usability issues relate to difficulties with navigating website/logging in, finding the correct food, and portion size estimation. • Does not include a pre-check for accuracy/quality based on overall energy intakes. • Does not include warnings for unusual values/over/under consumption. • Tool does not assess usual salt intake.
Intake24	<ul style="list-style-type: none"> • Uses a researcher website that lets researchers manage the survey and data with the potential to install and provide the system locally. • Great flexibility in terms of adapting language, interface, food and nutrient databases, respondent feedback reports, including support from the Intake24 	<ul style="list-style-type: none"> • Intake24 does not include a participant request/reminder management system. This has to be provided separately. • Currently not able to use recipe function. • Costs associated with support for localisation/internationalisation.

	<p>team.</p> <ul style="list-style-type: none"> • Originally developed for people over the age of 10 years old but adaptations were also made to use the tool from the age of 1.5 years (data not published). • Validated tool that uses a multiple-pass format (less studies conducted compared to ASA24) including the option to include important prompts, checks for low energy reports, and long time gaps between meals. • Takes on average 20 minutes to complete. • Tool is easy to use for participants with some issues reported that are not specifically related to Intake24 but more common to 24-hour dietary recall tools in general (less studies conducted compared to ASA24). • Tool is compatible with common browsers and can be used on desktops, laptops, tablets and smartphones. • No licence fee. • Dietary intake data is available ongoing on the web platform. • Tool can record dietary supplement intake. • Tool can include a follow-up survey URL or external online questionnaire can embed a participant's unique URL to link to Intake24. • Used in the UK, is currently being adapted for use in Australia in their next national nutrition survey, and a New Zealand version is available. • Tool has been adapted for offline use and could be used to enter recall data collected on paper. • Tool can be used effectively by interviewers. 	<ul style="list-style-type: none"> • Common user usability issues relate to difficulties with navigating website/logging in and finding the correct food. • Does not include a warning for unusual values/over consumption. • Tool does not assess usual salt intake (adaptation could be considered).
Myfood24	<ul style="list-style-type: none"> • Uses a web interface for researchers to manage the survey. • Great flexibility in terms of adapting language, 	<ul style="list-style-type: none"> • Not validated with children under 10 years. • May be expensive as standard pricing model, annual licence fee, plus a fee per anticipated use of the system for research

	<p>interface, food databases, and respondent feedback reports including support from the Myfood24 team.</p> <ul style="list-style-type: none"> • Originally developed for people over the age of 10 years old and usability studies show appropriate for adolescents and older people. • Validated tool (against biomarkers) that uses a multiple-pass format including the option to include important prompts, forgotten foods at the end, and warning for unusually large quantities. • Takes on average 15 minutes to complete. • Tool is mostly easy to use for participants. • Can be used on common browsers, accessed via a computer, laptop, or mobile. • Can include dietary supplements (not included in nutrient data). • Tool includes a recipe builder function. • Can link to online questionnaires. • Used in large studies in the UK, an Australian version is available, and a New Zealand baby/infant food database exists. • Can be completed offline (Android device only) and can be administered by an interviewer. 	<p>use, need to contact Myfood24 for a quote.</p> <ul style="list-style-type: none"> • Some usability-related comments state that searching for a food is challenging which may be due to the underlying food list database (lack of home-made foods, lack of specific brands etc). • Only includes few prompts and can only be added for additional fee. • Foods can only be added to the food list database by the Myfood24 team and with a development fee. • Does not include warnings for underconsumption or unusual energy intake. • Tool does not assess usual salt intake.
--	--	--

Based on best available information. Where needed, information was verified with authors/developers.

ASA24: the Automated Self-Administered 24-hour; UK: United Kingdom; US: United States.