

The ropelength conjecture of alternating knots

BY YUANAN DIAO

*Dept. of Mathematics and Statistics, University of North Carolina at Charlotte, Charlotte,
9201 University City Blvd., NC 28223, U.S.A.*

(Received 03 August 2022; revised 20 March 2024; accepted 23 October 2024)

Abstract

A long standing conjecture states that the ropelength of any alternating knot is at least proportional to its crossing number. In this paper we prove that this conjecture is true. That is, there exists a constant $b_0 > 0$ such that $R(K) \geq b_0 Cr(K)$ for any alternating knot K , where $R(K)$ is the ropelength of K and $Cr(K)$ is the crossing number of K . In this paper, we prove that this conjecture is true and establish that $b_0 > 1/56$.

2020 Mathematics Subject Classification: 57K10 (Primary); 57K14 (Secondary)

1. Introduction

The ropelength of a knot K is defined as the minimum length among all knots in the ambient isotopic class of K with unit thickness [3, 8, 11]. A fundamental question in geometric knot theory asks how the ropelength $R(K)$ of a knot K is related to $Cr(K)$, the crossing number of K [1–4, 6]. In the particular case that K is an alternating knot, a long standing conjecture, well known at least to researchers who study the ropelength problem, states that the ropelength of K is at least proportional to $Cr(K)$. We prove that this conjecture is true. More specifically, we prove that there exists a constant $b_0 > 1/56$ such that $R(K) > b_0 Cr(K)$ for any alternating knot K .

2. Reverse parallel links of alternating knots

Let K be a knot in S^3 . The regular neighbourhood of K on any orientable surface M embedded in S^3 on which K lies is an annulus. Let A be such an annulus. The link formed by the boundaries K' , K'' of A is called a *reverse parallel link* of K if K' , K'' are assigned opposite orientations. The linking number f between K' and K'' when they are assigned parallel orientations is called the *framing* of K [12, 13]. The framing f is independent of the orientation of K and the ambient isotopy class of A in S^3 depends only on K and the framing. Furthermore, the reverse parallel links of K are characterised by their framing numbers. That is, two reverse parallel links of K are ambient isotopic if and only if they have the same linking number. We shall denote a reverse parallel link of K with framing f by \mathbb{K}_f following [9].

Let K be an alternating knot with a reduced diagram D . Let $c(D)$ be the number of crossings in D and $w(D)$ the writhe of D . It is known that $c(D) = Cr(K)$ is the crossing number

of K . Also, all crossings of D are positive with respect to one checkerboard shading of D and are all negative with respect to the other checkerboard shading of D . Let $v_+(D)$ be the number of regions in the shading with respect to which all crossings are positive, and $v_-(D)$ be the number of regions in the complementary shading with respect to which all crossings are negative, then $v_+(D) + v_-(D) = c(D) + 2$ [10]. The following results provide the key to solving the ropelength conjecture for alternating knots.

THEOREM 2.1. [9] *Let K be an alternating knot and D a reduced diagram of K . Let $c(D)$, $w(D)$, $v_+(D)$ and $v_-(D)$ be as defined above, then the braid index of \mathbb{K}_f , denoted by $\mathbf{b}(\mathbb{K}_f)$, is given by the following formula:*

$$\mathbf{b}(\mathbb{K}_f) = \begin{cases} c(D) + 2 + a(D) - f, & \text{if } f < a(D), \\ c(D) + 2, & \text{if } a(D) \leq f \leq b(D), \\ c(D) + 2 - b(D) + f, & \text{if } f > b(D), \end{cases}$$

where $a(D) = -v_-(D) + w(D)$ and $b(D) = v_+(D) + w(D)$. In particular, $\mathbf{b}(\mathbb{K}_f) \geq c(D) + 2$ for any f .

THEOREM 2.2. [5] *Let L be an oriented link and $\mathbf{b}(L)$ be its braid index. Then $\ell(L_c) \geq \mathbf{b}(L)$ where $\ell(L_c)$ is the minimum length among all links in the ambient isotopic class of L that are embedded in the cubic lattice of \mathbb{R}^3 .*

3. The ropelengths of alternating knots

THEOREM 3.1. *There exists a constant $b_0 > 1/56$ such that for any alternating knot K , $R(K) \geq b_0 Cr(K)$.*

Proof. Let K_c be a knot on the cubic lattice that is ambient isotopic to K and has the minimum length among all lattice knots in the ambient isotopic class of K . Let $\ell(K_c)$ be the length of K_c , then we have $R(K) > (1/14)\ell(K_c)$ as a consequence from the proof of [7, lemma 1].

Now, the set $\{\mathbf{x} + t((1/2, 1/2, 1/2): \mathbf{x} \in K_c, 0 \leq t \leq 1\}$ is an embedding of the annulus $\{1 \leq x^2 + y^2 \leq 2: x, y \in \mathbb{R}\}$ into \mathbb{R}^3 with K_c and $K'_c = ((1/2, 1/2, 1/2)) + K_c$ as its boundary curves. Assigning K_c and K'_c opposite orientations yields a reverse parallel link L_c of K . Let \mathbb{K}_c and \mathbb{K}'_c be the lattice knots obtained from K_c and K'_c by scaling up with a scaling factor of 2, and let \mathbb{L}_c be the link formed by them. Notice that $\ell(\mathbb{L}_c) = 4\ell(K_c)$. By Theorem 2.1, we have $\mathbf{b}(\mathbb{L}_c) \geq Cr(K) + 2$. Then by Theorem 2.2, we have $\ell(K_c) \geq (Cr(K) + 2)/4$. Therefore $R(K) > (1/14)\ell(K_c) \geq (Cr(K) + 2)/56$.

Remark 3.2. The construction in the proof of [7, lemma 1] can in fact be improved to yield a better bound $R(K) > (1/12)\ell(K_c)$. This would allow us to improve the estimation of the constant b_0 to be at least $1/48$. It is certainly possible to further improve b_0 by not using the lattice knots as a mid step.

Remark 3.3. Theorem 2.1 only applies to alternating knots, therefore the ropelength conjecture remains open in general for alternating links with two or more components.

Acknowledgement. The author is grateful to Hugh Morton for the key development of the theory in our joint work [9].

REFERENCES

- [1] G. BUCK. Four-thirds power law for knots and links. *Nature* **392** (1998), 238–239.
- [2] G. BUCK and J. SIMON. Thickness and crossing number of knots. *Topology Appl.* **91** (3) (1999), 245–257.
- [3] J. CANTARELLA, R. KUSNER and J. SULLIVAN. On the minimum ropelength of knots and links. *Invent. Math.* **150** (2002), 257–286.
- [4] E. DENNE, Y. DIAO and J. SULLIVAN. Quadrisecants give new lower bounds for the ropelength of a knot. *Geom. Topol.* **10** (2006), 1–26.
- [5] Y. DIAO. Braid Index Bounds Ropelength from Below. *J. Knot Theory Ramifications*, **29** (4) (2020), 2050019.
- [6] Y. DIAO, C. ERNST, A. POR and U. ZIEGLER. The ropelengths of knots are almost linear in terms of their crossing numbers. *J. Knot Theory Ramifications*, **28** (14), 1950085, 2019. DOI: [10.1142/S0218216519500858](https://doi.org/10.1142/S0218216519500858).
- [7] Y. DIAO, C. ERNST and E. J. JANCE VAN RENSBURG. Upper Bounds on Linking Number of Thick Links. *J. Knot Theory Ramifications* **11**(2) (2002), 199–210.
- [8] Y. DIAO, C. ERNST and E. J. JANCE VAN RENSBURG. Thicknesses of knots. *Math. Proc. Camb. Phil. Soc.* **126** (2) (1999), 293–310.
- [9] Y. DIAO and H. MORTON. The Braid Indices of the Reverse Parallel Links of Alternating Knots. *Algebr. Geom. Topol.*, **24**(5) (2024), 2957–2970.
- [10] L. H. KAUFFMAN. State models and the Jones polynomial. *Topology* **26**(3) (1987), 395–407.
- [11] R. LITHERLAND, J. SIMON, O. DURUMERIC and E. RAWDON. Thickness of knots. *Topology Appl.* **91** (1999), 233–244.
- [12] I. J. NUTT. Embedding knots and links in an open book III. On the braid index of satellite links. *Math. Proc. Camb. Phil. Soc.* **126** (1999), 77–98.
- [13] L. RUDOLPH. A congruence between link polynomials. *Math. Proc. Camb. Phil. Soc.* **107** (1990), 319–327.