

Short Communication

The importance of selectively logged forests for tiger *Panthera tigris* conservation: a population density estimate in Peninsular Malaysia

D. MARK RAYAN and SHARIFF WAN MOHAMAD

Abstract To obtain information on density of tiger *Panthera tigris* in selectively logged forest, a 9-month camera-trapping survey was conducted over elevations of 190–850 m in lowland and hill/upper dipterocarp and lower montane forests in Gunung Basor Forest Reserve, Peninsular Malaysia. Capture-recapture sampling methods were used to estimate tiger population density in the Reserve. The number of individual tigers captured was six. Using the M_h jackknife estimator the average capture probability per sampling occasion was 0.28 and the corresponding estimate of population size $8 \pm \text{SE } 1.89$. The overall probability of photo-capturing a tiger present in the sampled area was 0.75. Using an approach based on distances between photo-captures, a buffer width of 3.22 km and an effectively sampled area of 308 km² was estimated. This resulted in a density estimate of $2.59 \pm \text{SE } 0.71$ adult tigers per 100 km². The results indicate that selectively logged forests such as Gunung Basor Forest Reserve have the potential to accommodate a high density of tigers. Decision makers and conservation planners should not therefore perceive selectively logged forests to have limited conservation value. Further research on the ecology of tigers and their prey in selectively logged forests is urgently needed. Such research would enable conservationists to recommend tiger-friendly management guidelines for sustainable forest management and thereby significantly contribute to tiger conservation in Malaysia.

Keywords Camera-trap, capture-recapture, forest reserve, Malaysia, *Panthera tigris*, selective logging, tiger.

It has been estimated that there are c. 500 wild tigers *Panthera tigris* remaining in Peninsular Malaysia (Kawanishi et al., 2003). Although 45% of Malaysia is still forested (Kawanishi et al., 2003), the country's apex predator is gravely threatened by habitat loss, forest fragmentation, prey depletion, poaching and retaliatory killing (Locke, 1954; Elagupillay, 1983; Kawanishi et al., 2003, 2006). To assess the status of tigers camera-trapping surveys at

nine sites were conducted over 1997–1999 (Lynam et al., 2007) but only naive density estimates (0.51–1.95 100 km⁻²) could be derived because of insufficient data for capture-recapture analysis.

Apart from a study conducted in primary forest in Taman Negara National Park, Peninsular Malaysia, over 1999–2001 (Kawanishi & Sunquist, 2004), robust density estimates of tigers are not available for other forest types in Malaysia. There is therefore an urgent need to obtain information on the density of tigers in other forest types to formulate an effective national tiger conservation strategy. To provide such information for a disturbed habitat we adapted the capture-recapture framework (Nichols & Karanth, 2002) to determine the population density of tigers in a selectively logged dipterocarp forest in Peninsular Malaysia.

This study was conducted in Gunung Basor Forest Reserve, in Jeli District, in north-east Peninsular Malaysia in the state of Kelantan. This Reserve is part of a contiguous landscape that falls under a Class 1 Tiger Conservation Landscape (Dinerstein et al., 2006). The area is undulating (150–1,840 m) with floristic zones ranging from lowland dipterocarp and hill/upper dipterocarp forest to lower montane forest. The dipterocarp forest of this forest reserve has been selectively logged on several occasions since the 1970s and more recently during 2004–2006.

Following reconnaissance surveys for tiger signs (September–October 2004) a total of 15 camera-trap locations were selected encompassing an area of c. 120 km². Camera-trapping, with passive infra-red camera-traps (CamTrakker, Georgia, USA), was conducted between October 2004 and July 2005. All camera-traps were operational for 24 hours per day. Each camera-trap location had two camera-traps positioned on opposite sides of a trail to photograph both flanks of any tiger simultaneously (Karanth & Nichols, 2002). Camera-trap locations were at elevations of 190–850 m, and distance between locations was 1.8–6.0 km.

Capture-recapture methods (Nichols & Karanth, 2002) were used to estimate total abundance. A closure test was conducted with the software CAPTURE (Otis et al., 1978; Rexstad & Burnham, 1991) to investigate whether the closed population assumption was violated. The jackknife estimator (Otis et al., 1978) under the heterogeneity model (M_h) was used to estimate population size because it has performed

D. MARK RAYAN (Corresponding author) and SHARIFF WAN MOHAMAD
WWF-Malaysia, 49, Jalan SS23/15, Taman SEA, 47400 Petaling Jaya, Selangor,
Malaysia. E-mail mdarmaraj@wwf.org.my

Received 24 October 2007. Revision requested 3 December 2007.
Accepted 3 April 2008.

well in other tiger camera-trap studies (Karanth, 1995; Karanth & Nichols, 1998; Karanth et al., 2004).

Trap effort during each trapping session was not equal because of instances of camera-trap malfunction and camera-trap damage by elephants. A sampling occasion was therefore defined according to each monthly trapping period (Karanth, 1995). There was no loss in the detection of tigers by conforming to this definition of sampling occasion. Density estimates were generated by dividing tiger abundance (\hat{N} , from *CAPTURE*) by the effectively sampled area, $\hat{A}(\hat{W})$. The effectively sampled area was estimated using the strip width buffer method (Wilson & Anderson, 1985; Karanth & Nichols, 1998).

The statistical test for population closure in *CAPTURE* supported the assumption that the sampled population was closed for the study period ($z = -0.629$, $P = 0.26$). The number of individual tigers captured (M_{t+i}) was six (Table 1). Using the M_h jackknife estimator the average capture probability per sampling occasion (\hat{p}) was 0.2812 and the corresponding estimate of population size ($\hat{N}(\hat{SE}(\hat{N}))$) was 8(1.89). Thus, the overall probability of photo-capturing a tiger in the sampled area (M_{t+1}/\hat{N}) was 0.75. The

TABLE 1 Capture histories of the six individual adult tigers camera-trapped from December 2004 to July 2005, and the number of trap nights per month (i.e. the sampling effort).

	No. of occasions							
	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July
Male 1	1	1	1	1	0	1	1	1
Male 2	1	0	0	0	0	0	0	0
Male 3	0	0	0	0	0	0	1	0
Female 1	1	1	0	1	0	0	0	0
Female 2	1	1	0	0	1	0	1	0
Female 3	0	0	0	0	0	1	0	1
No. of trap nights	241	236	411	418	398	372	346	74

polygon formed by the outermost camera-traps was 122.9 km² (Fig. 1) with a buffer width (\hat{W}) of 3.22 km and an estimated effectively sampled area $\hat{A}(\hat{W})$ of 308 km². The estimated density ($\hat{D}(\hat{SE})$) was 2.59 (0.71) adult tigers per 100 km².

These results show that the selectively logged forests of Gunung Basor Forest Reserve contain a population density




FIG. 1 (A) The state of Kelantan in the north-east of Peninsular Malaysia (shaded), with the location of Jeli District indicated in the west of the state, (B) the Tiger Conservation Landscape of Jeli, with Gunung Basor Forest Reserve indicated, and (C) the study area in Gunung Basor Forest Reserve, with the study area, effectively sampled area, and excluded areas (orchards, plantations and human settlements).

TABLE 2 Comparison of capture-recapture statistics used in deriving density estimates in this and Kawanishi & Sunquist's (2004) study in Peninsular Malaysia.

Study site	Kawanishi & Sunquist (2004)			
	Merapoh	Kuala Terenggan	Kuala Koh	This study
Number of months, T	9	9	9	8
Closure test, P	0.19	0.94	0.073	0.26
Selection criteria under model M_h	0.86	1.00	1.00	0.82
Number of animals captured, M_{t+1}	5	5	6	6
Estimated average capture probability per sampling occasion, \hat{p}	0.13	0.16	0.22	0.28
Estimated capture probability over all sampling occasions, (M_{t+1}/\hat{N})	0.71	1.00	1.00	0.75
Population estimate \pm SE, $\hat{N}(\widehat{SE}(\hat{N}))$	7 \pm 1.92	5 \pm 2.35	6 \pm 2.44	8 \pm 1.89
95% confidence interval of estimate	6–14	5–20	6–21	7–15
Study area (km ²)	165.47	164.20	151.41	122.90
Boundary width (km)	3.03 ¹	4.11 ¹	2.59 ¹	3.22 ²
Effectively sampled area (km ²)	353.60	452.50	317.24	308.00
Estimated tiger density \pm SE (km ⁻²)	1.98 \pm 0.54	1.10 \pm 0.52	1.89 \pm 0.77	2.59 \pm 0.71

¹Boundary width calculated using the Absolute Maximum Distance Moved method

²Boundary width calculated using the Mean Maximum Distance Moved method

of tigers c. 30% higher than the highest estimate derived by Kawanishi & Sunquist (2004) in Taman Negara National Park, a protected primary forest (Table 2). In other protected primary forests in South-east Asia estimated tiger densities are 1–4 per 100 km² (Indonesia: O'Brien et al., 2003; Thailand: Simcharoen et al., 2007). Although logged forests are disturbed habitats, the response of tigers to the direct and indirect impacts of logging is poorly known. Selective logging may actually improve tiger habitat (Miquelle et al., 1999) as the disruption of the forest canopy increases sunlight to the forest floor and thus increases browse availability to tiger prey (Davies et al., 2001).

We did not conduct transect sampling to quantify absolute abundance of prey because this method does not provide sufficient sample sizes to estimate prey abundance (Kawanishi, 2002). Nevertheless, relative abundance indices based on photo encounter rates of two prey species (barking deer *Muntiacus muntjak* and wild boar *Sus scrofa*) appear to support a predictive model of tiger abundance as a function of prey (Karanth et al., 2004) as the indices of these two species are at least three and six times higher, respectively, than that of tigers in Gunung Basor Forest Reserve (Darmaraj, 2007). Reliance on cattle depredation is unlikely to account for the reserve's high population density of tigers because an average of only eight cases of such predation per year over 1993–2003 was recorded within the entire state of Kelantan (Badrul, 2003).

Our results illustrate the potential of selectively logged forests to accommodate a high population density of tigers. There is a tendency in Malaysia to perceive selectively logged-over forest as having limited conservation and economic value; this fallacy has probably led to the degazetting of forest

reserves and subsequent conversion to other land uses (e.g. for plant commodity crops such as oil palm). As tigers have large habitat requirements the effects of such conversion, leading to fragmentation and isolation of forest reserves, will severely affect the long-term viability of tiger populations across the landscape. Our results demonstrate the need for further research on tiger ecology in selectively logged forests to inform decision makers and conservation planners of the conservation value of such habitats. We hope that future research will highlight the role of selectively logged forests for tiger conservation and aid in providing tiger-friendly management guidelines for sustainable forest management in Malaysia.

Acknowledgements

We thank WWF-Netherlands for financial assistance and the Department of Wildlife and National Parks and the Forestry Department of Kelantan for their support. We are also indebted to Kae Kawanishi, Ullas Karanth, WWF-Malaysia colleagues and others for their valuable input.

References

- BADRUL, A.M.S. (2003) The trend of tiger-human conflicts in Peninsular Malaysia from 1991 to 2003. *Journal of Wildlife and Parks*, 21, 103–109.
- DARMAJAJ, M.R. (2007) *Tiger Monitoring Study in Gunung Basor Forest Reserve, Jeli, Kelantan*. Unpublished Report. WWF-Malaysia, Petaling Jaya, Malaysia.
- DAVIES, G., HEYDON, M., LEADER-WILLIAMS, N., MACKINNON, J. & NEWING, H. (2001) The effects of logging on tropical forest ungulates. In *The Cutting Edge—Conserving Wildlife in Logged*

- Tropical Forest* (eds R.A. Fimbel, A. Grajal & J.G. Robinson), pp. 93–24. Columbia University Press, New York, USA.
- DINERSTEIN, E., LOUCKS, C., HEYDLAUFF, A., WIKRAMANAYAKE, E., BRYJA, G., FORREST, J. et al. (2006) *Setting Priorities for the Conservation and Recovery of Wild Tigers: 2005–2015: A User's Guide*. WWF, WCS, Smithsonian, and NFWF-STF, Washington, DC, & New York, USA.
- ELAGUPILLAY, S. (1983) Livestock depredation problem by tigers in cattle farms. *Journal of Wildlife and Parks*, 2, 136–144.
- KARANTH, K.U. (1995) Estimating tiger *Panthera tigris* populations from camera-trap data using capture-recapture models. *Biological Conservation*, 71, 333–338.
- KARANTH, K.U., CHUNDAWAT, R.S., NICHOLS, J.D. & KUMAR, N.S. (2004) Estimation of tiger densities in the tropical dry forests of Panna, Central India, using photographic capture-recapture sampling. *Animal Conservation*, 7, 285–290.
- KARANTH, K.U. & NICHOLS, J.D. (1998) Estimation of tiger densities in India using photographic captures and recaptures. *Ecology*, 79, 2852–2862.
- KARANTH, K.U. & NICHOLS, J.D. (eds) (2002) *Monitoring Tigers and their Prey: A Manual for Researchers, Managers and Conservationists in Tropical Asia*. Centre for Wildlife Studies, Bangalore, India.
- KARANTH, K.U., NICHOLS, J.D., KUMAR, N.S., LINK, W.A. & HINES, J.E. (2004) Tigers and their prey: predicting carnivore densities from prey abundance. *Proceedings of the National Academy of Sciences of the USA*, 101, 4854–4858.
- KAWANISHI, K. (2002) *Population status of tigers (Panthera tigris) in a primary rainforest of Peninsular Malaysia*. PhD thesis, University of Florida, Gainesville, USA.
- KAWANISHI, K., SITI, H.Y., ABDUL KADIR, A.H. & TOPANI, R. (2003) Distribution and potential population size of the tiger in Peninsular Malaysia. *Journal of Wildlife and Parks*, 21, 29–50.
- KAWANISHI, K. & SUNQUIST, M.E. (2004) Conservation status of tigers in a primary rainforest of Peninsular Malaysia. *Biological Conservation*, 120, 329–344.
- KAWANISHI, K., SUNQUIST, M.E. & JASMI, A. (2006) Effects of human activities on the tiger-prey community in Taman Negara National Park, Peninsular Malaysia and their implications for conservation. In *Management and Status of Resources in Protected Areas of Peninsular Malaysia* (eds O. Sahir, Y. Siti Hawa, S. Elagupillay, M.N. Shukor, A. Norhayati & A.M.S. Shahrul), pp. 175–198. Department of Wildlife and National Parks, Kuala Lumpur, Malaysia.
- LOCKE, A. (1954) *The Tigers of Terengganu*. Museum Press Ltd, London, UK.
- LYNAM, A.J., LAIDLAW, R., WAN SHAHARUDDIN, W.N., ELAGUPILLAY, S. & BENNETT, E.L. (2007) Assessing the conservation status of the tiger *Panthera tigris* at priority sites in Peninsular Malaysia. *Oryx*, 41, 454–462.
- MIQUELLE, D.G., SMIRNOV, E.N., MERRILL, T.W., MYSLENKOV, A.E., QUIGLEY, H.B., HORNOCKER, M.G. & SCHLEYER, B. (1999) Hierarchical spatial analysis of Amur tiger relationships to habitat and prey. In *Riding the Tiger: Tiger Conservation in Human-dominated Landscapes* (eds J. Seidensticker, S. Christie & P. Jackson), pp. 71–99. Cambridge University Press, Cambridge, UK.
- NICHOLS, J.D. & KARANTH, K.U. (2002) Statistical concepts: estimating absolute densities of tigers using capture-recapture sampling. In *Monitoring Tigers and their Prey: A Manual for Researchers, Managers and Conservationists in Tropical Asia* (eds K.U. Karanth & J.D. Nichols), pp. 121–137. Centre for Wildlife Studies, Bangalore, India.
- O'BRIEN, T.G., KINNAIRD, M.F. & WIBISONO, H.T. (2003) Crouching tigers, hidden prey: Sumatran tiger and prey populations in a tropical forest landscape. *Animal Conservation*, 6, 131–139.
- OTIS, D.L., BURNHAM, K.P., WHITE, G.C. & ANDERSON, D.R. (1978) Statistical inference from capture data on closed animal populations. *Wildlife Monographs*, 62, 1–135.
- REXSTAD, E. & BURNHAM, K.P. (1991) *User's Guide for Interactive Program CAPTURE. Abundance Estimation of Closed Animal Populations*. Colorado State University, Fort Collins, USA.
- SIMCHAROEN, S., PATTANAVIBOOL, A., KARANTH, K.U., NICHOLS, J.D. & KUMAR, N.S. (2007) How many tigers *Panthera tigris* are there in Huai Kha Khaeng Wildlife Sanctuary, Thailand? An estimate using photographic capture-recapture sampling. *Oryx*, 41, 447–453.
- WILSON, K.R. & ANDERSON, D.R. (1985) Evaluation of two density estimators of small mammal population size. *Journal of Mammalogy*, 66, 13–21.

Biographical sketches

MARK RAYAN DARMARAJ is a field biologist who has been working for WWF-Malaysia for the past 5 years. He currently designs and conducts wildlife research in several selectively logged forests in Peninsular Malaysia using camera traps. His particular interests are the population ecology of large mammals, especially tigers. WAN MOHAMAD SHARIFF, also with WWF-Malaysia, is continuing research on the ecology of tigers using camera traps. His other interests are the ecology of medium to large terrestrial mammals, especially other species of felid.