

**Aims.** To ensure that service users in in-patient secure services have prolactin, vitamin D and HbA1c monitoring as per current best practice guidance.

**Background.** Service users prescribed antipsychotic medication are at risk of developing raised prolactin levels and metabolic syndrome. In both sexes, long-standing hyperprolactinaemia can lead to low bone mineral density with an increased risk of developing osteoporosis.

In recent years there has been increasing controversy on the increase in Vitamin D monitoring despite the poor evidence for complications from vitamin D deficiency in adults. Not undertaking this test in the absence of symptoms will potentially reduce anxiety for service users could save £17 per test and £50 for a 12-week course of Vitamin D supplementation. Local and national guidance indicate Vitamin D monitoring should only be done in symptomatic people.

**Method.** Fifty-five service users in the five in-patient wards had their electronic records and pathology results reviewed over a one-year period. All service users were expected to have a minimum of an annual HbA1c and prolactin level but to only have vitamin D monitoring if symptomatic for deficiency.

**Result.** Although 100% of service users in MSU were tested, vitamin D testing was consistently undertaken without documented clinical evidence of deficiency. The ranges across all units were: prolactin (72–1384 mU/L), HbA1c (30–90 mmol/mol) and vitamin D (15–124 nmol/L). Local reference ranges are prolactin (53–360 mU/L), HbA1c (<48 mmol/mol) and Vitamin D (50–120 nmol/L).

Prolactin levels were highest on the male medium secure wards.

The other two units had significantly less testing with prolactin and HbA1c levels being the least measured (18% of service users on male LSU and 23% on the female ward respectively). Vitamin D testing on these two wards were 38% on the female ward and 18% on the male ward for both tests.

**Conclusion.** Northside House has a dedicated physical health team and this is likely to explain its 100% score. However, vitamin D testing was being undertaken automatically rather than based on symptoms.

The recommendation is to add prolactin and HbA1c to the physical screens done before CPA meetings for all service users prescribed an antipsychotic but to stop Vitamin D testing in the absence of clinical symptoms of vitamin D deficiency.

### Advanced nurse practitioners- the missing link on old age psychiatry inpatient wards?

Grace Lydia Goss<sup>1\*</sup>, Priya Gowda<sup>2</sup> and Danika Rafferty<sup>2</sup>

<sup>1</sup>Aneurin Bevan University Health Board and <sup>2</sup>Cwm Taf Morgannwg University Health Board

\*Corresponding author.

doi: 10.1192/bjo.2021.869

**Aims.** 1. This project aimed to review the medical cover available to an Old Age Psychiatry inpatient ward.

2. To discuss with ward staff their view for potential improvements and areas of clinical development.

3. To review the potential of a Full Time Nurse Practitioner role on the ward.

One junior doctor (CT1 or equivalent) was allocated to cover the ward whilst balancing their other training needs and clinical commitments. The inpatient ward was based in a community hospital with no onsite medical team. The patients mostly had complex medical needs and multiple comorbidities.

**Method.** The Junior Doctor's timetable and the time allocated to the ward was reviewed. Questionnaires were conducted with nursing staff to assess their views on the support of physical health cover. The patient notes were analysed for the time taken to review patients after falls over a one month period.

**Result.** There were 14.5 hours allocated to ward cover. An additional 4 hours was provided by another visiting junior doctor totalling 18.5 hours per week- 11% of the time. This figure does not account for annual leave, on call commitments or study days whereby there was no additional cover.

A short survey completed by ward staff showed- (1 = Very Poor/Difficult 5 = Excellent/Easy)

- They rated medical cover of physical health needs on ward 7 as 1.3.
- They found contacting a Doctor to discuss a physical problem as 1.7- with particular concern for OOH.
- It was rated to be extremely difficult for a same day review of physical health problems- 1.7
- It was rated extremely difficult to get a physical review following a fall on ward 7- 1.4
- Continuity of care for the patients on ward 7 was rated as 1.6.

The patient case files reviewed over a one month period showed x8 falls. These took on average 14 hours before having a review.

**Conclusion.** Medical cover for the old age psychiatry inpatient ward was inconsistent and a challenge for a single trainee to manage alongside their other clinical commitments and training needs. A case was proposed to management with an SBAR for a Full Time Advanced Nurse Practitioner which has been approved. This role should provide patients with appropriate cover of their physical health needs. It will allow the junior doctor to work alongside them on the ward supporting each other to provide optimal care for the inpatients.

### Audit on venous thromboembolism risk assessment in mental health inpatient wards

Rahul Malhotra

Bryn Y Neuadd Hospital, Betsi Cadwaladr University Health Board

doi: 10.1192/bjo.2021.870

**Aims.** From May 2015 NHS organisations in Wales are expected to report the number of VTE cases associated with hospital admissions which are possible hospital acquired thrombosis (HAT) per calendar month. NICE Quality Standards (QS3) recommend that All patients, on admission, receive an assessment of VTE and bleeding risk using the clinical risk assessment criteria described in the national tool.

**Background.** VTE is a condition in which a blood clot (thrombus) forms in a vein, most commonly in the deep veins of the legs, known as a deep vein thrombosis (DVT). The thrombus can dislodge from its original site and travel in the blood (embolism). If it becomes lodged in the lungs, a condition known as a pulmonary embolism (PE) arises and can cause sudden death. Hospital acquired thrombosis is avoidable and unfortunately kills patients under our care.

**Method.** Collected data using a standardised form for 131 patients from 3 inpatient mental health units on documentation of a VTE risk assessment in the inpatient notes. For those patients who had a documented risk assessment, further data were collected on documentation of contraindications, risk factors, sign and date of prescription and the appropriateness of prescribing.

**Conclusion.** 8% of patients from one mental health unit (n = 48) had a documented risk assessment in the notes. The subsections of