



European First-Episode Schizophrenia Network

EDITED BY SHÔN LEWIS

EDITORIAL

- s1 The European First-Episode Schizophrenia Network**
S. Lewis
-

PAPERS

- s3 First-episode schizophrenia, bipolar disorder and other psychoses in a rural Irish catchment area: incidence and gender in the Cavan–Monaghan study at 5 years**
P. J. Scully, J. F. Quinn, M. G. Morgan, A. Kinsella, E. O'Callaghan, J. M. Owens and J. L. Waddington
-
- s10 Independent course of childhood auditory hallucinations: a sequential 3-year follow-up study**
S. Escher, M. Romme, A. Buiks, P. Delespaul and J. van Os
-
- s19 Family history of psychiatric disorders and age at first contact in schizophrenia: an epidemiological study**
M. Byrne, E. Agerbo and P. B. Mortensen
-
- s26 Distinguishing characteristics of subjects with good and poor early outcome in the Edinburgh High-Risk Study**
E. C. Johnstone, R. Cosway and S. M. Lawrie
-
- s30 Subjective and objective neuropsychological abnormalities in a psychosis prodrome clinic**
M. Hambrecht, M. Lammertink, J. Klosterkötter, E. Matuschek and R. Pukrop
-
- s38 Executive dysfunction in first-episode schizophrenia and relationship to duration of untreated psychosis: the West London Study**
E. Joyce, S. Hutton, S. Mutsatsa, H. Gibbins, E. Webb, S. Paul, T. Robbins and T. Barnes
-
- s45 Improvement in cognitive functioning in patients with first-episode psychosis during treatment with quetiapine: an interim analysis**
K. P. Good, I. Kiss, C. Buiteman, H. Woodley, Q. Rui, D. Whitehorn and L. Kopala
-
- s50 Neurological soft signs in first-episode psychosis: a systematic review**
P. Dazzan and R. M. Murray
-
- s58 Structural magnetic resonance imaging in patients with first-episode schizophrenia, psychotic and severe non-psychotic depression and healthy controls. Results of the Schizophrenia and Affective Psychoses (SAP) project**
R. K. R. Salokangas, T. Cannon, T. van Erp, T. Ilonen, T. Taiminen, H. Karlsson, H. Lauerma, K.-M. Leinonen, E. Wallenius, A. Kaljonen, E. Syvälahti, H. Vilkkman, A. Alanen and J. Hietala
-
- s66 Brain morphology in antipsychotic-naïve schizophrenia: a study of multiple brain structures**
W. Cahn, H. E. Hulshoff Pol, M. Bongers, H. G. Schnack, R. C. W. Mandl, N. E. M. van Haren, S. Durston, H. Koning, J. A. van der Linden and R. S. Kahn
-
- s73 Duration of untreated psychosis and pathways to care in first-episode psychosis. Investigation of help-seeking behaviour in primary care**
A. Skeate, C. Jackson, M. Birchwood and C. Jones
-
- s78 Randomised controlled trial of early detection and cognitive therapy for preventing transition to psychosis in high-risk individuals. Study design and interim analysis of transition rate and psychological risk factors**
A. P. Morrison, R. P. Bentall, P. French, L. Walford, A. Kilcommons, A. Knight, M. Kreutz and S. W. Lewis
-
- s85 Neurocognitive dimensions characterising patients with first-episode psychosis**
S. Friis, K. Sundet, B. R. Rund, P. Vaglum and T. H. McGlashan
-
- s91 Randomised controlled trial of cognitive-behavioural therapy in early schizophrenia: acute-phase outcomes**
S. Lewis, N. Tarrrier, G. Haddock, R. Bentall, P. Kinderman, D. Kingdon, R. Siddle, R. Drake, J. Everitt, K. Leadley, A. Benn, K. Grazebrook, C. Haley, S. Akhtar, L. Davies, S. Palmer, B. Faragher and G. Dunn
-
- s98 OPUS study: suicidal behaviour, suicidal ideation and hopelessness among patients with first-episode psychosis. One-year follow-up of a randomised controlled trial**
M. Nordentoft, P. Jeppesen, M. Abel, P. Kassow, L. Petersen, A. Thorup, G. Krarup, R. Hemmingsen and P. Jørgensen
-