

Which characteristics of nursing home residents influence differences in malnutrition prevalence? An international comparison of The Netherlands, Germany and Austria

Noémi C. van Nie-Visser^{1*}, Judith Meijers¹, Jos Schols^{1,2}, Christa Lohrmann³, Sabine Bartholomeyczik⁴, Marieke Spreuwenberg¹ and Ruud Halfens¹

¹Department of Health Services Research, Faculty of Health, Medicine and Life Sciences, School for Public Health and Primary Care (CAPHRI), Maastricht University, PO Box 616, 6200 MD, Maastricht, The Netherlands

²Department of General Practice, Faculty of Health, Medicine and Life Sciences, School for Public Health and Primary Care (CAPHRI), Maastricht University, PO Box 616, 6200 MD, Maastricht, The Netherlands

³Institute of Nursing Science, Medical University Graz, Billrothgasse 6, 8010 Graz, Austria

⁴Institute of Nursing Science, University Witten/Herdecke, Stokumerstraße 12, Witten, Germany

(Submitted 14 December 2012 – Final revision received 22 August 2013 – Accepted 24 September 2013 – First published online 18 November 2013)

Abstract

Prevalence rates of malnutrition vary considerably internationally, partly due to differences in measurement methodology and instruments. In the present study, the same measurement methodology and instruments were used in The Netherlands, Germany and Austria. The aim of the present study was to investigate whether resident characteristics influence possible differences in malnutrition prevalence between countries. The study followed a cross-sectional, multi-centre design that measured malnutrition in nursing home residents from The Netherlands, Germany and Austria. Resident data were gathered using a standardised questionnaire. Malnutrition was operationalised using BMI, unintentional weight loss and nutritional intake. Data were analysed using an association model. The prevalence rates of malnutrition in The Netherlands, Germany and Austria were 18.3, 20.1 and 22.5%, respectively. The multivariate generalised estimating equation (GEE) logistic regression analysis showed that sex, age, care dependency, the mean number of diseases and some specific diseases were influencing factors for whether the resident was malnourished or not. The OR of malnutrition in the three countries declined after including the influencing factors resulting from the multivariate GEE analysis. The present study reveals that differences in the prevalence rates of malnutrition in nursing homes in The Netherlands, Germany and Austria are influenced by different resident characteristics. Since other country-related factors could also play an important role in influencing differences in the prevalence rates of malnutrition between the countries (structural and process factors of malnutrition care policy). We recommend the investigation of these factors in future studies.

Key words: Malnutrition: Resident characteristics: Nursing homes

Malnutrition is an important and still rather under-recognised problem in health care^(1–8). Malnutrition refers to negative deviations from a normal nutritional status, and it has been defined as inadequate nutritional status, undernourishment due to poor dietary intake, poor appetite, muscle wasting and weight loss⁽⁹⁾. Elia⁽¹⁰⁾ defined malnutrition as a nutritional condition in which an insufficient or disproportionate intake of energy, protein and other nutrients adversely affects tissue/body form (shape, size and composition) and function, and clinical outcomes.

Malnutrition increases the chance of complications. It worsens the immune function, leading to a higher risk of infections, and impairs wound healing. Moreover, malnutrition

impairs the quality of life and increases the length of hospital stay and costs of health care^(11–17).

Prevalence rates of malnutrition vary enormously internationally^(7,18–22), since in European nursing homes, the prevalence rates of malnutrition have been found to range between 2 and 74%^(23–25). These variations can partly be explained by differences in methodology and instruments used to measure malnutrition, and also resident characteristics can have an influence⁽²⁶⁾. For instance, age, sex, morbidity and care dependency are related to malnutrition^(4,9,19,25–29), as well as to infections⁽³⁰⁾, physical disabilities⁽³¹⁾ and polypharmacy^(32,33).

The aim of the present study was to investigate whether resident characteristics influence possible differences in

Abbreviation: LPZ, Landelijke Prevalentiemeting Zorgproblemen (National Prevalence Measurement of Care Problems).

* **Corresponding author:** N. C. van Nie-Visser, fax +31 43 388 41 62, email n.vannie@maastrichtuniversity.nl

malnutrition prevalence between countries, when using the same measurement methodology and instruments. The present hypothesis was that various resident characteristics influence differences in the prevalence rates of malnutrition between The Netherlands, Germany and Austria.

The following research questions will be investigated: (1) What is the prevalence rate of malnutrition in nursing homes in The Netherlands, Germany and Austria? (2) Are the characteristics of malnourished residents different in the three countries? (3) Which resident characteristics influence malnutrition? (4) Is the prevalence of malnutrition in nursing homes in The Netherlands, Germany and Austria different when controlling for the resident characteristics that influence the difference in malnutrition prevalence?

Methods

For the present study, data were used from the National Prevalence Measurement of Care Problems (in Dutch, Landelijke Prevalentiemeting Zorgproblemen (LPZ)), which is executed annually. Since 2004, the LPZ has been measuring the prevalence, prevention and treatment of malnutrition and quality indicators of nutritional care. It involves an annually conducted measurement in different health care settings (hospitals, long-term care and home care)⁽³⁴⁾. In 2008, the LPZ measurement expanded internationally to Germany and Austria (LPZ-International). In each country, data are gathered with the same instruments according to the same procedure. These countries conduct the same standardised measurement, supported by the project group of the LPZ⁽³⁵⁾. In each country, the coordination of the LPZ is carried out by a national project group led by a national coordinator. The Dutch LPZ project group facilitates each participating country with all documents and a website in their own language to enable them to promote, support and carry out the measurement. Each year, the national coordinators have an international research group meeting to discuss relevant issues and updates concerning possible changes in questionnaires, measurement procedures and cooperations^(20,36,37).

Design

The LPZ uses a cross-sectional, multi-centre design. For the present study, the data of LPZ-International collected in April 2009 and April 2010 in Dutch, German and Austrian nursing homes were analysed.

Instrument

Data were gathered using a standardised questionnaire at the patient level. As demographic data, age, sex, date of admission, co-morbidity, care dependency, weight, height and unintentional weight loss were measured. Malnutrition was operationalised and validated according to Meijers⁽³⁸⁾ and Meijers *et al.*^(7,39). Residents were qualified as malnourished if they met one of the following criteria: (1) BMI ≤ 20 kg/m² (age >65 years); (2) unintentional weight loss (more than 6 kg in the previous 6 months or more than 3 kg in the last

month); (3) no nutritional intake for 3 d or reduced intake for more than 10 d combined with a BMI ranging between 20 and 23.9 kg/m² (age >65 years).

Care dependency was measured with the Care Dependency Scale^(40,41). This scale consists of fifteen items, with a five-point Likert scale, and is validated for different settings in several countries^(42–45).

Since the original questionnaire and instruction material were in Dutch, these were translated by a professional translator into German. This translation was discussed by the Dutch project group (who spoke also German) with the project group in Germany and Austria until consensus was reached about the translation. The questionnaire was adapted to cultural differences. For instance, the nomenclatures for departments and professions that are present in German and Austrian nursing homes were adjusted to the local situation.

Sample

All Dutch, German and Austrian nursing homes were invited by (e)mail (including a flyer) and through publications in several professional journals to take part voluntarily in the LPZ measurement.

All residents of the participating nursing homes were invited to participate and included if they (or their legal representatives) gave informed consent. To get a more homogeneous sample, residents were included if they were at least 65 years old. Only those residents who were present at the day of the measurement and who were able to participate in the study were included. Residents were excluded when refusing to participate, not being available at the ward, being comatose or too ill and being terminal. In addition, data from 2009 of residents who participated both in 2009 and 2010 were excluded.

The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by different ethical committees in the different countries. In addition, the LPZ team received ethical approval from the medical ethical committee of the Maastricht University Medical Centre in The Netherlands (oral informed consent). The ethical committee related to the Institute of Nursing Science at Witten/Herdecke University gave its approval for the measurement in Germany and the medical ethical committee of the Medical University Graz approved to carry out the study in Austria (Germany and Austria: written informed consent).

Data collection

The participating nursing homes chose a coordinator who was responsible for the measurement within the institution. The coordinators were trained collectively by each national LPZ project group on how to organise data collection, and how to use the questionnaire and the specially designed Internet data-entry program. Subsequently, all the coordinators trained the health care professionals who would perform the data collection. For this purpose, the coordinators received a protocol and training package from the Dutch project



group to support them in training the health care professionals.

The assessment of the residents took place always by pairs of health care professionals (nurses, dietitians or doctors), one working on the resident's ward and one independent observer from another ward to enhance reliability.

Data analyses

Statistical analyses were performed using SPSS version 19 (SPSS, Inc.). χ^2 tests, Student's *t* test or ANOVA (with *post hoc* analyses using the Bonferroni method) and OR were used to describe the differences in (malnourished) resident characteristics between The Netherlands, Germany and Austria. Resident characteristics involved variables such as age, sex, length of stay, type and number of diseases and care dependency. Univariate logistic regression analyses were performed to describe the relationship of each baseline independent variable with the prevalence of malnutrition. Independent variables were country (0 = The Netherlands, 1 = Germany and 2 = Austria) and resident characteristics such as age, sex, length of stay, type and number of diseases and care dependency. For identifying differences in malnutrition prevalence between the countries, *P* values were based on two-sided tests, and the cut-off point for statistical significance was set at $P < 0.05$.

A univariate logistic generalised estimating equation (GEE) regression analysis was performed to estimate the OR of countries regarding the prevalence of malnutrition. The dependent variable was malnourished/not malnourished; the independent variables were two dummy variables indicating country (with The Netherlands as the reference category). GEE analysis corrects for the dependency of observations of individuals within institutions by adding a 'within-subject correlation structure' to the regression model. An exchangeable correlation structure was used, which means that correlations between individuals within the institutions are assumed to be the same. For building the association model, all variables that were significantly different between the three countries and related to malnutrition (both with $P < 0.10$) were seen as possible influencing variables (or confounders) in the GEE analyses. For that purpose, in the multivariate logistic regression analysis, all factors that were related to country and malnutrition difference (with $P < 0.10$) were added to the univariate model step by step so that the mean of both regression coefficients of the dummy variables for country changed. Only the covariates that led to a significant change (more than 10% of the regression coefficients) were included⁽⁴⁶⁾.

In the final multivariate model (corrected model), the OR of malnutrition in nursing homes in The Netherlands, Germany and Austria were estimated when controlling for the influencing resident characteristics. In this analysis, we focused on the change in the OR of malnutrition between the countries in the uncorrected model (univariate, without controlling for influencing resident characteristics) compared with the corrected model (multivariate, controlling for the observed influencing resident characteristics, see Table 5). Before the

multivariate analysis, data were assessed for congruence with regression assumptions.

Results

Response

In the present study, 214 nursing homes with 19876 residents were included in the analyses: 133 nursing homes from The Netherlands (*n* 14 123), sixty-one nursing homes from Germany (*n* 3973) and twenty nursing homes from Austria (*n* 1780). The response rate was significantly higher in The Netherlands (92.9%) than in Germany (82.9%) and Austria (80.8%). The reasons for not taking part in the measurement were refusing to participate (64.3%), not being available at the ward (27%), being comatose or too ill (5.7%) and being terminal (3.0%).

Resident characteristics

In Table 1, the characteristics of the included residents are shown separately for The Netherlands, Germany and Austria. Dutch residents were more often male, had a shorter mean length of stay, were less dependent of care and had fewer diseases than residents in Germany and Austria.

The most prevalent diseases in all the three countries were dementia (42.1% in The Netherlands, 55.2% in Germany and 60.8% in Austria), CVD (41.2% in The Netherlands, 70.0% in Germany and 59.0% in Austria) and motor disorders (27.0% in The Netherlands, 41.9% in Germany and 41.7% in Austria).

Malnutrition prevalence

The prevalence of malnutrition also differed significantly between the countries ($P < 0.05$; Table 2).

In Germany and Austria, the prevalence was somewhat higher than in The Netherlands (20.0 and 22.7% *v.* 18.0%, respectively).

Relationship between resident characteristics and malnutrition

Table 3 shows the prevalence of the different characteristics for the malnourished and not malnourished residents. Malnourished residents had more diseases and were older, more care dependent and more often female than those not malnourished. Furthermore, a significant difference was found between the malnourished and not malnourished residents concerning the type of prevalent diseases such as infectious disease, cancer, diabetes mellitus, blood disease, dementia, diseases of the digestive tract, injury resulting from accidents and total hip replacement. No significant difference was found in the length of stay.

To calculate which resident characteristics influenced the differences found in the prevalence rates of malnutrition between the countries, the factors that showed a significant difference between the countries and between the malnourished

Table 1. Resident characteristics, number and kind of disease, care dependency and malnutrition prevalence (Mean values and standard deviations; percentages)

	The Netherlands		Germany		Austria		<i>P</i>
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Nursing homes	133	62.2	61	28.5	20	9.3	
Residents	14 123	71.0	3973	20.0	1780	9.0	
Sex							<0.0001*†‡
Male	3717	26.3	868	21.8	255	14.3	
Female	10 409	73.7	3105	78.2	1525	85.7	
Age (years)							<0.0001*†‡
Mean	84		83		85		
SD	7		8		8		
Age categories							<0.0001
65–74 years	1311	9.3	634	16.0	187	10.5	
75–84 years	5432	38.5	1396	35.1	557	31.3	
≥ 85 years	7383	52.3	1943	48.9	1036	58.2	
Length of stay (d)							<0.0001*†‡
Median	631		767		694		
Mean	1017		1407		1195		
SD	1463		2129		1953		
Care dependency							<0.0001*†‡
Dependent	8341	59.1	2927	73.7	1431	80.4	
Independent	5783	40.9	1045	26.3	349	19.6	
Diseases							
Infectious disease	173	1.2	72	1.8	58	3.3	<0.0001*†‡
Cancer	996	7.1	325	8.2	148	8.3	0.016*
Endocrine/nutritional/metabolic disease	645	4.6	497	12.5	230	12.9	<0.0001*†
Diabetes mellitus	2554	18.1	1103	27.8	329	18.5	<0.0001*‡
Blood disease	416	2.9	176	4.4	87	4.9	<0.0001*†
Psychological disorder	1734	12.3	1269	31.9	372	20.9	<0.0001*†‡
Dementia	5943	42.1	2192	55.2	1082	60.8	<0.0001*†‡
Nervous system disorder	1262	8.9	617	15.5	264	14.8	<0.0001*†
Eye/ear disorder	2728	19.3	552	13.9	330	18.5	<0.0001*‡
Spinal cord lesion/paraplegia	29	0.2	10	0.3	8	0.4	0.132
CVD	5819	41.2	2782	70.0	1051	59.0	<0.0001*†‡
CVA/hemiparesis	2677	19.0	714	18.0	319	17.9	0.265
Respiratory disorder/disease	1966	13.9	402	10.1	173	9.7	<0.0001*†
Diseases of the digestive tract	1610	11.4	1101	27.7	367	20.6	<0.0001*†‡
Disorder/disease of kidney/urinary tract	1658	11.7	1029	25.9	379	21.3	<0.0001*†‡
Skin disorder	903	6.4	249	6.3	133	7.5	0.187
Motor disorder	3814	27.0	1663	41.9	742	41.7	<0.0001*†
Congenital disorders	162	1.1	101	2.5	49	2.8	<0.0001*†
Injury resulting from accidents	493	3.5	202	5.1	153	8.6	<0.0001*†‡
Total hip replacement	769	5.4	222	5.6	128	7.2	0.011†‡
Other non-specified diseases	1116	7.9	452	11.4	208	11.6	<0.0001*†
Prevalent diseases							<0.0001*†‡
Mean	2.6		3.87		3.64		
SD	1.57		1.68		1.98		

CVA, cerebrovascular accident.

* Values were significantly different between The Netherlands and Germany.

† Values were significantly different between The Netherlands and Austria.

‡ Values were significantly different between Germany and Austria.

and not malnourished residents ($P < 0.1$) were incorporated in a multivariate GEE analysis.

Possible influencing variables (GEE analyses and association model)

Tables 1 and 3 show that sex, age, age categories, mean number of diseases and care dependency, as well as infectious disease, cancer, diabetes mellitus, blood disease, dementia, CVD, respiratory disease, diseases of the digestive tract, motor disorder, injury resulting from accidents, total hip replacement or other non-specified diseases were possible

confounders or influencing variables ($P < 0.01$). After entering these variables step by step in the univariate model, in the final multivariate model (corrected model; Table 4) the variables care dependency, CVD, diseases of the digestive tract, age, dementia, diabetes mellitus, sex, mean number of diseases, respiratory disease and other non-specified diseases were included as influencing variables for the difference in malnutrition prevalence in the three countries. The two most influencing resident characteristics (confounders) were care dependency and CVD. The confounders resulting from the analyses influence the OR of countries.

Table 2. Prevalence of malnutrition

(Number of prevalence of malnutrition and percentages; odds ratios and 95% confidence intervals)

	Prevalence of malnutrition		<i>P</i>	OR	95% CI
	<i>n</i>	%			
Total (<i>n</i> 19 771)	3729	18.9			
The Netherlands (<i>n</i> 14 021)*	2530	18.0			
Germany (<i>n</i> 3972)	795	20.0	0.05	1.137	1.040, 1.242
Austria (<i>n</i> 1778)	404	22.7	0.0001	1.335	1.186, 1.504

* Reference group.

After controlling for these variables in the final multivariate model (corrected model), the OR of malnutrition difference between the three countries declined. The OR of malnutrition prevalence declined when comparing the univariate model (without controlling for predictive resident characteristics) with the multivariate model (controlling for influencing resident characteristics) (see Table 4) between the countries (see Table 5). The OR of malnutrition between The Netherlands and Germany declined from 1.137 to 1.065

($P=0.201$). The OR of malnutrition between The Netherlands and Austria declined from 1.335 to 1.085 ($P=0.213$). After controlling for the influencing resident characteristics (confounders), there were no differences any more; the OR declined to 1, implying that the differences in the prevalence rates of malnutrition declined.

The difference was not significant any more between The Netherlands and Germany and Austria when controlling for these influencing resident characteristics.

Discussion

The present unique large-scale study explored whether resident characteristics influence possible differences in malnutrition prevalence between The Netherlands, Germany and Austria, when using the same measurement methodology and instrument.

The prevalence of malnutrition differed significantly between the countries. The highest prevalence was found in Austria (22.7%, The Netherlands 18.0% and Germany 20.0%). These prevalence findings are within the range of earlier internationally reported malnutrition prevalence rates. Data collected by the Nutrition Day survey showed a malnutrition prevalence of 16.7% in nursing home residents

Table 3. Patients with malnutrition (M+) /without malnutrition (M-) and patient characteristics (Odds ratios and 95% confidence intervals)

	M+ (<i>n</i> 3729)	M- (<i>n</i> 16 042)	<i>P</i>	OR	95% CI
Sex (%)			<0.0001	1.420	1.306, 1.543
Male	19.4	25.5			
Female	80.6	74.5			
Mean age (years)	85.1	84.0	<0.0001	1.026	1.021, 1.031
Age categories (%)			<0.0001	1.221	1.156, 1.289
65–74 years	8.6	11.2			
75–84 years	34.1	37.8			
≥ 85 years	57.2	51.0			
Mean length of stay (days)	1143	1184	0.101	1	1.000, 1.000
Care dependency (%)			<0.0001	2.190	2.016, 2.379
Dependent	22.8	11.9			
Independent	77.2	88.1			
Diseases (%)					
Infectious disease	1.9	1.4	0.026	1.354	1.036, 1.768
Cancer	8.5	7.1	0.003	1.214	1.066, 1.382
Endocrine/nutritional/metabolic disease	7.5	6.8	0.136	1.109	0.968, 1.272
Diabetes mellitus	14.7	21.4	<0.0001	0.633	0.574, 0.699
Blood disease	4.0	3.3	0.027	1.231	1.024, 1.482
Psychological disorder	16.3	17.1	0.253	0.946	0.859, 1.041
Dementia	56.3	44.1	<0.0001	1.632	1.519, 1.754
Nervous system disorder	11.2	10.7	0.402	1.050	0.935, 1.176
Eye/ear disorder	17.8	18.3	0.491	0.968	0.882, 1.062
CVD	44.9	49.6	<0.0001	0.830	0.773, 0.892
CVA/hemiparesis	17.9	18.8	0.176	0.938	0.855, 1.029
Respiratory disorder/disease	14.1	12.5	0.011	1.145	1.032, 1.269
Diseases of the digestive tract	18.5	14.8	<0.0001	1.303	1.187, 1.431
Disease of kidney/urinary tract	15.6	15.4	0.747	1.016	0.921, 1.121
Skin disorder	6.3	6.5	0.574	0.959	0.828, 1.110
Motor disorder	33.0	30.9	0.015	1.099	1.019, 1.186
Congenital disorders	1.4	1.6	0.323	0.858	0.634, 1.162
Injury resulting from accidents	5.7	4.0	<0.0001	1.458	1.242, 1.711
Total hip replacement	6.5	5.4	0.012	1.207	1.042, 1.399
Other non-specified diseases	10.4	8.6	0.001	1.207	1.042, 1.399
Mean number of prevalent diseases	3.02	2.90	<0.0001	1.031	1.011, 1.051

CVA, cerebrovascular accident.

Table 4. Generalised estimating equation (GEE) – association model

	B1*	B2†	95 % CI (B1)	95 % CI (B2)	P (B1)	P (B2)
Uncorrected model‡	-0.289	-0.128	-0.408, -0.170	-0.217, -0.039	0.00	0.05
Corrected model§	-0.079	-0.063	-0.204, 0.046	-0.161, 0.034	0.213	0.201

* B1, comparing Austria with The Netherlands.

† B2, comparing Germany with The Netherlands.

‡ Variables in model: country (The Netherlands as a reference group, Germany and Austria) and malnutrition prevalence.

§ Corrected for variables: care dependency, age, sex, mean number of diseases and specific diseases (CVD, disease of the digestive tract, dementia, diabetes mellitus, respiratory disorder and other non-specified diseases).

in Germany⁽⁶⁾. Furthermore, a study by Tannen *et al.*⁽⁴⁷⁾ has shown comparable rates (15.1%) collected in German nursing homes, and another study has shown the same rates (15.7%) in Austrian hospitals⁽⁴⁸⁾. A study in different aged care residents in Australia has shown a much higher prevalence rate. The prevalence of malnutrition across these facilities varied from 31.8 to 72.1%⁽¹⁹⁾. In interpreting the differences between these studies, it must be realised that each study uses its own methodology as well as definition and operationalisation of malnutrition.

Resident characteristics that were both related to country and malnutrition were seen as possible influencing variables (or confounders) in the GEE analyses. Being female, being older, having more diseases and having diseases such as CVD, disease of the digestive tract, dementia, diabetes mellitus, respiratory disorder and other non-specified diseases influence the chance of getting malnourished.

The present study revealed that the two largest confounders are being more care dependent and having a CVD. This finding is confirmed in other studies showing that malnourished residents are more care dependent. Other studies confirm the present findings on the relationship between malnutrition and CVD^(49,50). Furthermore, the remaining confounders found in the GEE analyses have also been indicated by other studies. Residents have more often several diseases such as disease of the digestive tract and respiratory disorder^(31,51). Other studies have shown that women are more at risk of getting malnourished⁽⁵²⁾. Gaskill *et al.*⁽¹⁹⁾ found a relationship between malnutrition and an increased age and

high level of care needs. Suominen *et al.*⁽⁵¹⁾ described similar patient-related factors that explain malnutrition in nursing home residents in Finland. Their logistic regression analyses have shown that impaired functioning, swallowing difficulties, dementia and constipation are associated with being malnourished⁽⁵¹⁾. This is also in line with the present findings. Although the present study shows that resident characteristics influence differences in the prevalence rates of malnutrition between countries, it has never been studied using the same method and definition on a large scale in different countries.

There was no significant difference between The Netherlands, Germany and Austria after controlling for these influencing variables. In addition to resident characteristics, other influencing variables could also play a role in the difference in malnutrition prevalence between countries. Possible influencing variables could be differences in health care structure as nutritional care policies (e.g. nutritional screening policy, implementation of a nutritional care protocol/guideline, the policy of discussing malnourished residents in a multidisciplinary team) and care processes (e.g. preventive and treatment measures used). A study by Meesterberends *et al.*⁽⁵³⁾ has revealed that six factors, including resident-related, nursing-related and structure-related factors, explain the differences in the incidence rates of pressure ulcer between nursing homes in The Netherlands and Germany. Future studies must be performed to assess the specific contribution of these structural and process factors to differences in the prevalence rates of malnutrition in different countries.

Table 5. OR of malnutrition in The Netherlands, Germany and Austria, controlling for influencing resident characteristics (Odds ratios and 95 % confidence intervals)

	P	OR	95 % CI
Univariate			
The Netherlands*			
Germany	0.05	1.137	1.040-1.242
Austria	0.001	1.335	1.186-1.504
Multivariate†			
The Netherlands			
Germany	0.201	1.065	0.967-1.174
Austria	0.213	1.083	0.955-1.227

* Reference group.

† Model includes variables: care dependency, age, sex, mean number of diseases and specific diseases (CVD, disease of the digestive tract, dementia, diabetes mellitus, respiratory disorder and other non-specified diseases).

Limitations

In the present study, data from three countries were analysed with different sample sizes. While the institutions participated voluntarily, no information was available about the degree of representativeness of the samples. However, to date, no other studies with such a large number of patients have been published.

Since there is no globally accepted 'gold' standard for malnutrition, we based the present study on a definition that meets those factors for which consensus exists⁽³⁹⁾.

Finally, it might be possible that more and other resident characteristics that were not taken into account in the present study are likely to influence the risk of getting malnourished. Therefore, we assume that structural and process factors of malnutrition could also play a role.

Conclusion

Malnutrition is still a considerable problem; about 20% of all nursing home residents in the present study were malnourished. There are differences between countries, which can be explained by resident characteristics. Since other country-related factors such as structural and process factors of malnutrition could also play an important role in influencing differences in the prevalence rates of malnutrition between countries, we recommend the investigation of these factors in future studies.

Acknowledgements

We thank Nutricia Advanced Medical Nutrition for providing an unrestricted grant. Nutricia Advanced Medical Nutrition had no role in the design, analysis or writing of this article.

N. v. N., J. M., J. S. and R. H. contributed to the design of the study. N. v. N., J. M., J. S., C. L., S. B. and R. H. helped in the data collection. N. v. N., J. M., M. S. and R. H. conducted the data analyses. N. v. N., J. M., J. S., C. L., M. S. and R. H. wrote the manuscript. M. S. provided significant advice and consultation concerning statistical issues. N. v. N., J. M., J. S., C. L., M. S. and R. H. approved the final manuscript.

There are no conflicts of interest to report.

References

1. Waitzberg DL, Caiaffa WT & Correia MI (2001) Hospital malnutrition: the Brazilian National Survey (IBRANUTRI): a study of 4000 patients. *Nutrition* **17**, 573–580.
2. Correia IM & Campos ACL (2003) The prevalence of hospital malnutrition in Latin America: the multicenter ELAN study. *Nutrition* **19**, 823–825.
3. Kruijenga HM, Wierdsma NJ, Van Bokhorst-de van der Schueren MAE, *et al.* (2003) Screening of nutritional status in The Netherlands. *Clin Nutr* **22**, 147–152.
4. Stratton RJ, Green CJ & Elia M (2003) *Disease-Related Malnutrition: An Evidence-Based Approach To Treatment*, 1st ed. Cambridge, MA: CABI Publishing.
5. Pirlich M, Schutz T, Norman K, *et al.* (2006) The German hospital malnutrition study. *Clin Nutr* **25**, 563–572.
6. Valentini L, Schindler K, Schlaffer R, *et al.* (2009) The first Nutrition Day in nursing homes: participation may improve malnutrition awareness. *Clin Nutr* **28**, 109–116.
7. Meijers JMM, Schols JMGA, Van Bokhorst-de van der Schueren MAE, *et al.* (2009b) Malnutrition prevalence in the Netherlands: results of the annual Dutch National Prevalence Measurement of Care. *Br J Nutr* **101**, 417–423.
8. Vanderwee K, Clays E, Bocquaert I, *et al.* (2010) Malnutrition and associated factors in elderly hospital patients: a Belgian cross-sectional, multi-centre study. *Clin Nutr* **29**, 469–476.
9. Chen CC, Schilling LS & Lyder CH (2001) A concept analysis of malnutrition in the elderly. *J Adv Nurs* **36**, 131–142.
10. Elia M (2000) *Guidelines for Detection and Management of Malnutrition*. Maidenhead: Malnutrition Advisory Group (MAG), Standing Committee of BAPEN.
11. Green CJ (1999) Existence, causes and consequences of disease-related malnutrition in the hospital and the community, and clinical and financial benefits of nutritional intervention. *Clin Nutr* **18**, 3–28.
12. Elia M, Stratton R, Russel C, *et al.* (2005) *The Costs of Disease-Related Malnutrition in the UK and Economic Considerations for the Use of Oral Nutritional Supplements (ONS) in Adults*. Redditch: BAPEN.
13. Russel C (2007) The impact of malnutrition on healthcare costs and economic considerations for the use of oral nutritional supplements. *Clin Nutr Suppl* **2**, 25–32.
14. Arvanitakis M, Beck A, Coppens P, *et al.* (2008) Nutrition in care homes and homecare: how to implement adequate strategies (report of the Brussel Forum (22–23 November 2007)). *Clin Nutr Suppl* **2**, 25–32.
15. Norman K, Pichard C, Lochs H, *et al.* (2008) Prognostic impact of disease-related malnutrition. *Clin Nutr* **27**, 481–488.
16. Banks M, Bauer J, Graves N, *et al.* (2010) Malnutrition and pressure ulcer risk in adults in Australian health care facilities. *Nutrition* **26**, 896–901.
17. Meijers JMM, Halfens RJG, Wilson L, *et al.* (2012) Estimating the cost associated with malnutrition in Dutch nursing homes. *Clin Nutr* **31**, 65–68.
18. Donini LM, De Felice MR & Cannella C (2007) Nutritional status determinants and cognition in the elderly. *Arch Gerontol Geriatr* **44**, Suppl. 1, 143–153.
19. Gaskill D, Black LJ, Isenring EA, *et al.* (2008) Malnutrition prevalence and nutrition issues in residential aged care facilities. *Australas J Ageing* **27**, 189–194.
20. Bartholomeyczik S, Reuther S, Luft L, *et al.* (2010) Prävalenz von Mangelernährung, Maßnahmen und Qualitätsindikatoren in deutschen Altenpflegeheimen – erste Ergebnisse einer landesweiten Pilotstudie (Prevalence of malnutrition, intervention and quality indicators in German nursing homes – first results of a nationwide pilot study). *Gesundheitswesen* **72**, 1–7.
21. Kaiser MJ, Bauer JM, Rämisch C, *et al.* (2010) Frequency of malnutrition in older adults: a multinational perspective using the mini nutritional assessment. *J Am Geriatr Soc* **58**, 1734–1738.
22. Westergren A, Lindholm C, Axelsson C, *et al.* (2008) Prevalence of eating difficulties and malnutrition among persons within hospital care and special accommodations. *J Nutr Health Aging* **12**, 39–43.
23. Volkert D, Kreuel K, Heseker H, *et al.* (2004) Energy and nutrient intake of young-old, old-old and very-old elderly in Germany. *Clin Nutr* **58**, 1190–1200.
24. Pauly L, Stehle P & Volkert D (2007) Nutritional situation of elderly nursing home residents. *Z Gerontol Geriatr* **40**, 3–12.
25. Meijers JMM, Halfens RJG, Van Bokhorst-de van der Schueren MAE, *et al.* (2009a) Malnutrition in Dutch healthcare: prevalence, prevention, treatment and quality indicators. *Nutrition* **25**, 512–519.
26. Westergren A, Wann-Hansson C, Bergh Börgdal E, *et al.* (2009) Malnutrition prevalence and precision in nutritional care differed in relation to hospital volume – cross-sectional survey. *Nutrition* **8**, 20.
27. Lewis EJ & Stacey JB (1990) Nutritional assessment in the elderly. In *Geriatric Nutrition*, pp. 73–87 [JE Morley, Z Glick and LZ Rubenstein, editors]. New York: Raven Press.
28. Perissinotto E, Pisent C, Sergi G, *et al.* (2002) Anthropometric measurements in the elderly, age and gender differences. *Br J Nutr* **87**, 177–186.
29. Aliabadi M, Kimiagar M, Ghayour-Mobarhan M, *et al.* (2008) Prevalence of malnutrition in free living elderly people in Iran: a cross-sectional study. *Asia Pac J Clin Nutr* **17**, 285–289.
30. Vitale E (2011) An evaluation of the association of malnutrition with nosocomial infections in elderly patients. *Immunopharmacol Immunotoxicol* (epublication ahead of print version).

31. Oliveira MRM, Fogaça KCP & Leandro-Merhi VA (2009) Nutritional status and functional capacity of hospitalized elderly. *Nutr J* **8**, 54.
32. Heuberger RA & Caudell K (2011) Polypharmacy and nutritional status in older adults. A cross-sectional study. *Drug Aging* **28**, 315–323.
33. Jyrkkä J, Mursu J, Enlund H, *et al.* (2012) Polypharmacy and nutritional status in elderly people. *Curr Opin Clin Nutr Metab Care* **15**, 1–6.
34. Halfens RJG, Meesterberends E & Meijers JMM, *et al.* (2011) Rapportage resultaten: Landelijke Prevalentiemeting Zorgproblemen 2011 Annual Report. Maastricht University, Maastricht, The Netherlands.
35. Nie van NC, Schols MGA, Lohrmann C, *et al.* (2013) An international prevalence measurement of care problems: study design. *J Adv Nurs* **69**, e18–e29.
36. Schönherr S, Halfens R, Meijers J, *et al.* (2012) Structural and process indicators of nutritional care: a comparison between Austrian hospitals and nursing homes. *Nutrition* **28**, 868–873.
37. Nie van NC, Meijers JMM, Schols MGA, *et al.* (2011) Comparing quality of nutritional care in Dutch and German nursing homes. *J Clin Nurs* **20**, 2501–2508.
38. Meijers JMM (2009) Awareness of malnutrition in healthcare. The Dutch perspective PhD Thesis, Maastricht University, School for Public Health and Primary Care, Maastricht, The Netherlands.
39. Meijers JMM, Van Bokhorst-de van der Schueren MAE, Schols JMGA, *et al.* (2010) Defining malnutrition: mission or mission impossible. *Nutrition* **26**, 432–440.
40. Dijkstra A (1998) Care dependency – an assessment instrument for use in long term care facilities PhD Thesis, Groningen: Rijksuniversiteit Groningen.
41. Lohrmann C (2003) Pflegeabhängigkeitsskala: ein Einschätzungsinstrument für Heime und Kliniken – Eine methodologische Studie PhD Thesis, Berlin Center of Humanities and Health Science, Charité-Universitätsmedizin, Berlin, Germany.
42. Dijkstra A, Buist G, Moorer P, *et al.* (1999) Construct validity of the Nursing Care Dependency Scale. *J Clin Nurs* **8**, 380–388.
43. Dijkstra A, Brown L, Havens B, *et al.* (2000) An international psychometric testing of the Care Dependency Scale. *J Adv Nurs* **31**, 944–952.
44. Dijkstra A, Coleman M, Tomas C, *et al.* (2003) Cross-cultural psychometric testing of the Care Dependency Scale with data. *J Adv Nurs* **43**, 181–187.
45. Dijkstra A, Tiesinga LJ, Plantinga L, *et al.* (2005) Diagnostic accuracy of the Care Dependency Scale. *J Adv Nurs* **50**, 410–416.
46. Twisk JWR (2010) *Inleiding in de toegepaste biostatistiek* (Introduction to Applied Biostatistics). Amsterdam: Reed Business.
47. Tannen A, Schutz T, Nie van NC, *et al.* (2008) Mangelernährung in deutschen Pflegeheimen und Krankenhäusern – Pflegebedarf und pflegerische Versorgung (Malnutrition in German nursing homes and hospitals – care dependency and nursing interventions). *Aktuelle Ernährungsmedizin* **33**, 177–183.
48. Tannen A & Lohrmann C (2012) Malnutrition in Austrian hospital patients. Prevalence, risk factors, nursing interventions, and quality indicators: a descriptive multicentre study. *J Adv Nurs* **69**, 1840–1849.
49. Colín-Ramírez E, Orea-Tajeda A, Castillo-Martínez L, *et al.* (2011) Malnutrition syndrome, but not body mass index, is associated to worse prognosis in heart failure patients. *Clin Nutr* **30**, 753–758.
50. Teh R, Wham C, Kerse N, *et al.* (2010) How is the risk of undernutrition associated with cardiovascular disease among individuals of advanced age? *J Nutr Health Aging* **14**, 737–743.
51. Suominen M, Muurinen S, Routasalo P, *et al.* (2005) Malnutrition and associated factors among aged residents in all nursing homes in Helsinki. *Eur J Clin Nutr* **59**, 578–583.
52. Hickson M (2006) Malnutrition and ageing. *Postgrad Med J* **82**, 2–8.
53. Meesterberends E, Halfens R, Spreeuwenberg M, *et al.* (2013) Do patients in Dutch nursing homes have more pressure ulcers than patients in German nursing homes? A prospective multi-centre cohort study. *JAMDA* **14**, 605–610.