

Optimum feeding and growth in preterm neonates

J. E. Harding*, J. G. B. Derraik, M. J. Berry, A. L. Jaquiery, J. M. Alsweiler, B. E. Cormack and F. H. Bloomfield

Liggins Institute, University of Auckland, Auckland, New Zealand

Approximately 10% of all babies worldwide are born preterm, and preterm birth is the leading cause of perinatal mortality in developed countries. Although preterm birth is associated with adverse short- and long-term health outcomes, it is not yet clear whether this relationship is causal. Rather, there is evidence that reduced foetal growth, preterm birth and the long-term health effects of both of these may all arise from a suboptimal intrauterine environment. Further, most infants born preterm also experience suboptimal postnatal growth, with potential adverse effects on long-term health and development. A number of interventions are used widely in the neonatal period to optimise postnatal growth and development. These commonly include supplementation with macronutrients and/or micronutrients, all of which have potential short-term risks and benefits for the preterm infant, whereas the long-term health consequences are largely unknown. Importantly, more rapid postnatal growth trajectory (and the interventions required to achieve this) may result in improved neurological outcomes at the expense of increased cardiovascular risk in later life.

Received 13 July 2012; Revised 1 November 2012; Accepted 27 November 2012; First published online 8 January 2013

Key words: fetal growth restriction, infant diet, infant growth, infant, premature, neonatal nutrition

Introduction

It is estimated that 10% of all births are preterm, and this incidence is increasing worldwide.¹ Every year, over 1 million infants die because of complications associated with preterm birth, which is the leading cause of neonatal mortality worldwide and the second leading cause of death among children under 5 years of age.¹ Preterm babies are also frequently born small-for-gestational-age, and there is a large body of epidemiological evidence indicating that reduced size at birth is linked to increased risk of developing a number of diseases in later life. These include hypertension,² impaired glucose tolerance,³ obesity^{4,5} and coronary heart disease.^{6,7}

Earlier studies focused on reduced size at birth in babies born at term, which was assumed to result from impaired foetal growth. However, more recent studies suggest that preterm birth, independent of foetal growth rate and even at late preterm gestations, is itself associated with increased blood pressure,⁸ altered blood pressure regulation,⁹ diabetes and insulin resistance,^{10–14} as well as ischaemic heart disease and stroke¹⁵ in later life. The adverse short- and long-term neurological and social outcomes of preterm birth are well known.^{16,17} Nonetheless, it is becoming increasingly apparent that the adverse metabolic and neurodevelopmental outcomes following preterm birth not only affect those born at the extremes of viability but appear to be present as a graded response across preterm gestations, right up to and including late preterm births.^{8,18}

Both infants born preterm and those who experience intrauterine growth restriction (IUGR) are exposed to early environmental stress and suboptimal nutrition. This insult becomes apparent clinically during the later stages of pregnancy in IUGR infants, whereas for those born preterm this nutritional restriction occurs early in postnatal life, which is comparable to the third trimester of pregnancy.¹⁹ However, preterm infants are also much more likely to experience poor growth *in utero*, and those infants born extremely preterm (24–32 weeks) appear to grow poorly as early as the first trimester of pregnancy.²⁰ As a result, the rate of IUGR is much greater among infants born preterm than those born at term,²¹ and it has been estimated that 40% of preterm babies have also experienced IUGR.²² The origins of a significant proportion of IUGR, such as that secondary to placental vascular disease, also lie in early pregnancy.

The causes of preterm birth are multi-factorial;²³ however, irrespective of the underlying causes, most infants born preterm also experience suboptimal postnatal growth. Further, although this suboptimal growth is often most profound in the immediate postnatal period,²⁴ it may also persist through childhood and adolescence. After the neonatal period, this suboptimal growth may be in part a result of alterations in the endocrine regulation of growth, which appears to be disrupted in preterm children and adolescents, so that growth is no longer associated with its normal hormone regulators.¹⁹ However, a major contributor to postnatal faltering growth in infants born preterm is nutritional disruption due to inadequate nutrient intake.²⁵ It is common practice in neonatal units worldwide to increase intravenous nutrition incrementally in preterm infants over the 1st week of life and to introduce enteral feeds

*Address for correspondence: J. E. Harding, Liggins Institute, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand.
(Email j.harding@auckland.ac.nz)

cautiously. When compared with estimated placental nutrient transfer *in utero*,²⁵ there is an inevitable and substantial nutrient deficit, particularly for nitrogen, by the end of the 1st week.²⁶ This deficit occurs even if the latest recommended nutritional intakes are met, and contributes to early postnatal growth faltering. There are potentially serious consequences for the preterm infant, as postnatal malnutrition not only hinders somatic growth but may also adversely affect organ growth, structure and function.²²

Poor early postnatal growth is likely to be associated with adverse neurodevelopmental outcomes,^{24,27} but this relationship may be modifiable by other factors such as breastfeeding. In a recent large population study, infants born very preterm who were breastfed at the time of hospital discharge had significantly better neurodevelopmental outcomes than those who did not receive mothers' milk, despite an increased risk for poor early postnatal weight gain.²⁸ Thus, both early diet and early growth appear to be important mediators of later developmental outcomes.²⁸ However, it is important to appreciate that, in studies on breastfeeding, there are often confounding factors; in these studies, <20% of babies were breastfed at discharge, and maternal demographics were very different among breastfed and non-breastfed babies.

Cause and effect

Although preterm birth itself is associated with adverse short- and long-term health outcomes, it is not yet clear whether this relationship is causal. Rather, there is evidence that reduced foetal growth, preterm birth and the long-term health effects of both of these may all arise from a suboptimal intrauterine environment. In addition, there is some evidence that decreased foetal growth and shorter gestation length may also be linked to common antecedents, suggesting that further research into this area is warranted. In humans, for example, poor maternal nutrition before pregnancy or in the periconceptional period is associated with an increased risk for preterm birth.²³ Mothers who are underweight before pregnancy are also more likely to deliver preterm.^{29,30} Twins are born both early and small,³¹ outcomes that seem to have their origin in early pregnancy³² and which appear to be linked.³³ Further, poor maternal nutrition is also associated with poor long-term health in the offspring.³⁴

Animal studies have provided more conclusive evidence of the causal relationships involved. In sheep, maternal undernutrition around the time of conception led to preterm delivery in a proportion of animals.³⁵ Maternal undernutrition before conception also impaired the normal development of insulin resistance during pregnancy, which in turn was related to impaired foetal growth.³⁶ However, following periconceptional undernutrition, even the lambs born at term had evidence of disrupted regulation of early postnatal growth,³⁷ and in adulthood had altered body composition,¹⁰⁹ impaired glucose tolerance³⁸ and blunted adrenal response to corticotrophic stimulation.³⁹ These data

suggest that nutritional factors before and around the time of conception, rather than preterm birth itself, may be the primary cause of subsequent adverse outcomes.

Interventions

At an equivalent gestational age, a preterm infant has a very different nutritional intake from that supplied by the placenta to a foetus. During the second and the beginning of the third trimester of pregnancy, the estimated placental transfer to the foetus is 3.6–4.8 g/kg.d of protein and 12–14 g/kg.d of glucose.⁴⁰ *In utero*, protein uptake by the foetus exceeds the amount needed for protein accretion, and the excess is oxidised to produce energy. The total foetal requirement for fatty acids at mid-gestation is ~1 g/kg.d, suggesting that oxidation of fatty acids for energy is relatively unimportant in foetal life.⁴⁰ In comparison, guidelines for intravenous nutrition in preterm infants recommend high intakes for both fat (3.0–4.0 g/kg.d) and carbohydrate (up to 18 g/kg.d), with a much lower protein content (1.5–4.0 g/kg.d) than the estimated placental transfer.^{41,42} However, in clinical practice, these nutrient intakes are difficult to meet with current intravenous nutrition solutions and delivery systems. In reality, many preterm infants receive very low protein intakes in the first few days after birth,^{43,44} resulting in large deficits in energy and protein.^{45,46} Further, the amount and ratio of macronutrients given intravenously to preterm infants differ from the estimated *in utero* supply. These differences may promote the accretion of fat mass rather than lean body mass and may explain some of the observed differences in the body composition between preterm and term babies.^{47–49}

Higher protein and energy intakes in preterm infants during the first few days after birth reduce growth faltering at discharge.^{50–55} Other studies report favourable effects of earlier and/or higher total protein intake on the incidence of chronic lung disease,^{56–58} although most studies found no differences in clinical outcomes.^{50,59–61} Nonetheless, a nutritional approach that matches the estimated foetal macronutrient intakes has yet to be developed. Furthermore, protein intake is not the only determinant of growth, as an adequate balance of macronutrients and micronutrients is crucial for optimal growth and development in early postnatal life.⁶²

Breast milk is widely recognised to be the best enteral feed, but it is inadequate to sustain postnatal growth of preterm babies at intrauterine growth rates. Importantly, the concentration of protein in expressed breast milk appears to be even lower than previously estimated.^{63,64} Consequently, commercial breast milk fortifiers are commonly added to supplement the infant diet with additional quantities of protein and several other nutrients.^{41,65} However, even with breast milk fortification, recommended enteral protein intakes are seldom achieved.^{66–68} In addition, consensus recommendations for enteral protein intake were recently increased to 4.0–4.5 g/kg.d, resulting in the reformulation of several commercial breast milk fortifiers to increase the supplementary amount of added protein to

1.0–1.2 g per 100 ml breast milk and 2.5 g per 100 ml in some preterm formulae.⁶⁵

There is growing evidence that manipulation of the macronutrient composition of milk leads to changes in weight gain and proportionality in the infant.^{69,70} Systematic reviews report that both protein and multi-component breast milk fortification increase short-term growth parameters.^{71,72} However, the level of protein fortification in the studies included in these reviews was lower than that supplied by newly available fortifiers, and the long-term effects of higher protein fortifiers and formulas are yet to be determined.

There is evidence that postnatal faltering growth is associated with adverse neurodevelopmental outcomes,^{24,27} and improved growth over the first few months after birth has been shown to be associated with better outcomes at 18 months of age.⁷³ There is also evidence that nutrient-enriched formula provided to preterm infants after discharge may lead to improved later development.^{73–75} A randomised trial of nearly 300 small-for-gestational-age infants born at term also showed significant gains in length and head circumference on an enriched formula.⁷⁶ However, this trial reported adverse developmental outcomes at 9 months of age, although these seem to have disappeared at 18 months.⁷⁷

Further investigation is also required into the long-term effects of other dietary components in preterm neonates. For example, supplementation of breast milk with fat alone has little effect on postnatal growth.⁷⁸ However, additional long-chain polyunsaturated acids provided in breast milk via maternal supplementation are associated with improved problem-solving subscores (but not overall developmental scores) at the age of 6 months,⁷⁹ and in the Bayley mental development index at the age of 18 months in girls but not in boys.⁸⁰ A subgroup of children enrolled in this trial underwent an assessment of language skills, behaviour and temperament in early childhood (2–5 years of age); there were no significant differences between those who received docosahexaenoic acid supplementation and those who did not.⁸¹ Data on possible later cardiovascular effects of these interventions are not yet available.

Similarly, some neonatal units routinely prescribe calcium supplementation for very preterm babies to facilitate appropriate bone growth. There are few good data to support this;¹¹⁰ small trials have not found a benefit in bone mineral density by hospital discharge.¹¹¹ Supplementation of breast milk with a multi-nutrient fortifier for 12 weeks after discharge did result in higher calcium and phosphate intakes and increased skeletal growth but not bone density.⁸² However, supplementation reduced milk intake in these infants, so that protein and energy intakes and body composition at the end of the intervention period were comparable between supplemented and unsupplemented infants. Nonetheless, it is conceivable that neonatal calcium intake may improve later cardiovascular outcomes, as maternal calcium supplementation during pregnancy has been shown to be associated with lowered blood pressure in the offspring.⁸³

In lambs, ewe milk fortification (with a nutritional supplement analogous to breast milk fortifiers used clinically) for the first 2 weeks of life led to gestation- and sex-specific effects on early growth. For example, supplements altered growth in male but not female lambs, and resulted in increased weight-for-length (and by inference adiposity) at weaning in lambs born preterm but decreased adiposity in lambs born at term.⁸⁴ Some studies in humans have also shown differential effects in girls and boys following nutritional supplementation early in life.^{85,86}

Importantly, nutritional supplementation given in the neonatal period may also affect later endocrine function. In sheep, for example, early nutritional supplementation altered glucose–insulin axis function in females, independent of growth.⁸⁴ Further, although supplementation reduced the insulin-secretory response to both glucose and arginine in female lambs born preterm, the response was increased in females born at term. In humans, diet in the neonatal period has also been shown to influence later glucose–insulin axis function in babies. When term infants of diabetic women were fed on either maternal milk or banked non-diabetic breast milk, those with the greatest exposure to ‘diabetic’ milk in the 1st (but not 2nd) week after birth had the greatest risk of glucose intolerance at 2 years,⁸⁷ irrespective of the total duration of breastfeeding.⁸⁸ The enteroinsular axis is known to be active at birth⁸⁹ and to be modified by both maturity⁹⁰ and diet.⁹¹ It is therefore plausible that exposure to specific dietary components may modify pancreatic development in the newborn period. In addition, preterm infants have immature gastrointestinal tract function and experience delayed initiation of enteral feeding, as well as considerable variation in the type, composition and method of administration of milk feeds in the early neonatal period,⁹² all of which may have significant implications for later metabolic function.

Apart from nutritional interventions, common neonatal morbidities and their treatment can also affect nutritional intake and growth. For example, in a prospective birth cohort from the United States, delivery by Caesarean section at term was associated with a twofold increased risk of obesity at 3 years of age.⁹³ In preterm infants, corticosteroids are used to treat ventilator-dependent chronic lung disease but can significantly impair growth in preterm babies.⁹⁴ Hyperglycaemia may affect over 50% of extremely low-birth-weight infants⁹⁵ and is associated with increased mortality and morbidity.^{95,96} Hyperglycaemia in these babies is often managed by reducing glucose intake,⁹⁷ which may further impair growth. A recent clinical trial found that tight glycaemic control with insulin in hyperglycaemic preterm babies improved weight gain and mitigated the expected reduction in head circumference, suggesting a possible improvement in brain growth.⁹⁸ However, treatment reduced linear growth, suggesting an increase in fat mass. In addition, tight glycaemic control increased the risk for hypoglycaemia,⁹⁸ which is associated with poor neurodevelopmental outcomes in preterm babies.⁹⁹

Potential risks and benefits

The majority of infants born preterm undergo accelerated growth after discharge, so that 85% reach the normal height range by 2 years of age.¹⁰⁰ Nonetheless, nutritional interventions are commonly introduced to further accelerate their growth, even though the long-term benefits and costs of such interventions are still unclear. In fact, a number of studies have suggested that accelerated growth in early childhood may actually be detrimental to health in the long-term.

For example, in a cohort of very preterm babies, those with growth rates in the highest tertile over the first 2 years of life had lower insulin sensitivity and higher blood pressure as young adults, compared with those in the lowest tertile.¹⁴ Further, babies randomised to receive an enriched formula who exhibited accelerated neonatal growth had higher systolic blood pressure at 6–8 years of age than did babies fed a standard formula.¹⁰¹ Accelerated neonatal growth for as little as 2 weeks after birth was associated with higher levels of a marker of insulin resistance in adolescence,¹⁰² as well as with alterations in vasodilatation suggestive of impaired cardiovascular health.¹⁰³ The US Collaborative Perinatal Project studied nearly 56,000 pregnancies and showed that an increase in growth percentile during the first 7 years was associated with increased blood pressure.¹⁰⁴ A review of 21 studies identified that early accelerated growth (involving an increase of at least 0.67 weight SDS) was associated with a twofold to threefold increase in risk of overweight or obesity.^{105,106} In addition, accelerated growth was associated with a greater likelihood of developing insulin resistance and subsequently the metabolic syndrome.¹⁰⁶ Thus, these findings have led some authors to question the wisdom of encouraging rapid weight gain in infants born preterm or IUGR.¹⁰⁴

In contrast, other studies have observed beneficial effects of accelerated growth on neurodevelopmental outcomes. Preterm babies randomised to an enriched formula within 48 h of birth and fed this formula for an average of <6 weeks showed a one-third reduction in the risk for cerebral palsy in childhood.⁷⁷ Further, breastfed preterm babies supplemented with human milk fortifier for 6 weeks had accelerated growth with no adverse effects on neurodevelopment.¹⁰⁷ Observational studies have also reported that preterm babies who have higher growth rates in the neonatal period have higher neurodevelopmental scores in later life.^{24,108}

The relationship between accelerated postnatal growth and long-term cognitive and metabolic health is difficult to tease out from the underlying causes of accelerated growth itself. It is also important to take into account the end result of accelerated growth for an individual, such as the relationship between final and starting growth (expressed as a Z-score or centile). Accelerated growth usually follows a period of faltering growth, whether this is before birth in IUGR babies, after birth in babies with postnatal growth faltering as for most very preterm babies, or both before and after birth as may occur in preterm babies born IUGR. There are currently no proven interventions for improving intrauterine growth

in IUGR fetuses, suggesting that attention should be focussed on preventing the faltering postnatal growth that is so prevalent in babies born preterm.

Conclusions

Preterm birth is known to be associated with numerous short- and long-term adverse health outcomes. However, it is unclear whether these outcomes are a direct consequence of preterm birth itself or of associated factors, such as perturbed antenatal and/or postnatal growth and nutrition. There is evidence that both prenatal and postnatal nutritional restriction may contribute to adverse outcomes, and a number of interventions in the neonatal period have been shown to improve postnatal growth and development, at least in the short-term. Nonetheless, evidence for the long-term consequences of these interventions, particularly with respect to metabolic and cardiovascular health, is lacking, and most evidence on the long-term outcomes of neonatal nutrition and growth is epidemiological. High-quality prospective research in this area has lagged behind research in other aspects of neonatal care, and this deficit needs to be addressed. Future studies should attempt to dissect out the consequences of accelerated growth from its antecedents and whether the long-term risks and benefits of neonatal nutrition and growth, especially accelerated growth, may differ according to the outcome studied. This applies particularly to neurodevelopmental *v.* metabolic/cardiovascular outcomes, as targeting improvements in one aspect may involve a trade-off with respect to another.

Acknowledgment

The authors would like to thank all members of the LiFEPATH Group for their contributions to the referenced studies. This work was supported by the Health Research Council of New Zealand, the New Zealand Lotteries Grants Board, and the Auckland Medical Research Foundation.

References

1. March of Dimes, Partnership for Maternal Newborn & Child Health, Save the Children, World Health Organization. Born Too Soon: The Global Action Report on Preterm Birth, 2012. World Health Organization: Geneva. http://www.who.int/pmnch/media/news/2012/preterm_birth_report/en/index.html.
2. Law CM, Shiell AW, Newsome CA, *et al.* Fetal, infant, and childhood growth and adult blood pressure: a longitudinal study from birth to 22 years of age. *Circulation*. 2002; 105, 1088–1092.
3. Newsome CA, Shiell AW, Fall CHD, *et al.* Is birth weight related to later glucose and insulin metabolism? A systematic review. *Diabet Med*. 2003; 20, 339–348.
4. Kensara OA, Wootton SA, Phillips DI, *et al.* Fetal programming of body composition: relation between birth weight and body composition measured with dual-energy X-ray absorptiometry and anthropometric methods in older Englishmen. *Am J Clin Nutr*. 2005; 82, 980–987.

5. Ong KK. Size at birth, postnatal growth and risk of obesity. *Horm Res.* 2006; 65(Suppl. 3), 65–69.
6. Roseboom TJ, van der Meulen JH, Osmond C, *et al.* Coronary heart disease after prenatal exposure to the Dutch famine, 1944–45. *Heart.* 2000; 84, 595–598.
7. Eriksson JG, Forsen T, Tuomilehto J, Osmond C, Barker DJ. Early growth and coronary heart disease in later life: longitudinal study. *BMJ.* 2001; 322, 949–953.
8. Crump C, Winkleby MA, Sundquist K, Sundquist J. Risk of hypertension among young adults who were born preterm: a Swedish national study of 636,000 births. *Am J Epidemiol.* 2011; 173, 797–803.
9. Bayrakci US, Schaefer F, Duzova A, Yigit S, Bakaloglu A. Abnormal circadian blood pressure regulation in children born preterm. *J Pediatr.* 2007; 151, 399–403.
10. Lawlor DA, Davey Smith G, Clark H, Leon DA. The associations of birthweight, gestational age and childhood BMI with type 2 diabetes: findings from the Aberdeen Children of the 1950s cohort. *Diabetologia.* 2006; 49, 2614–2617.
11. Irving RJ, Belton NR, Elton RA, Walker BR. Adult cardiovascular risk factors in premature babies. *Lancet.* 2000; 355, 2135–2136.
12. Hovi P, Andersson S, Eriksson JG, *et al.* Glucose regulation in young adults with very low birth weight. *N Engl J Med.* 2007; 356, 2053–2063.
13. Hofman PL, Regan F, Jackson WE, *et al.* Premature birth and later insulin resistance. *N Engl J Med.* 2004; 351, 2179–2186.
14. Rotteveel J, van Weissenbruch MM, Twisk JWR, Delemarre-Van de Waal HA. Infant and childhood growth patterns, insulin sensitivity, and blood pressure in prematurely born young adults. *Pediatrics.* 2008; 122, 313–321.
15. Koupil I, Leon DA, Lithell HO. Length of gestation is associated with mortality from cerebrovascular disease. *J Epidemiol Community Health.* 2005; 59, 473–474.
16. Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet.* 2008; 371, 261–269.
17. Moster D, Lie RT, Markestad T. Long-term medical and social consequences of preterm birth. *N Engl J Med.* 2008; 359, 262–273.
18. MacKay DF, Smith GCS, Dobbie R, Pell JP. Gestational age at delivery and special educational need: retrospective cohort study of 407,503 schoolchildren. *PLoS Med.* 2010; 7, e1000289.
19. Rowe DL, Derraik JGB, Robinson E, Cutfield WS, Hofman PL. Preterm birth and the endocrine regulation of growth in childhood and adolescence. *Clin Endocrinol.* 2011; 75, 661–665.
20. Smith GCS, Smith MFS, McNay MB, Fleming JEE. First-trimester growth and the risk of low birth weight. *N Engl J Med.* 1998; 339, 1817–1822.
21. Cooke RW. Conventional birth weight standards obscure fetal growth restriction in preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 2007; 92, F189–F192.
22. Cooke RJ. Postnatal growth and development in the preterm and small for gestational age infant. In *Importance of Growth for Health and Development* Nestlé Nutrition Institute Workshop Series Pediatric Program, vol. 65. (eds. Lucas A, Makrides M, Ziegler EE), 2010; pp. 85–95. Karger AG: Basel.
23. Bloomfield FH. How is maternal nutrition related to preterm birth? *Annu Rev Nutr.* 2011; 31, 235–261.
24. Ehrenkranz RA, Dusick AM, Vohr BR, *et al.* Growth in the neonatal intensive care unit influences neurodevelopmental and growth outcomes of extremely low birth weight infants. *Pediatrics.* 2006; 117, 1253–1261.
25. Embleton NE, Pang N, Cooke RJ. Postnatal malnutrition and growth retardation: an inevitable consequence of current recommendations in preterm infants? *Pediatrics.* 2001; 107, 270–273.
26. Thureen PJ, Melara D, Fennessey PV, Hay WW. Effect of low versus high intravenous amino acid intake on very low birth weight infants in the early neonatal period. *Pediatr Res.* 2003; 53, 24–32.
27. Latal-Hajnal B, von Siebenthal K, Kovari H, Bucher HU, Largo RH. Postnatal growth in VLBW infants: significant association with neurodevelopmental outcome. *J Pediatr.* 2003; 143, 163–170.
28. Rozé J-C, Darmaun D, Boquien C-Y, *et al.* The apparent breastfeeding paradox in very preterm infants: relationship between breast feeding, early weight gain and neurodevelopment based on results from two cohorts, EPIPAGE and LIFT. *BMJ Open.* 2012; 2, e000834.
29. Han Z, Mulla S, Beyene J, Liao G, McDonald SD. Maternal underweight and the risk of preterm birth and low birth weight: a systematic review and meta-analyses. *Int J Epidemiol.* 2011; 40, 65–101.
30. Smith GCS, Shah I, Pell JP, Crossley JA, Dobbie R. Maternal obesity in early pregnancy and risk of spontaneous and elective preterm deliveries: a retrospective cohort study. *Am J Public Health.* 2007; 97, 157–162.
31. Muhlhauser BS, Hancock SN, Bloomfield FH, Harding R. Are twins growth restricted? *Pediatr Res.* 2011; 70, 117–122.
32. Hancock SN, Oliver MH, McLean C, Jaquiere AL, Bloomfield FH. Size at birth and adult fat mass in twin sheep are determined in early gestation. *J Physiol.* 2012; 590, 1273–1285.
33. Hediger ML, Luke B, Gonzalez-Quintero VH, *et al.* Fetal growth rates and the very preterm delivery of twins. *Am J Obstet Gynecol.* 2005; 193, 1498–1507.
34. Painter RC, Roseboom TJ, Bleker OP. Prenatal exposure to the Dutch famine and disease in later life: an overview. *Reprod Toxicol.* 2005; 20, 345–352.
35. Bloomfield FH, Oliver MH, Hawkins P, *et al.* A periconceptional nutritional origin for noninfectious preterm birth. *Science.* 2003; 300, 606.
36. Jaquiere AL, Oliver MH, Rumball CWH, Bloomfield FH, Harding JE. Undernutrition before mating in ewes impairs the development of insulin resistance during pregnancy. *Obstet Gynecol.* 2009; 114, 869–876.
37. Jaquiere AL, Oliver MH, Bloomfield FH, Harding JE. Periconceptional events perturb postnatal growth regulation in sheep. *Pediatr Res.* 2011; 70, 261–266.
38. Todd SE, Oliver MH, Jaquiere AL, Bloomfield FH, Harding JE. Periconceptional undernutrition of ewes impairs glucose tolerance in their adult offspring. *Pediatr Res.* 2009; 65, 409–413.

39. Oliver MH, Bloomfield FH, Jaquier AL, et al. Periconceptual undernutrition suppresses cortisol response to arginine vasopressin and corticotropin-releasing hormone challenge in adult sheep offspring. *J Dev Orig Health Dis.* 2012; 3, 52–58.
40. Hay WW, Lucas A, Heird WC, et al. Workshop summary: nutrition of the extremely low birth weight infant. *Pediatrics.* 1999; 104, 1360–1368.
41. Tsang RC, Uauy R, Koletzko B, Zlotkin SH. *Nutrition of the Preterm Infant: Scientific Basis and Practical Guidelines*, 2nd edn, 2005. Digital Education Publishing, Inc.: Cincinnati.
42. Koletzko B, Goulet O, Hunt J, et al. Guidelines on paediatric parenteral nutrition of the European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) and the European Society for Clinical Nutrition and Metabolism (ESPEN), supported by the European Society of Paediatric Research (ESPR). *J Pediatr Gastroenterol Nutr.* 2005; 41, S1–S87.
43. Lapillonne A, Fellous L, Mokthari M, Kermorvant-Duchemin E. Parenteral nutrition objectives for very low birth weight infants: results of a national survey. *J Pediatr Gastroenterol Nutr.* 2009; 48, 618–626.
44. Kiefer AS, Wickremasinghe AC, Johnson JN, et al. Medical management of extremely low-birth-weight infants in the first week of life: a survey of practices in the United States. *Amer J Perinatol.* 2009; 26, 407–418.
45. Ahmed M, Irwin S, Tuthill DP. Education and evidence are needed to improve neonatal parenteral nutrition practice. *J Parenteral Enteral Nutr.* 2004; 28, 176–179.
46. Stewart J, Mason D, Smith N, Protopapa K, Mason M. A Mixed Bag: an enquiry into the care of hospital patients receiving parenteral nutrition. London: National Confidential Enquiry into Patient Outcome and Death; 2010. Report No.: 0956088236.
47. Kashyap S, Forsyth M, Zucker C, et al. Effects of varying protein and energy intakes on growth and metabolic response in low birth weight infants. *J Pediatr.* 1986; 108, 955–963.
48. Kashyap S, Schulze KF, Forsyth M, et al. Growth, nutrient retention, and metabolic response in low birth weight infants fed varying intakes of protein and energy. *J Pediatr.* 1988; 113, 713–721.
49. Schulze KF, Stefanski M, Masterson J, et al. Energy expenditure, energy balance, and composition of weight gain in low birth weight infants fed diets of different protein and energy content. *J Pediatr.* 1987; 110, 753–759.
50. Wilson DC, Cairns P, Halliday HL, et al. Randomised controlled trial of an aggressive nutritional regimen in sick very low birthweight infants. *Arch Dis Child Fetal Neonatal Ed.* 1997; 77, F4–F11.
51. Poindexter BB, Langer JC, Dusick AM, Ehrenkranz RA. Early provision of parenteral amino acids in extremely low birth weight infants: relation to growth and neurodevelopmental outcome. *J Pediatr.* 2006; 148, 300–305.
52. Martin CR, Brown YF, Ehrenkranz RA, et al. Nutritional practices and growth velocity in the first month of life in extremely premature infants. *Pediatrics.* 2009; 124, 649–657.
53. Valentine CJ, Fernandez S, Rogers LK, et al. Early amino-acid administration improves preterm infant weight. *J Perinatol.* 2009; 29, 428–432.
54. Senterre T, Rigo J. Optimizing early nutritional support based on recent recommendations in VLBW infants and postnatal growth restriction. *J Pediatr Gastroenterol Nutr.* 2011; 53, 536–542.
55. Rochow N, Fusch G, Muhlninghaus A, et al. A nutritional program to improve outcome of very low birth weight infants. *Clin Nutr.* 2012; 31, 124–131.
56. Radmacher PG, Rafail ST, Adamkin DH. Nutrition and growth in VLBW infants with and without bronchopulmonary dysplasia. *Neonatal Intensive Care.* 2004; 16, 22–26.
57. Geary C, Caskey M, Fonseca R, Malloy M. Decreased incidence of bronchopulmonary dysplasia after early management changes, including surfactant and nasal continuous positive airway pressure treatment at delivery, lowered oxygen saturation goals, and early amino acid administration: a historical cohort study. *Pediatrics.* 2008; 121, 89–96.
58. Ehrenkranz RA, Das A, Wraga LA, et al. Early nutrition mediates the influence of severity of illness on extremely LBW infants. *Pediatr Res.* 2011; 69, 522–529.
59. Ibrahim HM, Jeroudi MA, Baier RJ, Dhanireddy R, Krouskop RW. Aggressive early total parental nutrition in low-birth-weight infants. *J Perinatol.* 2004; 24, 482–486.
60. Clark RH, Chace DH, Spitzer AR, et al. Effects of two different doses of amino acid supplementation on growth and blood amino acid levels in premature neonates admitted to the neonatal intensive care unit: a randomized, controlled trial. *Pediatrics.* 2007; 120, 1286–1296.
61. Tan MJ, Cooke RW. Improving head growth in very preterm infants – a randomised controlled trial I: neonatal outcomes. *Arch Dis Child Fetal Neonatal Ed.* 2008; 93, F337–F341.
62. Wahlqvist ML, Tienboon P. Growth and ageing. In *Nutrition and Metabolism*, 2nd edn (eds. Lanham-New SA, Macdonald IA, Roche HM), 2011; pp. 119–154. Wiley-Blackwell: Chichester.
63. de Boo HA, Harding JE. Protein metabolism in preterm infants with particular reference to intrauterine growth restriction. *Arch Dis Child Fetal Neonatal Ed.* 2007; 92, F315–F319.
64. Arslanoglu S, Moro GE, Ziegler EE. Preterm infants fed fortified human milk receive less protein than they need. *J Perinatol.* 2009; 29, 489–492.
65. Agostoni C, Buonocore G, Carnielli V, et al. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2010; 50, 85–91.
66. Henriksen C, Westerberg AC, Ronnestad A, et al. Growth and nutrient intake among very-low-birth-weight infants fed fortified human milk during hospitalisation. *Br J Nutr.* 2010; 102, 1179–1186.
67. Ehrenkranz RA, Younes N, Lemons JA, et al. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics.* 1999; 104, 280–289.
68. Cormack BE, Bloomfield FH. Audit of feeding practices in babies <1200 g or 30 weeks gestation during the first month of life. *J Paediatr Child Health.* 2006; 42, 458–463.
69. Grote V, von Kries R, Closa-Monasterolo R, et al. Protein intake and growth in the first 24 months of life. *J Pediatr Gastroenterol Nutr.* 2010; 51, S117–S118.

70. Mennella JA, Ventura AK, Beauchamp GK. Differential growth patterns among healthy infants fed protein hydrolysate or cow-milk formulas. *Pediatrics*. 2011; 127, 110–118.
71. Kuschel C, Harding J. Multicomponent fortified human milk for promoting growth in preterm infants. *Cochrane Database Syst Rev*. 2004, CD000343.
72. Kuschel CA, Harding JE. Protein supplementation of human milk for promoting growth in preterm infants. *Cochrane Database Syst Rev*. 2000, CD000433.
73. Lucas A, Morley R, Cole TJ, *et al*. Early diet in preterm babies and developmental status at 18 months. *Lancet*. 1990; 335, 1477–1481.
74. Henderson G, Fahey T, McGuire W. Nutrient-enriched formula versus standard term formula for preterm infants following hospital discharge. 2007, CD004696.
75. Jeon GW, Jung YJ, Koh SY, *et al*. Preterm infants fed nutrient-enriched formula until 6 months show improved growth and development. *Pediatr Int*. 2011; 53, 683–688.
76. Fewtrell MS, Morley R, Abbott RA, *et al*. Catch-up growth in small-for-gestational-age term infants: a randomized trial. *Am J Clin Nutr*. 2001; 74, 516–523.
77. Morley R, Fewtrell MS, Abbott RA, *et al*. Neurodevelopment in children born small for gestational age: a randomized trial of nutrient-enriched versus standard formula and comparison with a reference breastfed group. *Pediatrics*. 2004; 113, 515–521.
78. Kuschel CA, Harding JE. Fat supplementation of human milk for promoting growth in preterm infants. *Cochrane Database Syst Rev*. 2000, CD000341.
79. Henriksen C, Haugholt K, Lindgren M, *et al*. Improved cognitive development among preterm infants attributable to early supplementation of human milk with docosahexaenoic acid and arachidonic acid. *Pediatrics*. 2008; 121, 1137–1145.
80. Makrides M, Gibson RA, McPhee AJ, *et al*. Neurodevelopmental outcomes of preterm infants fed high-dose docosahexaenoic acid. *JAMA*. 2009; 301, 175–182.
81. Smithers LG, Collins CT, Simmonds LA, *et al*. Feeding preterm infants milk with a higher dose of docosahexaenoic acid than that used in current practice does not influence language or behavior in early childhood: a follow-up study of a randomized controlled trial. *Am J Clin Nutr*. 2010; 91, 628–634.
82. Aimone A, Rovet J, Ward W, *et al*. Growth and body composition of human milk-fed premature infants provided with extra energy and nutrients early after hospital discharge: 1-year follow-up. *J Pediatr Gastroenterol Nutr*. 2009; 49, 456–466.
83. Belizan JM, Villar J, Bergel E, *et al*. Long-term effect of calcium supplementation during pregnancy on the blood pressure of offspring: follow up of a randomised controlled trial. *BMJ*. 1997; 315, 281–285.
84. Berry M. *The Long Term Consequences of Preterm Birth and Neonatal Growth Rates in Lambs*. PhD thesis. 2012. University of Auckland: Auckland.
85. Lucas A, Fewtrell MS, Morley R, *et al*. Randomized trial of nutrient-enriched formula versus standard formula for postdischarge preterm infants. *Pediatrics*. 2001; 108, 703–711.
86. Waber DP, Vuori-Christiansen L, Ortiz N, *et al*. Nutritional supplementation, maternal education, and cognitive development of infants at risk of malnutrition. *Am J Clin Nutr*. 1981; 34, 807–813.
87. Plagemann A, Harder T, Franke K, Kohlhoff R. Long-term impact of neonatal breast-feeding on body weight and glucose tolerance in children of diabetic mothers. *Diabetes Care*. 2002; 25, 16–22.
88. Rodekamp E, Harder T, Kohlhoff R, *et al*. Long-term impact of breast-feeding on body weight and glucose tolerance in children of diabetic mothers. *Diabetes Care*. 2005; 28, 1457–1462.
89. Gentz J, Persson B, Kellum M, Bengtsson G, Thorell J. Effect of feeding on intravenous glucose tolerance and insulin response in piglets during the first day of life. *Life Sci*. 1971; 10, 137–144.
90. Padidela R, Patterson M, Sharief N, Ghatei M, Hussain K. Elevated basal and post-feed glucagon-like peptide 1 (GLP-1) concentrations in the neonatal period. *Eur J Endocrinol*. 2009; 160, 53–58.
91. Lucas A, Boyes S, Bloom SR, Aynsley-Green A. Metabolic and endocrine responses to a milk feed in six-day-old term infants: differences between breast and cow's milk formula feeding. *Acta Paediatr*. 1981; 70, 195–200.
92. Hans DM, Pylipow M, Long JD, Thureen PJ, Georgieff MK. Nutritional practices in the neonatal intensive care unit: analysis of a 2006 neonatal nutrition survey. *Pediatrics*. 2009; 123, 51–57.
93. Huh SY, Rifas-Shiman SL, Zera CA, *et al*. Delivery by caesarean section and risk of obesity in preschool age children: a prospective cohort study. *Arch Dis Child*. 2012; 97, 610–616.
94. Bloomfield FH, Knight DB, Harding JE. Side effects of 2 different dexamethasone courses for preterm infants at risk of chronic lung disease: a randomized trial. *J Pediatr*. 1998; 133, 395–400.
95. Hays SP, Smith EO, Sunehag AL. Hyperglycemia is a risk factor for early death and morbidity in extremely low birth-weight infants. *Pediatrics*. 2006; 118, 1811–1818.
96. Alexandrou G, Skiöld B, Karlén J, *et al*. Early hyperglycemia is a risk factor for death and white matter reduction in preterm infants. *Pediatrics*. 2010; 125, e584–e591.
97. Alsweiler JM, Kuschel CA, Bloomfield FH. Survey of the management of neonatal hyperglycaemia in Australasia. *J Paediatr Child Health*. 2007; 43, 632–635.
98. Alsweiler JM, Harding JE, Bloomfield FH. Tight glycemic control with insulin in hyperglycemic preterm babies: a randomized controlled trial. *Pediatrics*. 2012; 129, 639–647.
99. Lucas A, Morley R, Cole TJ. Adverse neurodevelopmental outcome of moderate neonatal hypoglycaemia. *BMJ*. 1988; 297, 1304–1308.
100. Knops N, Sneeuw K, Brand R, *et al*. Catch-up growth up to ten years of age in children born very preterm or with very low birth weight. *BMC Pediatr*. 2005; 5, 26.
101. Singhal A, Cole TJ, Fewtrell M, *et al*. Promotion of faster weight gain in infants born small for gestational age: is there an adverse effect on later blood pressure? *Circulation*. 2007; 115, 213–220.

102. Singhal A, Fewtrell M, Cole TJ, Lucas A. Low nutrient intake and early growth for later insulin resistance in adolescents born preterm. *Lancet*. 2003; 361, 1089–1097.
103. Singhal A, Cole TJ, Fewtrell M, Deanfield J, Lucas A. Is slower early growth beneficial for long-term cardiovascular health? *Circulation*. 2004; 109, 1108–1113.
104. Hemachandra AH, Howards PP, Furth SL, Klebanoff MA. Birth weight, postnatal growth, and risk for high blood pressure at 7 years of age: results from the Collaborative Perinatal Project. *Pediatrics*. 2007; 119, e1264–e1270.
105. Ong KK, Loos RJF. Rapid infancy weight gain and subsequent obesity: systematic reviews and hopeful suggestions. *Acta Paediatr*. 2006; 95, 904–908.
106. Ong KK. Catch-up growth in small for gestational age babies: good or bad? *Curr Opinion Endocrinol Diabetes Obes*. 2007; 14, 30–34.
107. Lucas A, Fewtrell MS, Morley R, et al. Randomized outcome trial of human milk fortification and developmental outcome in preterm infants. *Am J Clin Nutr*. 1996; 64, 142–151.
108. Franz AR, Pohlandt F, Bode H, et al. Intrauterine, early neonatal, and postdischarge growth and neurodevelopmental outcome at 5.4 years in extremely preterm infants after intensive neonatal nutritional support. *Pediatrics*. 2009; 123, e101–e109.
109. Jaquiere AL, Oliver MH, Honeyfield-Ross M, Harding JE, Bloomfield FH. Periconceptional undernutrition in sheep affects adult phenotype only in males. *J Nutr Metabol*. 2012; doi: 10.1155/2012/123610.
110. Kuschel CA, Harding JE, Kumaran VS. Calcium and phosphorus supplementation of human milk for preterm infants. *Cochrane Database Syst Rev*. 2001, CD003310.
111. Wauben IP, Atkinson SA, Grad TL, Shah JK, Paes B. Moderate nutrient supplementation of mother's milk for preterm infants supports adequate bone mass and short-term growth: a randomized, controlled trial. *Am J Clin Nutr*. 1998; 67, 465–472.