

# PERMUTATIONS RELATED TO SECANT, TANGENT AND EULERIAN NUMBERS

BY  
MORTON ABRAMSON

**1. Introduction.** It is well known that

$$(1) \quad \sum_{n=0}^{\infty} A_n \frac{x^n}{n!} = \sec x + \tan x$$

where  $A_n$  denotes the number of “up-down” or alternating permutations

$$(2) \quad e_1 < e_2 > e_3 < e_4 > \dots$$

of  $1, 2, \dots, n$ . The numbers  $A_{2n}$  and  $A_{2n+1}$  are known as the secant and tangent numbers respectively and  $A_{2n} = (-1)^n E_{2n}$ , where  $E_n$  is the Euler number.

A brief history of the numbers  $A_n$  is given by Gould [63] who notes that alternating permutations were enumerated in closed form by D. André [7] as long ago as 1879 and that a detailed exposition of the method is given by E. Netto [69, pp. 105–112] where various references in the literature are given. By first obtaining a recurrence relation and then solving a differential equation, Blundon [9] gives a short self-contained derivation of (1).

The familiar Eulerian numbers  $A_r(n)$  given explicitly by

$$(3) \quad A_r(n) = \sum_{i=0}^r (-1)^i \binom{n+1}{i} (r+1-i)^n,$$

with the generating function

$$(4) \quad 1 + \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{r=0}^{n-1} A_r(n) y^r = \frac{1-y}{e^{(y-1)x} - y},$$

enumerate those permutations  $e_1 e_2 \dots e_n$  of  $1, 2, \dots, n$  containing precisely  $r$  rises, a rise being a pair  $e_i < e_{i+1}$ ,  $i = 1, \dots, n-1$ . (Often an initial rise is considered to occur at  $e_1$  but this is not done here.) The Eulerian numbers were discussed in 1755 by Euler [54]. Discussions of the Eulerian numbers are also given by Carlitz and Scoville [38], Comtet [45], Foata and Schützenberger [56] and Riordan [72].

---

Received by the editors July 18, 1978.

The sec-tan and Eulerian numbers, their properties, combinatorial interpretations as permutation problems, related numbers and problems, generalizations, extensions and refinements have been studied extensively in the literature. Papers involving both numbers, include [22, 23, 26, 58]. Papers relating more to the sec-tan numbers include [7]–[10] incl., [17, 19, 20, 24, 25, 27, 29, 31, 32, 36, 37, 46, 49, 51, 52, 53, 55, 57, 59, 64, 74, 78]. Papers relating more to Eulerian numbers include [1] to [6] incl., [11]–[16] incl., [18, 21, 24, 28, 29, 30, 33, 34, 35], [38]–[44] incl., [47, 48, 50, 60, 62, 65, 66, 67, 68, 71, 73, 75, 79, 80, 81].

Stanley [76], [77] and Gessel [61] in his Ph.D. thesis under the supervision of Stanley unify many disparate topics in combinatorics. Included in their works are generalizations of the sec-tan and Eulerian numbers which contain as special cases many results obtained by others. More precisely, a general theory for the enumeration of order-reversing maps of finite ordered sets into chains is developed in [76], a unified method involving Möbius functions associated with binomial posets for enumerating permutations of sets and multisets is given in [77] and a general theory of enumeration of finite sequences by generating functions is developed in [61].

In this paper, we give a simple, elementary and purely combinatorial derivation of the following generalization of the sec-tan and Eulerian numbers. This is done by noting a natural relationship between the restricted permutations considered and corresponding restricted distributions of distinct objects into distinct cells.

In a permutation  $e_1 e_2 \cdots e_n$  of  $1, 2, \dots, n$  we define a run of length  $l \geq 1$  to be a sequence

$$e_{i+1} < e_{i+2} < \cdots < e_{i+l}$$

satisfying  $e_i > e_{i+1}$ ,  $e_{i+l} > e_{i+l+1}$  with  $e_0 = n + 1$ ,  $e_{n+1} = 0$ . For example, the permutation 3 4 1 6 2 7 9 8 5 contains precisely five runs,

$$3 < 4, 1 < 6, 2 < 7 < 9, 8, 5,$$

of lengths 2, 2, 3, 1, 1 respectively.

Note that a permutation of  $1, 2, \dots, n$  with precisely  $r$  runs contains precisely  $n - r$  rises and precisely  $r - 1$  falls, a fall being a pair  $e_i > e_{i+1}$ ,  $i = 1, 2, \dots, n - 1$ .

Denote by  $A_r(k, n, t_1, t_2)$ ,  $0 \leq r \leq n$ ,  $k, t_1, t_2 \geq 1$ , the number of permutations of  $1, 2, \dots, kn + t_1 + t_2$  with (a) precisely  $n + 2 - r$  runs, (b) the lengths of the first and last runs not less than  $t_1$  and  $t_2$  respectively and congruent to  $t_1 \pmod k$  and  $t_2 \pmod k$  respectively and (c) the lengths of all the other  $n - r$  runs congruent to  $0 \pmod k$ . That is,  $A_r(k, n, t_1, t_2)$  is the number of permutations of  $1, 2, \dots, kn + t_1 + t_2$  satisfying

$$(5) \quad e_1 < e_2 < \cdots < e_{m_1} > e_{m_1+1} < e_{m_1+2} < \cdots < e_{m_1+m_2} > \cdots > \cdots$$

with

$$(6) \quad t_1 \leq m_1 \equiv t_1 \pmod{k}, \quad t_2 \leq m_{n+2-r} \equiv t_2 \pmod{k}, \quad 0 < m_i \equiv 0 \pmod{k},$$

$i = 2, \dots, n + 1 - r$  where  $m_i$  is the length of the  $i$ th run,  $i = 1, 2, \dots, n + 2 - r$ .

We define  $A_{n+1}(k, n, t_1, t_2) = 1$ .

In §2 we give an explicit formula (19) for  $A_r(k, n, t_1, t_2)$  and also show that

$$(7) \quad \sum_{n=0}^{\infty} \sum_{r=0}^{n+1} A_r(k, n, t_1, t_2) y^r \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} \\ = (y-1) \left[ \frac{\varphi_{k,t_1}(x, y)\varphi_{k,t_2}(x, y)}{y - \varphi_{k,0}(x, y)} + \varphi_{k,t_1+t_2}(x, y) \right]$$

where

$$(8) \quad \varphi_{k,t}(x, y) = \sum_{j=0}^{\infty} \frac{(y-1)^j x^{t+jk}}{(t+jk)!}, \quad t = 0, 1, 2, \dots$$

The special cases  $A_0(k, n-1, k, t)$  and  $A_0(k, n-1, k, k)$  noted in §4 are results of Carlitz [25]. These special cases involve the Olivier functions  $\varphi_{k,t}(x, 0)$  defined by (8) with  $y = 0$ . See Carlitz [10] for some arithmetic properties of these functions.

**2. Distributions of  $1, 2, \dots, nk + t_1 + t_2$  into distinct cells.** Denote by  $g(k, n, t_1, t_2, p)$ ,  $n \geq p - 2 \geq 0$ ,  $t_1 \geq 1$ ,  $t_2 \geq 1$ , the number of distributions.

$$(9) \quad e_1 < e_2 < \dots < e_{m_1} \mid e_{m_1+1} < \dots < e_{m_1+m_2} \mid \dots \mid \dots < e_{kn+t_1+t_2} \\ \text{cell 1} \quad \text{cell 2} \quad \dots \quad \text{cell } p$$

of the integers  $1, 2, \dots, kn + t_1 + t_2$  into  $p$  distinct cells such that

$$(10) \quad t_1 \leq m_1 \equiv t_1 \pmod{k}, \quad t_2 \leq m_p \equiv t_2 \pmod{k} \quad \text{and} \quad 0 < m_i \equiv 0 \pmod{k}$$

for  $i = 2, \dots, p - 1$ , where  $m_i$  denotes the number of objects in cell  $i$ . Integers in a cell are arranged in rising order. Then

$$(11) \quad g(k, n, t_1, t_2, p) \\ = \sum (kn + t_1 + t_2)! / (i_1 k + t_1)! (i_2 k)! (i_3 k)! \dots (i_{p-1} k)! (i_p k + t_2)!$$

where the summation is over all  $i_1 \geq 0$ ,  $i_p \geq 0$ ,  $i_2 > 0$ ,  $i_3 > 0, \dots, i_{p-1} > 0$  satisfying  $i_1 + \dots + i_p = n$  and the exponential generating function is

$$(12) \quad \sum_{n=p-2}^{\infty} g(k, n, t_1, t_2, p) \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} \\ = \left( \sum_{i=1}^{\infty} \frac{x^{ik}}{(ik)!} \right)^{p-2} \sum_{j=0}^{\infty} \frac{x^{t_1+jk}}{(t_1+jk)!} \sum_{l=0}^{\infty} \frac{x^{t_2+lk}}{(t_2+lk)!}$$

We define  $g(k, n, t_1, t_2, 1) = 1$ .

In a distribution (9) satisfying (10) define the distribution property  $P_i$  to be, “ $e_{t_1+ik}$  and  $e_{t_1+ik+1}$  are in the same cell”,  $i = 0, 1, 2, \dots, n$ .

LEMMA. *The number of distributions (9) satisfying (10) and satisfying precisely  $j$  of the  $n + 1$  distribution properties  $P_0, P_1, \dots, P_n$  is  $g(k, n, t_1, t_2, n + 2 - j)$ .*

**Proof.** Suppose a distribution (9) satisfying (10) satisfies precisely  $j$  of the  $n + 1$  properties with cell  $i$  satisfying precisely  $j_i (\geq 0)$  of the properties. Then cell  $i$  contains  $(j_i + 1)k$  integers,  $i = 2, \dots, p - 1$ , cell 1 contains  $j_1 k + t_1$  integers and cell  $p$  contains  $j_p k + t_2$  integers. Since  $j_1 + \dots + j_p = j$  and  $(j_1 k + t_1) + (j_2 + 1)k + (j_3 + 1)k + \dots + (j_{p-1} + 1)k + (j_p k + t_2) = kn + t_1 + t_2$  it follows that  $p = n + 2 - j$ .

3. **The numbers  $A_r(k, n, t_1, t_2)$ .** There are  $(kn + t_1 + t_2)! / (k!)^n t_1! t_2!$  linear arrangements

$$(13) \quad e_1 e_2 \cdots e_{kn+t_1+t_2}$$

of  $1, 2, \dots, kn + t_1 + t_2$ ,  $n \geq 0$ ,  $k, t_1, t_2 \geq 1$  satisfying

$$(14) \quad \begin{aligned} & e_1 < e_2 < \cdots < e_{t_1}, \\ & e_{t_1+1} < e_{t_1+2} < \cdots < e_{t_1+k} \\ & e_{t_1+k+1} < e_{t_1+k+2} < \cdots < e_{t_1+k2} \\ & \vdots \\ & e_{t_1+k(n-1)+1} < e_{t_1+k(n-1)+2} < \cdots < e_{t_1+kn} \\ & e_{t_1+kn+1} < e_{t_1+kn+2} < \cdots < e_{t_1+kn+t_2}. \end{aligned}$$

Denote by  $A_r(k, n, t_1, t_2)$  the number of arrangements (13) satisfying (14) and satisfying precisely  $r$  of the  $n + 1$  arrangement properties

$$(15) \quad e_{t_1} < e_{t_1+1}, e_{t_1+k} < e_{t_1+k+1}, e_{t_1+k2} < e_{t_1+k2+1}, \dots, e_{t_1+kn} < e_{t_1+kn+1}$$

denoted by  $P_0, P_1, \dots$ , respectively.

Denote by  $N(P_{u_1}, P_{u_2}, \dots, P_{u_j})$  the number of arrangements (13) satisfying (14) and satisfying the  $j$  arrangement properties  $P_{u_1}, \dots, P_{u_j}$  (and possibly other of the properties). Let

$$(16) \quad s(j) = \sum N(P_{u_1}, P_{u_2}, \dots, P_{u_j})$$

where the summation is taken over all  $j$ -combinations  $0 \leq u_1 < u_2 < \dots < u_j \leq n$ . By the general principle of inclusion and exclusion we have

$$(17) \quad A_r(k, n, t_1, t_2) = \sum_{j=0}^{n+1-r} (-1)^j \binom{r+j}{j} s(r+j)$$

with  $s(0) = (kn + t_1 + t_2)! / (k!)^n t_1! t_2!$ . We now show that

$$(18) \quad s(j) = g(k, n, t_1, t_2, n + 2 - j).$$

**Proof of (18).** Upon removing the bars of a distribution (9) satisfying (10) and also satisfying precisely the  $j$  distribution properties  $P_{u_1}, \dots, P_{u_j}$  we obtain an arrangement counted by  $N(P_{u_1}, \dots, P_{u_j})$  where the arrangement properties  $P_{u_1}, \dots, P_{u_j}$  correspond to the distribution properties  $P_{u_1}, \dots, P_{u_j}$ . Further every arrangement counted by  $N(P_{u_1}, \dots, P_{u_j})$  can be obtained this way. Using the Lemma in §2, (18) follows.

Hence, (17) and (18) give the formula

$$(19) \quad A_r(k, n, t_1, t_2) = \sum_{j=0}^{n+1-r} (-1)^j \binom{r+j}{j} g(k, n, t_1, t_2, n+2-r-j),$$

and the ordinary generating function is

$$(20) \quad \sum_{r=0}^{n+1} A_r(k, n, t_1, t_2) y^r = \sum_{j=0}^{n+1} g(k, n, t_1, t_2, n+2-j) (y-1)^j.$$

**Proof of (7).** From (20), the left side of (7)

$$\begin{aligned} &= \sum_{n=0}^{\infty} (y-1)^{n+1} \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} + \sum_{n=0}^{\infty} \sum_{m=2}^{n+2} g(k, n, t_1, t_2, m) (y-1)^{n+2-m} \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} \\ &= (y-1) \varphi_{k,t_1+t_2}(x, y) + \sum_{m=2}^{\infty} \sum_{n=m-2}^{\infty} g(k, n, t_1, t_2, m) \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} (y-1)^{n+2-m} \\ &= (y-1) \varphi_{k,t_1+t_2}(x, y) + \sum_{m=2}^{\infty} (\varphi_{k,0}(x, y) - 1)^{m-2} \\ &\quad \times \varphi_{k,t_1}(x, y) \varphi_{k,t_2}(x, y) (y-1)^{2-m} \text{ (by (12))} \\ &= (y-1) \varphi_{k,t_1+t_2}(x, y) + \frac{(y-1)}{y - \varphi_{k,0}(x, y)} \varphi_{k,t_1}(x, y) \varphi_{k,t_2}(x, y) \\ &= \text{right side of (7)}. \end{aligned}$$

We also note the relation

$$\begin{aligned} A_r(k, n, t_1, t_2) - A_{r-1}(k, n-1, t_1, k+t_2) \\ = \binom{kn+t_1+t_2}{t_2} A_r(k, n-1, t_1, k) - A_r(k, n-1, t_1, k+t_2) \end{aligned}$$

where both sides are equal to the number of permutations counted by  $A_r(k, n, t_1, t_2)$  with the last run of length exactly  $t_2$ .

**4. Special Cases.** The special case  $A_0(k, n, t_1, t_2)$  counts those permutations of  $1, 2, \dots, kn+t_1+t_2$  with the first and last runs of lengths  $t_1$  and  $t_2$  respectively and each of the other runs of length  $k$ . By (7), with  $y=0$ , we have

$$(21) \quad \sum_{n=0}^{\infty} A_0(k, n, t_1, t_2) \frac{x^{kn+t_1+t_2}}{(kn+t_1+t_2)!} = \frac{\varphi_{k,t_1}(x, 0) \varphi_{k,t_2}(x, 0)}{\varphi_{k,0}(x, 0)} + \varphi_{k,t_1+t_2}(x, 0).$$

**Eulerian numbers.** The number of permutations of  $1, 2, \dots, n$  containing precisely  $r$  rises (or precisely  $n - r$  runs) is the well-known Eulerian number  $A_r(n) = A_r(1, n - 2, 1, 1)$ . Noting that

$$(22) \quad \varphi_{1,1}(x, y) = \frac{e^{(y-1)x} - 1}{y - 1},$$

and by (7), we obtain the known generating function (4). From (12) it follows that  $g(1, n - 2, 1, 1, p) = p!S(n, p)$  where

$$S(n, p) = \sum_{i=0}^{p-1} (-1)^i \binom{p}{i} (p - i)^n / p!$$

is the Stirling number of the second kind. By (19), use of a simple identity, and the fact that  $A_r(n)$  has the symmetric property  $A_r(n) = A_{n-1-r}(n)$  we obtain the familiar explicit formula

$$(23) \quad \begin{aligned} A_r(n) &= \sum_{i=0}^{n-1-r} (-1)^i \binom{r+i}{r} (n-r-i)! S(n, n-r-i) \\ &= \sum_{i=0}^{n-1-r} (-1)^i \binom{n+1}{i} (n-r-i)^n. \\ &= \sum_{i=0}^r (-1)^i \binom{n+1}{i} (r+1-i)^n. \end{aligned}$$

**The numbers  $A_r(k, n, k, t)$ .** Using notation similar to that of Carlitz [25] we let  $A_{r,k,t}(kn+t) = A_r(k, n-1, k, t)$  and define  $A_{0,k,t}(k0+t) = 1$ . Then, for  $0 \leq r \leq n-1$ ,  $A_{r,k,t}(kn+t)$  is the number of permutations of  $1, 2, \dots, kn+t$  with (a) exactly  $n+1-r$  runs and (b) the length of each of the first  $n-r$  runs congruent to  $0 \pmod k$  and the length of the last run not less than  $t$  and congruent to  $t \pmod k$ . Let  $A_{n,k,t}(kn+t) = 1$ . Letting  $t_1 = k$  and  $t_2 = t$  in (7) and noting that

$$(24) \quad \begin{aligned} \varphi_{k,k}(x, y) &= \frac{\varphi_{k,0}(x, y) - 1}{y - 1} \\ \varphi_{k,k+t}(x, y) &= \frac{1}{y-1} \left( \varphi_{k,t}(x, y) - \frac{x^t}{t!} \right) \end{aligned}$$

we obtain

$$(25) \quad \sum_{n=0}^{\infty} \sum_{r=0}^n A_{r,k,t}(kn+t) y^r \frac{x^{kn+t}}{(kn+t)!} = \frac{(y-1)\varphi_{k,t}(x, y)}{y - \varphi_{k,0}(x, y)}.$$

Letting  $t = k$  in (25) we obtain

$$(26) \quad 1 + \sum_{n=0}^{\infty} \sum_{r=0}^n A_{r,k,k}(kn+k) y^r \frac{x^{kn+k}}{(kn+k)!} = \frac{y-1}{y - \varphi_{k,0}(x, y)}$$

with  $A_{r,k,k}(kn+k)$  being the number of permutations of  $1, 2, \dots, k(n+1)$  having exactly  $n+1-r$  runs and with the length of each run a positive multiple of  $k$ .

In the case  $r=0$ , by (21) or (25), we obtain

$$(27) \quad \sum_{n=0}^{\infty} A_{0,k,t}(kn+t) \frac{x^{kn+t}}{(kn+t)!} = \frac{\varphi_{k,t}(x, 0)}{\varphi_{k,0}(x, 0)}$$

and letting  $t=k$  in (27) we have

$$(28) \quad 1 + \sum_{n=0}^{\infty} A_{0,k,k}(kn+k) \frac{x^{kn+k}}{(kn+k)!} = \frac{1}{\varphi_{k,0}(x, 0)},$$

where  $A_{0,k,t}(kn+t)$  is the number of permutation of  $1, 2, \dots, kn+t$  with the last run of length  $t$  and each of the other runs of length  $k$ . Both (27) and (28) are results of Carlitz [22] and [25] where he notes that  $\varphi_{k,t}(x, 0)$  are the Olivier functions. See also Gessel [16, p. 51, example 2] and Stanley [74].

**Permutations with runs of even length.** The number of permutations of  $1, 2, \dots, 2n+2$  with precisely  $n+1-r$  runs and with each of the runs of even length is  $A_{r,2,2}(2n+2)$  and by (26),

$$(29) \quad 1 + \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} A_{r,2,2}(2n+2) y^r \frac{x^{2n+2}}{(2n+2)!} = \frac{y-1}{y-\varphi_{2,0}(x, y)} \\ = \frac{y-1}{y-\cos(x\sqrt{1-y})}.$$

The number of permutations of  $1, 2, \dots, 2n+1$  with precisely  $n+1-r$  runs and with each of the first  $n-r$  runs of even length and the last run of odd length is  $A_{r,2,1}(2n+1)$  with  $A_{n,2,1}(2n+1)=1$  and by (25) it follows that

$$(30) \quad \sum_{n=0}^{\infty} \sum_{r=0}^n A_{r,2,1}(2n+1) y^r \frac{x^{2n+1}}{(2n+1)!} = \frac{(y-1)\varphi_{2,1}(x, y)}{y-\varphi_{2,0}(x, y)} \\ = \frac{\sqrt{(1-y)}\sin(x\sqrt{1-y})}{\cos(x\sqrt{1-y})-y}.$$

Let  $A_{r,n}$  denote the number of permutations of  $1, 2, \dots, n$  with precisely  $[(n+1)/2]-r$  runs and with each of even length if  $n$  is even and all but the last run even if  $n$  is odd. By combining (29) and (30) we obtain

$$(31) \quad 1 + \sum_{n=1}^{\infty} \sum_{r=0}^{(n-1)/2} A_{r,n} y^r \frac{x^n}{n!} = (y-1) \frac{(1+\varphi_{2,1}(x, y))}{(y-\varphi_{2,0}(x, y))}$$

**Up-down permutations.** The cases  $A_{0,2,2}(2n+2)$  and  $A_{0,2,1}(2n+1)$  are the numbers of up-down permutations for the even and odd cases respectively and

(29) and (30) reduce, respectively, to

$$(32) \quad 1 + \sum_{n=0}^{\infty} A_{0,2,2}(2n+2) \frac{x^{2n+2}}{(2n+2)!} = \secant x$$

and

$$(33) \quad \sum_{n=0}^{\infty} A_{0,2,1}(2n+1) \frac{x^{2n+1}}{(2n+1)!} = \text{tangent } x.$$

Combining (32) and (33) and defining  $A_0 = 1$  we obtain (1).

#### REFERENCES

1. Morton Abramson, *Some generalized Eulerian numbers*, Proc. Fifth Southeastern Conference on Combinatorics, graph theory and computing, Florida Atlantic University, Boca Raton, Florida, 1974, pp. 159–172.
2. Morton Abramson, *A simple solution of Simon Newcomb's problem*, J. of Comb. Theory, "A", **18** (1975), pp. 223–225.
3. Morton Abramson, *A note on permutations with fixed pattern*, J. of Comb. Theory, "A", **19** (1975), pp. 237–239.
4. Morton Abramson, *Enumeration of sequences by levels and rises*, Discrete Math., **12** (1975), pp. 101–112.
5. Morton Abramson, *Sequences by number of w-rises*, Canad. Math. Bull., **18** (1975), pp. 317–319.
6. Morton Abramson and David Promislow, *Enumeration of arrays by column rises*, J. of Comb. Theory, "A", **24** (1978), pp. 247–250.
7. D. André, *Développements de sec x et de tan x*, C. R. Acad. Sci. Paris, **88** (1879), pp. 965–967.
8. D. André, *Sur les permutations alternées*, J. Math. Pures Appl., **7** (1881), p. 167.
9. W. J. Blundon, *Solution to "A permutation problem"*, 4755 [1957, 596], Amer. Math. Monthly, **65** (1958), p. 533.
10. L. Carlitz, *Some arithmetic properties of the Oliver functions*, Math. Ann., **128** (1955), pp. 412–419.
11. L. Carlitz, *Eulerian numbers and polynomials*, Math. Magazine, **32** (1959), pp. 247–260.
12. L. Carlitz, *Eulerian numbers and polynomials of higher order*, Duke Math. J., **27** (1960), pp. 401–423.
13. L. Carlitz, *A note on Eulerian numbers*, Arch. Math., **14** (1963), pp. 383–390.
14. L. Carlitz, *Extended Bernoulli and Eulerian numbers*, Duke Math. J., **31** (1964), pp. 667–689.
15. L. Carlitz, *Enumeration of sequences by rises and falls: a refinement of the Simon Newcomb problem*, Duke Math. J., **39** (1972), pp. 267–280.
16. L. Carlitz, *Eulerian numbers and operators*, The Theory of Arithmetic Functions, pp. 65–70, Lecture Notes in Math., Vol. **251**, Springer, Berlin, 1972.
17. Leonard Carlitz, *Enumeration of a special class of permutations by rises*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz., No. **412–460** (1973), pp. 189–196.
18. L. Carlitz, *Enumeration of permutations by rises and cycle structure*, J. Reine Angew. Math., **262/263** (1973), pp. 220–233.
19. L. Carlitz, *Enumeration of up-down permutations by number of rises*, Pacific J. Math., **45** (1973), pp. 49–58.
20. L. Carlitz, *Enumeration of up-down sequences*, discrete Math., **4** (1973), pp. 273–286.
21. L. Carlitz, *Eulerian numbers and operators*, collect. Math., **24** (1973) pp. 175–200.
22. L. Carlitz, *Permutations with prescribed pattern*, Math. Nachr., **58** (1973), 31–53.



23. L. Carlitz, *Permutations and sequences*, *Advances in Math.*, **14** (1974), pp. 92–120.
24. L. Carlitz, *A combinatorial property of  $q$ -Eulerian numbers*, *Amer. Math. Monthly*, **82** (1975), pp. 51–54.
25. L. Carlitz, *Generating functions for a special class of permutations*, *Proc. Amer. Math. Soc.*, **47** (1975), pp. 251–256.
26. L. Carlitz, *Permutations, sequences, and special functions*, *SIAM Rev.*, **17** (1975), pp. 298–321.
27. L. Carlitz, *Combinatorial property of a special polynomial sequence*, *Canad. Math. Bull.*, **20** (1977), pp. 183–188.
28. L. Carlitz, D. C. Kurtz, R. Scoville, and O. P. Stackelberg, *Asymptotic properties of Eulerian numbers*, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, **23** (1972), pp. 47–54.
29. L. Carlitz and J. Riordan, *Congruences for Eulerian Numbers*, *Duke Math. J.*, **20** (1953), pp. 339–343.
30. L. Carlitz, D. P. Roselle and R. A. Scoville, *Permutations and sequences with repetitions by number of increases*, *J. Comb. Theory*, **1** (1966), pp. 350–374.
31. L. Carlitz and Richard Scoville, *Tangent numbers and operators*, *Duke Math. J.*, **39** (1972), pp. 413–429.
32. L. Carlitz and Richard Scoville, *Up-down sequences*, *Duke Math. J.*, **39** (1972), pp. 583–598.
33. L. Carlitz and Richard Scoville, *Enumeration of rises and falls by position*, *Discrete Math.*, **5**, (1973), pp. 45–59.
34. L. Carlitz and Richard Scoville, *Enumeration of permutations by rises, falls, rising maxima, and falling maxima*, *Acta Math. Acad. Sci. Hungar.*, **25** (1974), pp. 269–277.
35. L. Carlitz and Richard Scoville, *Generalized Eulerian numbers: combinatorial applications*, *J. Reine Angew. Math.*, **265** (1974), pp. 110–137.
36. L. Carlitz and Richard Scoville, *Enumeration of up-down permutations by upper records*, *Monatshefte für Mathematik*, **79** (1975), pp. 3–12.
37. L. Carlitz and Richard Scoville, *Generating functions for certain types of permutations*, *J. Combinatorial Theory “A”*, **18** (1975), pp. 262–275.
38. L. Carlitz and Richard Scoville, *Eulerian numbers and operators*, *Fibonacci Quart.*, **13** (1975), pp. 71–83.
39. L. Carlitz and Richard Scoville, *Some permutation problems*, *J. Comb. Theory “A”*, **22** (1977), pp. 129–145.
40. L. Carlitz, Richard Scoville and Theresa Vaughan, *Enumeration of permutations and sequences with restrictions*, *Duke Math. J.*, **40** (1973), pp. 723–741.
41. L. Carlitz, R. Scoville and T. Vaughan, *Enumeration of pairs of permutations and sequences*, *Bull. Amer. Soc.*, **80** (1974), pp. 881–884.
42. L. Carlitz, R. Scoville, and T. Vaughan, *Enumeration of pairs of permutations*, *Discrete Math.*, **14** (1976), pp. 215–239.
43. L. Carlitz, R. Scoville and T. Vaughan, *Enumeration of pairs of sequences by rises, falls, and levels*, *Manuscripta Math.*, **19** (1976), pp. 211–243.
44. L. Carlitz and T. Vaughan, *Enumeration of sequences of given specification according to rises, falls, and maxima*, *Discrete Math.*, **8** (1974), pp. 147–167.
45. L. Comtet, *Analyse Combinatoire*, vol. **1**, pp. 63–64, vol. **2**, pp. 82–86, Paris, 1970.
46. N. E. Debruijn, *Permutations with given ups and downs*, *Nieuw Archief voor Wiskunde*, **18** (1970), pp. 61–65.
47. J. F. Dillon and D. P. Roselle, *Eulerian numbers of higher order*, *Duke Math. J.*, **35** (1968), pp. 247–256.
48. J. F. Dillon and D. P. Roselle, *Simon Newcomb’s problem*, *SIAM J. Appl. Math.*, **17** (1969), pp. 1086–1097.
49. Robert Donaghey, *Alternating permutations and binary increasing trees*, *J. Comb. Theory, “A”*, **18** (1975), pp. 141–148.
50. F. B. Drane and D. P. Roselle, *A sequence of polynomials related to the Eulerian polynomials*, *Utilitas Math.*, **9** (1976), pp. 33–37.

51. Dominique Dumont, *Interprétations combinatoires des nombres de Genocchi*, Duke Math. J., **41** (1974), pp. 305–318.
52. R. C. Entringer, *A combinatorial interpretation of the Euler and Bernoulli numbers*, Nieuw Arch. V. Wiskunde, **14** (1966), pp. 241–246.
53. R. C. Entringer, *A note on enumeration of permutations of  $(1, \dots, n)$  by number of maxima*, Gac. Mat. (Madrid), **23** (1971), pp. 67–69.
54. L. Euler, *Institutiones Calculi Differentialis*, St. Petersburg, 1755.
55. Dominique Foata, *Groupes de réarrangements et nombres d'Euler*, C. R. Acad. Sci. Paris Sér. A-B, **275** (1972), A 1147-A 1150.
56. Dominique Foata and Marcel-P. Schützenberger, *Théorie géométrique des polynômes eulériens*, in “Lecture Notes in Mathematics,” No. **138**, Springer-Verlag, New York, 1970.
57. D. Foata and M. P. Schützenberger, *Nombres d'Euler et permutations alternantes*, A Survey of Combinatorial Theory, pp. 173–187, North-Holland, Amsterdam, 1973.
58. Dominique Foata and Volker Strehl, *Euler numbers and variations of permutations*, Colloquio Internazionale sulle Teorie Combinatorie (Roma, 1973), Tomo **I**, pp. 119–131. Atti dei Convegni Lincei, No. **17**, Accad. Naz. Lincei, Roma, 1976.
59. H. O. Foulkes, *Enumeration of permutations with prescribed up-down and inversion sequences*, Discrete Math., **15** (1976), pp. 235–252.
60. H. O. Foulkes, *A nonrecursive combinatorial rule for Eulerian numbers*, J. Comb. Theory, “A”, **22** (1977), pp. 246–248.
61. Ira Martin Gessel, *Generating Functions and Enumeration of sequences*, Ph.d. Thesis, M.I.T. 1977.
62. Ira Gessel and Richard P. Stanley, *Stirling Polynomials*, J. of Comb. Theory, **24** (1978), pp. 24–33.
63. H. W. Gould, *Comment on Problem 67-5, “up-down” permutations*, SIAM Review, **10** (1968), pp. 225–226.
64. H. W. Gould, *Explicit formulas for Bernoulli numbers*, Amer. Math. Monthly, **79** (1972), pp. 44–51.
65. Allan Wm. Johnson, Jr., *Solution to, “A difference equation in two variables,” Problem E 2609 [1976, 567]*, Amer. Math. Monthly, **84** (1977), pp. 826–827.
66. Robert E. Knop, *A note on hypercube partitions*, J. Comb. Theory, “A”, **15** (1973), pp. 338–342.
67. Germain Kreweras, *Sur une extension du problème dit “de Simon Newcomb”*, C.R. Acad. Sci. Paris Sér. A-B, **263** (1966), pp. A43–A45.
68. David C. Kurtz, *A note on concavity properties of triangular arrays of numbers*, J. of Comb. Theory “A”, **13** (1972), pp. 135–139.
69. E. Netto, *Lehrbuch der Combinatorik*, Berlin, 1927.
70. L. Olivier, *Bemerkungen über eine Art von Funktionen, welche Eigenschaften haben, wie die Cosinus und Sinus*, J. Reine Angew. Math., **2** (1827), pp. 243–251.
71. J. Riordan, *Triangular permutation numbers*, Proc. Amer. Math. Soc., **2** (1951), pp. 429–432.
72. J. Riordan, *An Introduction to Combinatorial Analysis*, J. Wiley, New York, 1958.
73. D. P. Roselle, *Permutations by number of rises and successions*, Proc. Amer. Math. Soc., **19** (1968), pp. 8–16.
74. J. Rosen, *The number of product-weighted lead codes for ballots and its relation to the Ursell-functions of the Linear Ising Model*, J. Comb. Theory, “A”, **20** (1976), p. 377.
75. B. Sorin, *La distribution combinatoire du nombre de différences premières positives*, Rev. Inst. Internat. Statist., **39** (1971), pp. 9–20.
76. Richard P. Stanley, *Ordered Structures and Partitions*, Memoirs of the Amer. Math. Soc., No. **119**, Amer. Math. Soc., Providence, R. I., 1972.
77. Richard P. Stanley, *Binomial posets, Möbius inversion, and permutation enumeration*, J. Combinatorial Theory “A”, **20** (1976), pp. 336–356.
78. Volker Strehl, *Enumeration of alternating permutations according to peak sets*, J. of Comb. Theory, “A”, **24** (1978), pp. 238–240.

79. S. Tanny, *A probabilistic interpretation of Eulerian numbers*, *Duke Math. J.*, **40** (1973), pp. 717–722.

80. Don West, *Solution to “The Smith college diploma problem,” Problem E 2404 [1973, 316]*, *Amer. Math. Monthly*, **81** (1974), pp. 286–289.

81. J. Worpitzky, *Studien über die Bernoullischen und Eulerschen Zahlen*, *J. für die reine und angewandte Math.*, **94** (1883), pp. 203–232.

DEPARTMENT OF MATHEMATICS

YORK UNIVERSITY  
4700 KELLE STREET  
DOWNSVIEW, ONT.  
M3J 1P3.