

Applying welfare science to bottlenose dolphins (*Tursiops truncatus*)

ILK Clegg^{*†‡}, CE Van Elk[§] and F Delfour^{†‡}

[†] Laboratoire d'Éthologie Expérimentale et Comparée EA4443, Université Paris 13, Villetaneuse, France

[‡] Equipe Delphinarium, Parc Astérix, BP8, 60128 Plailly, France

[§] Dolfinarium Harderwijk, Strandboulevard 1, 3841 AB Harderwijk, The Netherlands

* Contact for correspondence and requests for reprints: isabella.clegg@leec.univ-paris13.fr

Abstract

Animal welfare science is a burgeoning field, but research on cetaceans (whales, dolphins and porpoises) is lacking. Bottlenose dolphins (*Tursiops truncatus*) are the most well-known and studied cetaceans, particularly in captivity, and thus are used in this review as a model for other cetacean species. Despite the public interest and need for such research, studies specifically investigating dolphin welfare are lacking. This review uses the three broad categories of behaviour, health, and cognition, to discuss how dolphin welfare has been assessed thus far, and could be assessed in future. We present welfare indicators validated in other species that could be applied to dolphins, including innovative measures, such as cognitive appraisal of emotions. We provide a summary of practical recommendations for validating the indicators of bottlenose dolphin welfare. This paper aims to stimulate further research into dolphin welfare which could improve the lives of the animals themselves and ultimately support regulatory decisions. We recommend uniting expertise in cetology and welfare science in order to develop a holistic approach to dolphin welfare assessment.

Keywords: affective states, animal-based measures, animal welfare, bottlenose dolphins, cetaceans, welfare assessment

Introduction

Research into welfare assessments for zoo and aquarium (hereafter referred to as ‘zoo’) animals is increasing as farm animal welfare assessment is proven to be feasible and valid (Barber 2009; Whitham & Wielebnowski 2013). There is much support for the adaptation of farm animal measures to zoo animals (Swaigood 2007; Hill & Broom 2009; Mason & Veasey 2010), and zoological institutions are well set-up for conducting measures due to the individualised care and multiple daily interactions (Barber 2009).

In this review, we conceptualise welfare using Spruijt *et al*'s (2001) description of a “...balance between positive (reward, satisfaction) and negative (stress) experiences or affective states. The balance may range from positive (good welfare) to negative (poor welfare)”. This ‘feelings-based’ definition aligns with positions taken recently by many others (eg Yeates & Main 2008; Mason & Veasey 2010; Watters 2014; Dawkins 2015), and specifies measurement of both positive and negative welfare. Our review also prioritises animal-based over resource-based measures, since they are more likely to accurately reflect welfare (Webster 2005; Roe *et al* 2011; Whitham & Wielebnowski 2013). We also give equal consideration to indicators of positive and negative welfare (Désiré *et al* 2002; Paul *et al* 2005; Boissy *et al* 2007; Yeates & Main 2008).

There is very little existing research on the welfare of cetaceans (Ugaz *et al* 2013; Clegg *et al* 2015), in captivity or the wild. Given that public interest often stimulates research in the associated areas (eg with farm animal welfare; Rushen *et al* 2011), cetacean welfare studies are likely to increase markedly in the near future since the public's interest is at a high level and intensity (Grimm 2011; Ventre & Jett 2015). Although many questions posed are, in fact, ethical dilemmas (eg ‘Should we keep dolphins in captivity?’), objective data on the animals' welfare state would aid in these personal decisions (Jiang *et al* 2007; Clegg *et al* 2015). Published farm welfare assessments have shown how this type of data can be gathered, for instance the Welfare Quality® project for farm animals (2009a,b,c) and its subsequent cross-species adaptations (eg Mononen *et al* 2012; Clegg *et al* 2015; Barnard *et al* 2016).

While cetology, the study of cetaceans, has burgeoned (Hill & Lackups 2010), there are very few studies on cetacean welfare and methods of assessment (Galhardo *et al* 1996; Clark 2013; Ugaz *et al* 2013; Clegg *et al* 2015). Bottlenose dolphins (*Tursiops truncatus*) are the most studied cetacean species (Hill & Lackups 2010), and the most common in captivity (Pryor & Norris 1998; Wells & Scott 1999), thus we choose them as the focus species for this review of how welfare science could be applied to cetaceans. We arrange the relevant cetology knowledge

into behaviour, health-related and cognition research, all well-established specialities (Wells 2009). These three categories are derived from Webster's (2005) 'Triangulation' principle for the measurement of welfare, where accumulating information from each specialty increases overall validity. Importantly, while our review includes wild research and welfare applications (wild animal welfare should be measured: Jordan 2005; Ohl & van der Staay 2012), our discussions orientate towards captive dolphins since their environment is closely controlled by humans. The necessity for dolphin welfare research is clear: there are 250 bottlenose dolphins in the European Association of Zoos and Aquaria facilities (EAZA 2015), 444 animals listed in US and Canadian dolphinariums (in 2011; Ceta-base), and many others worldwide not registered on an official record, all maintained in a huge range of facility types that differentially impact welfare (Joseph & Antrim 2010).

This review compiles what we believe to be literature on wild and captive bottlenose dolphins most relevant to welfare, suggesting some farm and zoo animal approaches which could be adapted to cetaceans, with final recommendations on initial studies and how the dolphin welfare discipline might evolve. A strong focus is maintained on those areas of cetology that merit further investigation to answer questions on bottlenose dolphins' quality of life.

Published work on dolphin welfare

There are very few studies of dolphin welfare, either in captivity or the wild (Ugaz *et al* 2013; Clegg *et al* 2015). Thus, there are no validated measures, ie ones that we know are linked to positive or negative affective states, as yet. Given the dearth of welfare research, in some cases findings from other cetacean species are extrapolated to bottlenose dolphins.

Studies of wild dolphin welfare

Only a handful of studies have focused on welfare concepts with regard to wild cetaceans, and even within these direct mentioning of the word 'welfare' is rare. A popular topic has been assessments of the impacts of tourist boats on various cetacean species (eg Stockin *et al* 2008; Christiansen & Lusseau 2015), although the focus remains at population-level indicators. Long-term data revealed that rate and repetitions of wild bottlenose dolphin whistles were potential indicators of short-term stress (Esch *et al* 2009). A more recent study suggested that an upward shift in whistle frequency was linked to increased emotional arousal (Heiler *et al* 2016). Butterworth *et al* (2013) empirically evaluated dolphin welfare in the Taiji drive hunts, an annual harvesting of dolphins in Japan, but this research only concentrated on welfare at the point of death. In the first and only teaming of wild marine mammal research with animal welfare science to our knowledge, Butterworth *et al* (2012) used the Five Freedoms to discuss how entanglement affects individual animal welfare in a number of species, including dolphins.

Studies measuring captive dolphin welfare

Similarly, there are only a handful of captive dolphin studies that have endeavoured to develop welfare measures. Ugaz *et al* (2013) correlated salivary cortisol to behavioural parameters in 23 *T. truncatus*, concluding that welfare was better in open (enclosed area of the sea) than closed (artificial water and pool) facilities due to lower cortisol levels and less floating and circular swimming. Castellote and Fossa (2006) suggested acoustic activity as a welfare measure for belugas (*Delphinapterus leucas*) and found it dropped to low levels during stressful events, but they did not correlate it with other parameters and only studied two animals. In a multi-disciplinary approach, Waples and Gales (2002) looked at inappetence, social behaviour, lethargy, weight loss and blood parameters in three *T. truncatus* with substantially deteriorating welfare likely due to social stress, revealing useful associations although again limited by sample size.

The C-Well[®] Assessment

In the first development of a welfare assessment protocol for dolphins, Clegg *et al* (2015) studied 20 *T. truncatus* in three facilities and adapted a well-established farm animal assessment (Welfare Quality[®] 2009a,b,c) to this species (the C-Well[®] Assessment). The research used 36 multi-dimensional measures, 58% of which were animal-based, to yield individual welfare scores comparable on many different levels (eg by measure, criteria, in total; among pools, sex, age class, facilities). Although the measures were unweighted, they were validated through expert opinion and application in specific contexts, and have associated standardised methods and thresholds. Some are reviewed in the relevant categories in *Research relevant to dolphin welfare*.

Given the lack of existing studies on dolphin welfare indicators, the next section is a review of cetology disciplines relevant to welfare. We expanded Webster's (2005) welfare measurement categories to behaviour, health (from Webster's 'physiology') and cognition (from 'neurobiology'). Health, while still including physiology, encompasses longer-term welfare indicators, and cognition includes experimental psychology methods potentially valuable for assessing welfare.

Research relevant to dolphin welfare

Health

Health-welfare interface

Health and welfare interact directly and indirectly as part of a complex relationship (Walker *et al* 2012). A reasonable level of health is considered a prerequisite for good animal welfare (Webster 2005; Hill & Broom 2009), while poor health is a likely contributing factor to poor welfare (Fraser *et al* 1997; Dawkins 2004; Boissy *et al* 2007; Mason & Veasey 2010). But do all components of poor health affect welfare? We refer back to our definition of welfare (Spruijt *et al* 2001) to address this: the balance of affective states and health and should only be impacted when poor health

either directly impacts affective state through, for example, nausea, lethargy or pain, or indirectly through loss of function. Poor health (eg an asymptomatic tumour) does not always affect emotional state and hence welfare, as we define it (Fraser *et al* 1997; Mason & Veasey 2010).

Health parameters in dolphins

An infection or disease can cause pain and/or 'sickness behaviour', which includes inappetence, lethargy, depression, and anti-social behaviours, all of which have direct or indirect effects on affective state (Broom 1991; Millman 2007; Sneddon *et al* 2014). Dolphins tend to mask symptoms of pain and disease as a survival adaptation (Waples & Gales 2002; Castellote & Fossa 2006), which therefore may only become obvious when the health problem is severe. Perhaps, as a consequence, little is published about indicators of pain in dolphins, with exceptions for extreme situations, such as their behavioural response to killing methods in the wild (Butterworth *et al* 2013). Weary *et al* (2006) and Sneddon *et al* (2014) provide cross-species advice for identifying behavioural and physiological pain markers, such as studying behavioural differences after analgesia administration. Inappetence and lethargy in dolphins have been correlated with many different diseases and together are generally reliable as poor health indicators (Joseph *et al* 1986; Johnson *et al* 2009). However, they can also be caused by social stress or even reproductive events such as oestrus (Waples & Gales 2002), where the associated affective states may vary from negative to positive. Studies on the behavioural and haematological characteristics of inappetence, where differentiations are made depending on whether it was caused by poor health or social stressors, are much needed.

In lieu of reliable pain indicators, physical bodily damage has been used as a health-related welfare measure in other captive species (Broom 1991; Welfare Quality® 2009a,b,c; Mononen *et al* 2012), and for wild animals as well (eg Jordan 2005; Cattet *et al* 2008). Clegg *et al* (2015) proposed the percentage of rake marks (superficial lesions and scars caused by conspecifics in play, sexual and aggressive behaviours; Scott *et al* 2005) on the body as a welfare measure for bottlenose dolphins, since such marks can be used as a proxy indicator of aggression levels and social stress (Scott *et al* 2005; Orbach *et al* 2015). However, this measure requires further investigation, for example to differentiate rake-mark levels due to high levels of play and aggression while controlling for age and sex differences.

Longer-term measures of dolphin health could also be useful for assessing welfare. Body Condition Scoring (BCS), an assessment of the extent of body fat present (Roche *et al* 2009), has been favoured as a general welfare measure (eg wild: Mann & Kemps 2003; Pettis *et al* 2004; Cattet *et al* 2008; captive: Roche *et al* 2009; Welfare Quality® 2009a,b,c; Mononen *et al* 2012) and it has already been used in wild health assessments of *T. truncatus* (Fair *et al* 2014; Schwacke *et al* 2014). Joblon *et al* (2014) produced a standardised protocol using stranded short-beaked common dolphins (*Delphinus delphis*), and Clegg *et al* (2015)

developed a standardised BCS graphic for *T. truncatus* but did not test its reliability. The next step for these BCS tools is to correlate the results to other measures of affective state: Roche *et al* (2009) conducted this with cows, concluding that BCS may serve as a proxy indicator for hunger, satiety or feeling ill (leading to inappetence). Other health-related conditions such as diarrhoea, skin inflammation, eye condition, and coughing, have been used as farm animal welfare measures (Welfare Quality® 2009a,b,c; Mononen *et al* 2012), some of which were proposed for *T. truncatus* welfare (skin and eye condition, coughing; Clegg *et al* 2015), but have not been studied in relation to affective states. Haematological indices can be measures of disease states, especially when the pathology is advanced, but so far have had limited use in welfare assessments due to potential high inter- and intra-individual variation. Although wild dolphin health assessments have published their data and established baselines (Thomson & Geraci 1986; Dierauf & Gulland 2001; Wells 2009), and captive dolphin voluntary blood sampling is readily achievable using positive reinforcement training (Brando 2010), studies have not yet linked ranges of blood values to health-related welfare.

Population measures of longer-term health and welfare such as longevity and reproductive rate should also be considered (Dawkins 1998; Barber 2009). However, as with farm animals, parameters, such as high reproductive success, do not necessarily indicate that welfare is good (Dawkins 1980). For captive dolphin populations, baselines are being established by projects (notably in the US) allowing access to their valuable multi-species databases (Small & DeMaster 1995; Innes *et al* 2005; Venn-Watson *et al* 2011). Welfare conclusions from fitness measures should be supported by other data (Swaigood 2007), such as in Christiansen and Lusseau's study (2015) linking disturbance behaviour from whale-watching boats, body condition and foetal growth rate in minke whales (*Balaenoptera acutorostrata*). Data on the incidence and severity of diseases can also be used as population-level health parameters: such wild studies are available (eg Reif *et al* 2008; Schwacke *et al* 2014), but data are not published for captive dolphins. However, extensive records are kept for most captive dolphins (C van Elk, personal communication 2016) and, thus, peer-reviewed publications on the nature of their diseases would be beneficial for establishing standardised health assessments.

Physiological parameters

Measures of physiological responses can contribute to assessments of emotions and affective states (Désiré *et al* 2002; Webster 2005; Boissy *et al* 2007). Endocrine responses to stressors are most commonly used (eg Moberg & Mench 2000), but as interest increases in positive welfare, other markers are being considered: for example, the balance of sympathetic and parasympathetic systems (for a review, see Boissy *et al* 2007), and indicators of eustress (positive stressors, eg mate acquisition, experienced by the animal; Selye 1975). Within dolphin physiology research, numerous studies of physiological measures

of stress for wild *T. truncatus* (eg Ortiz & Worthy 2000; Fair *et al* 2014; Schwacke *et al* 2014) have provided useful baselines, which will start to elucidate individual variation and repeatability questions (Atkinson *et al* 2015). Sample collection in the wild is challenging since taking blood is not possible without restraint and faecal samples are difficult to obtain (Atkinson *et al* 2015). This area, specifically, is where training for voluntary samples in captivity has exceptional advantages; for example, voluntary saliva collection is feasible and can provide accurate cortisol measurements in *T. truncatus* (Pedernera-Romano *et al* 2006; Ugaz *et al* 2013). Other sampling protocols are also possible with training, such as blood, faecal, blow (expiration of air) and biopsy collection. As for terrestrial animals, marine mammals experience diurnal and seasonal variation in cortisol levels (for a review, see Atkinson *et al* 2015), which would need to be taken into account in any welfare assessment and suggests that a conservative range would need to be used in any conclusions made, as opposed to a single threshold. Additionally, a recent review advised caution when applying terrestrial animal stress models to marine mammals. While corticosteroid pathways seem to be similar, evidence indicates other neuroendocrine hormones (eg catecholamines) may be regulated very differently (Atkinson *et al* 2015). Further, long-term studies on cetaceans in captivity could start to answer such questions on hormone regulatory systems. Innovative new collection techniques enabling accurate animal identification (example with cetaceans: whale blow), and insightful behavioural correlations, must also guide future progress (Möstl & Palme 2002): advice very applicable to dolphin studies.

Behaviour

Social behaviour

Behavioural measures are an important component in welfare frameworks (Dawkins 2004; Maple 2007), with some believing that they are more informative about welfare than health since behaviours are likely more indicative of emotional state (Gonyou 1994; Joseph & Antrim 2010). There have been a number of long-term studies of wild dolphin behaviour (Wells 1991; Mann *et al* 2000; Parsons *et al* 2006), including social relationships within their fission-fusion societies (Mann *et al* 2000; Wells 2009). Surprisingly, ethological studies of captive populations have not, until recently, been commonplace (Dudzinski 2010). Social behaviour measures will foreseeably be one of the most important tools in assessing dolphin welfare: as highly social mammals (Pryor & Norris 1998; Mann *et al* 2000), they are susceptible to social stress. Sudden changes in conspecific associations, aggression levels and social isolation have been correlated with declines in welfare (Waples & Gales 2002). Excessive or abnormal aggression levels are used as farm animal welfare measures (Webster 2005; Welfare Quality® 2009b; Mononen *et al* 2012), and using existing ethograms of aggressive behaviours to analyse frequencies over time could reveal dolphins' 'excessive' and 'normal' thresholds (Samuels & Gifford 1997; Scott *et al* 2005). Increased quantity and severity of

rake marks could serve as a proxy indicator for levels of aggression and social stress (Waples & Gales 2002; Scott *et al* 2005). Clegg *et al*'s (2015) rake-mark assessment, currently using very conservative thresholds, might be validated to allow monitoring of aggression levels. Rake-mark quantification is an example of a method where collaboration between wild and captive researchers might be fruitful, since aggression is harder to observe in the wild and rake-mark levels could be used as proxy measures (Clegg *et al* 2015; Scott *et al* 2005).

While social stress is a negative consequence of being a social mammal, the highly social life of dolphins also has positive effects. Positive social (ie affiliative) behaviour has been proposed as an indicator of good welfare in other species (Boissy *et al* 2007; Buchanan-Smith *et al* 2013). Affiliative behaviour has been well-documented in wild *Tursiops* spp (Herman & Tavolga 1980 (early review); Connor *et al* 2000, 2006; Sakai *et al* 2006), and a little less so in captivity (Tamaki *et al* 2006; Dudzinski 2010). Gentle rubbing behaviours between dolphins are thought to be analogous to allogrooming in terrestrial mammals (Tamaki *et al* 2006; Kuczaj *et al* 2013) and may have potential as a measure of good welfare (Boissy *et al* 2007), along with synchronous swimming (thought to reflect social bonds; Connor *et al* 2006). Dudzinski (2010) and Kuczaj *et al* (2013) reviewed both wild and captive social affiliative behaviour, agreeing that in both settings tactile behaviours seem to be used to express emotions.

Play

Play behaviour is one of the strongest potential welfare indicators for animals, mainly because it is more likely to occur in the absence of threats and utilitarian needs (Bel'kovich *et al* 1991) and is linked to positive emotions (Held & Špinka [2011] reviewed link with welfare). Play is also likely to improve long-term fitness and health, as well as being socially contagious and therefore capable of spreading good welfare in groups (Held & Špinka 2011); these less-acknowledged benefits are especially relevant to the welfare of group-living dolphins. Despite this, play is not commonly used as a measure in welfare assessments (Welfare Quality® 2009a,b,c), most likely because of its inherent variability (Held & Špinka 2011) and difficulty of measurement. There is also evidence for some species that play may not always be linked to a positive emotional state (Blois-Heulin *et al* 2015).

Evidence for wild and captive dolphin play is abundant (for reviews, see Paulos *et al* 2010; Kuczaj & Eskelinen 2014), including copious examples of object play (recent papers: Kuczaj & Makecha 2008; Paulos *et al* 2010; Greene *et al* 2011; Delfour & Beyer 2012), and evidence of inventing games (Pace 2000). McCowan *et al* (2000) showed that captive dolphins monitored their bubble quality as well as 'planned' for the behaviour; suggesting involvement of conscious thought and appraisal and strengthening the notion that play impacts affective state. Hill and Ramirez (2014) studied play in 14 belugas (*Delphinapterus leucas*) over three years, revealing differences between adult and

young preferences and showing that bouts were longer and more diverse when enrichment devices were involved. Where play is studied in captivity, the influence of any prior conditioning should be noted: Neto *et al* (2016) showed that when trainers positively reinforced dolphins' interactions with toys, their interest in the objects increased outside of the sessions. This technique could be used to increase the benefits of the toys to the dolphins but, until we have other measures of positive affective states available in this situation, the motivation to play may be influenced. In any case, as with all species, standardised measurements are needed. A study of African elephants (*Loxodonta africana*) addressed this using a play index (Vicino & Marcacci 2015), and a similar approach might be possible with dolphins. Such a behavioural measure could easily be applicable to wild dolphin welfare assessments, for example to investigate whether exposure to more environmental or social stressors results in reduced play frequencies.

Abnormal behaviour

Abnormal behaviours are most often studied in the context of stereotypic behaviour, which has been most recently defined as "...repetitive behaviour induced by frustration, repeated attempts to cope, and/or CNS dysfunction" (Mason & Rushen 2008). Abnormal behaviours are seldom observed in wild animals, although Miller *et al* (2011a) suggest they observed stereotypic swimming in lemon sharks (*Negaprion brevirostris*). In one of the only studies describing abnormal behaviour in wild dolphins, the causes and effects of solitary living for *T. truncatus* were investigated, and certain aspects were concluded as abnormal (eg behaviour oriented excessively towards humans; Müller & Bossley 2002). Stereotypic behaviour is commonly investigated as a welfare measure for captive animals (for a review, see Mason & Rushen 2008). There are scarcely any published studies with captive dolphins, and the small handful existing are outdated (Gygax 1993 and Clark 2013 describe this literature), making it hard to identify any common explanatory variables. Stereotypic swimming has been discussed in the literature as a concern for captive dolphins. There are disparities among definitions of this behaviour (Gygax 1993; Miller *et al* 2011b), but terrestrial animal studies also suffered similar problems with pacing behaviours and found that the variability and possible functions must be meticulously analysed for it to be defined as a stereotypy (Mason & Rushen 2008). Sobel *et al* (1994) observed preferences in the directionality of circular swimming, but did not measure whether the route around the pool was fixed and whether the animal was vigilant at the time. There is little evidence correlating stereotypic swimming with other potential factors of welfare. For example, although Ugaz *et al* (2013) found that in closed facilities there were higher rates of circular swimming as well as higher cortisol levels, they did not statistically test for a correlation between these two factors. Clegg *et al* (2015) included a stereotypy measure but based thresholds on terrestrial animal frequencies, since there was no data from dolphin species. Since there are ongoing questions

about whether higher stereotypy frequency infers poorer welfare, even in terrestrial animals (Mason & Latham 2004; Dawkins 2006; Mason & Rushen 2008), future studies on this phenomenon in dolphins should aim to correlate suspected stereotypic behaviour with other indicators of welfare to validate it as an indicator. Basic work regarding the appearance of stereotypies is also needed, for example, the two main types of stereotypy defined in terrestrial mammals are 'oral' and 'movement' (Webster 2005; Mason & Rushen 2008), so a fundamental investigation would be to ascertain whether the same is true for dolphins.

Anticipatory behaviour

Recently, anticipatory behaviour (a measure of 'reward-sensitivity'; Spruijt *et al* 2001) has been used as a welfare measure in farm and zoo animals; low intensity anticipatory behaviour is thought to reflect positive affective states and high intensity anticipation indicates poorer welfare as the animal is heavily dependent on the reward (Spruijt *et al* 2001; Watters 2014). While one preliminary study focused on anticipatory behaviour in captive bottlenose dolphins (Jensen *et al* 2013), further work is necessary given the different training methods (ie conditioning to 'rewards') in dolphin facilities, which might ultimately be closely linked to welfare. Based on the reward-sensitivity paradigm it is likely that dolphins showing moderate anticipatory behaviour might experience positive affective states, while those that perform it for excessive amounts of time might be frustrated or have little other stimulation (Watters 2014). In order to evaluate its utility for welfare assessment, future studies with dolphins and other species should investigate the association between anticipatory behaviour frequencies and other welfare indicators, in order to understand what might qualify as 'excessive' levels.

Cognition

Emotions

Emotions are defined as intense, short-lived affective responses to an event, usually associated with specific physiological changes (Dantzer 1988). Animals with higher cognitive abilities may be capable of more complex emotions (eg guilt) (Paul *et al* 2005), and while this might result in increased chances of suffering, it could also lead to higher potential for positive affective states. Research beyond the 'basic emotions' (eg fear, rage, play; Mendl *et al* 2010) is essential to understand the valence and arousal levels of affective states which make up overall welfare (Leliveld *et al* 2013; Siegford 2013). Désiré *et al* (2002) and Boissy *et al* (2007) provide reviews on measuring animal emotion and the relevance to welfare.

Although dolphin emotion studies are scarce, there have been more studies on negative than positive ones. Most studies have looked at how sounds might reflect emotions, for example, burst pulse sounds have been associated with agonistic and aggressive behaviours (Overstrom 1983), and long-term etho-acoustical projects have made headway in pairing sounds to behaviour (eg Herzing 1996; Janik & Sayigh

2013). Animal emotion research is now widening to measure positive emotions as well (Boissy *et al* 2007), but there are no strongly supported indicators as yet in dolphins (Kuczaj *et al* 2013). Tactile behaviour was suggested by Dudzinski (2010) and Kuczaj *et al* (2013) to be linked to positive emotions in dolphins, but has not yet been analysed in conjunction with other indicators. Motivation and preference testing are applicable to captive dolphins and could reveal indicators of emotion (Gonyou 1994; Mendl *et al* 2010).

Environmental enrichment

This sub-section is applicable to dolphins under human care only. Environmental enrichment is any technique designed to improve biological functioning of captive animals through environmental modification (Newberry 1995). Bottlenose dolphins are good candidates for enhanced welfare through enrichment due to their cognitive abilities (Schusterman *et al* 2013) and propensity to, and creativity within, play (Kuczaj & Eskelinen 2014). Enrichment has been provided to captive dolphins for several decades, but there are few published studies describing the animals' responses (for a review, see Clark 2013). Furthermore, enrichment is often assumed to automatically enhance welfare even if it is unclear whether the animal's affective state will be improved (for reviews, see Swaisgood 2007; Würbel & Garner 2007). Enrichment strategies should be monitored to ensure that they are true enrichment by monitoring the animals' responses and looking for signs of habituation so that decisions can be made as to when, where and how the enrichment is presented again (Kuczaj *et al* 2002; Hoy *et al* 2010; Siegford 2013).

Recently, cognitive challenges have been presented as enrichment, with positive results as long as the animals possess the resources and skills to solve the problem (Meehan & Mench 2007; Siegford 2013). Clark (2013) supports cognitive enrichment with dolphin species, hypothesising that floating, simplistic objects are not sufficient to hold the dolphins' interest in the long-term. However, behaviour should be monitored to investigate whether this is indeed the case (Hill & Broom 2009), and such data, which shows responses to definable, repeatable contexts, could also aid in finding welfare indicators (Delfour & Beyer 2012). The Human-Animal Relationship (HAR) is only just beginning to be investigated in other species in relation to cognitive enrichment and welfare (Whitham & Wielebnowski 2013) and, due to the multiple, daily, and often close-contact interactions, is very likely to contribute to the welfare state of captive dolphins (Brando 2010; Clegg *et al* 2015). Future investigations assessing the HAR might aim to disentangle the effects of food reinforcement with the dolphins' attitude towards the humans themselves. An example of such an approach is shown by Perelberg and Schuster (2009), who demonstrated that outside of feeding sessions, a captive bottlenose dolphin group approached humans to receive rubs and petting in the absence of any other rewards. Given that many cetacean species show much tactile behaviour during intra-specific social affiliation (Dudzinski 2010; Kuczaj

et al 2013), investigating the frequency and dimensions of voluntary human contact by the animals, during and outside of training sessions, might be a meaningful measure of their affective states.

Cognitive measures of dolphin welfare

Dolphins' cognitive abilities are frequently compared to those of great apes (Delfour 2010; Schusterman *et al* 2013), and may be linked to their fusion-fission society (Connor *et al* 2000; Maze-Foley & Würsig 2002). Dolphins display co-operative hunting (Connor *et al* 2000), use tools (eg Smolker *et al* 1997), and recognise their mirror image (Reiss & Marino 2001; Delfour 2006). Studies of cognitive bias, which investigate how emotional experiences affect cognitive processes, may aid in our interpretation of welfare, and constitute measures themselves (Paul *et al* 2005; Mendl *et al* 2009). Given the dolphins' learning capabilities (Brando 2010), many of the non-invasive cognitive bias methods reviewed in Mendl *et al* (2009) used with other species could be adapted. Paul *et al* (2005) also reviewed evidence for memory and attention bias processes in animals, concluding that if confirmed they could have implications for measuring welfare.

In humans as well as non-humans, the brain hemispheres process information differently, producing lateralised behaviours, ie a preference for either the left or right eye or body part (Rogers 2002). It seems that animals may predominantly use the right hemisphere when stressed (see Rogers 2010), with Leliveld *et al*'s (2013) review going further to conclude that negative emotions are managed by the right hemisphere and positive emotions by the left ('emotional lateralisation'). Examples of lateralised behaviours in wild and captive cetaceans are common, eg during foraging (Clapham *et al* 1995; Silber & Fertl 1995), flipper-rubbing (Sakai *et al* 2006), and visual discrimination tasks (Yaman *et al* 2003; Delfour & Marten 2006). Most notably, Karenina *et al* (2010, 2013) showed that belugas and killer whales (*Orcinus orca*) placed calves on their right side in non-threatening situations, with killer whales preferring the left when the situation became increasingly threatening (in this case proximity to boats). Sakai *et al* (2006) suggested a link with positive affective state since the left pectoral fin and eye were favoured during affiliative flipper-rubbing behaviour in Indian Ocean bottlenose dolphins (*Tursiops aduncus*). These last examples concerning lateralised behaviour and affective states should form a basis for future research into welfare implications of this phenomenon.

In the field of cognition, in particular, but also within health and behaviour, researchers rarely study both wild and captive dolphins, resulting in a skewed perspective of particular topics in certain environments, and leading us to an initial recommendation to increase collaborative efforts and reviews (in accordance with Hill & Lackups 2010; Pack 2010). Finally, although we must understand the dolphins' cognitive abilities, we should do so bearing in mind their *umwelt*, ie their 'subjective universe', and the focus of etho-phenomenological studies (Delfour 2006, 2010). For example, an intermodal associative task was completed

very differently by bottlenose dolphin subjects due to the dominance hierarchy at the time dictating participation and mode of learning (Delfour & Marten 2006). Being cognisant of the dolphins' *umwelt* may help in determining what is important to the dolphins, and thus how to provide them with a good quality of life.

Considerations for developing dolphin welfare measures

In this section we review recommendations on the design of studies to investigate the measures discussed above. Welfare measures should be developed *in situ*, thus ensuring applicability to the dolphins and their environment (Dawkins 2006; Maple 2007). The measures must also be species-specific (Blokhuis 2008; Barber 2009; Hill & Broom 2009), and examine welfare at an individual level where possible (Siegford 2013). Zoological institutions have been advised to employ scientists specifically dedicated to assessing welfare (Maple 2007; Barber 2009), and facilities maintaining dolphins should take this step also.

The first proposed measures for *T. truncatus* should be validated through correlations with other parameters. Our review addresses the potential measures for validation which we identified with respect to cetacean health, behaviour and cognition (Table 1). When establishing welfare measures, studies of captive rather than wild dolphins, are more likely to be successful due to greater access to the subjects, their history, and their environment. International, inter-facility collaborations are vital to combat problems of low sample sizes and to control for inevitable environmental variation. For wild dolphin welfare indicators, long-term studies are the natural starting point since most have individual behavioural, physiological, as well as life history data (Wells 2009; Fair *et al* 2014). While it would be inaccurate to apply all measures for wild and captive animals without validation (Jordan 2005), it is likely that many welfare indicators, at least behaviourally, will be consistent between wild and captive *T. truncatus* since their repertoires show similarities (Mann *et al* 2000; Dudzinski 2010).

When validating measures, pre-existing conditions can be used where it is likely the animal has very good or poor welfare (Jordan 2005; Castellote & Fossa 2006; Whitham & Wielebnowski 2013). For example, transportation offers opportunities to assess welfare as it is assumed to induce a substantial, but short-term, welfare change for captive cetaceans (eg Castellote & Fossa 2006). Long-term states associated with social contexts may be more salient for welfare measurement: for example, the period after transport when the animals are introduced to a new group. Group changes are frequent enough in dolphinarium networks to provide adequate sample sizes for analysis. The selected behaviours and physiological parameters should then be measured during these events (and cognitive data if possible), with focal qualitative data (eg trainer ratings) taken concurrently to support the presumed change in welfare. Welfare measures should be conducted regularly, and also separately from full assessments. For example,

Table 1 Summary of the welfare-related topics in dolphin health, behaviour and cognition which merit further investigation in order to develop measures of welfare. Evidence supporting each topic has been taken from bottlenose dolphins (*Tursiops truncatus*) where possible, but where these were lacking studies from other cetacean species had to be used.

Category	Aspects meriting further investigation as dolphin welfare measures
Health	Epidemiological measures (eg mortality, reproductive success)
	Disease prevalence
	Body Condition Scoring
	Cortisol (and other 'stress' hormone) levels
Behaviour	Rake mark percentage cover
	Excessive aggression
	Affiliative behaviour
	Play
Cognition	Anticipatory behaviour
	Abnormal and stereotypic behaviours
	Emotions linked with sound production
	Indicators of basic emotions (eg fearful, playful, rage)
	Indicators of more complex emotions (eg contentedness, depression)
	Cognitive bias testing
Visual and behavioural laterality	

behavioural measures of welfare could be applied on a weekly basis to dolphin groups since behavioural monitoring has been advised as essential for ensuring good welfare (Maple 2007), and especially with captive dolphins (Waples & Gales 2002; Clegg *et al* 2015). Eventually, comparing results from measures and assessments between individuals can highlight associations with good or poor welfare, thus indicating where changes in management protocols should occur and stimulating improvements in welfare of the animals themselves.

Conclusion

We have reviewed the literature on animal welfare science and cetology in order to identify the most successful intersections for developing bottlenose dolphin welfare measures. A general theme is that collaborations, whether wild-captive, across different cetology fields, or between multiple captive facilities, are necessary if we want to address this multi-dimensional concept.

We suggest that indicators, such as cortisol levels, inappetence and bodily injuries, as well as body condition and population fitness measures in the longer term, may help us assess health-related welfare. Behavioural measures are

likely to be the most informative for dolphin welfare, and we have shown evidence that tactile affiliation, play, anticipatory behaviour and stereotypic behaviours may be closely linked to affective states. Cognitive measures reflect how behavioural and physiological components are integrated to form the affective states experienced by the animals, and thus recent techniques, such as cognitive bias testing hold much promise for welfare assessment.

Lastly, we identified practical recommendations for validating the first measures, concluding that although captive studies should take the lead, long-term wild studies are also rich sources of potential indicators. Any proposed measures should be tested in situations likely to elicit changes in welfare with adequate sample sizes to allow the major environmental variations to be controlled for. Established measures would allow facility managers to monitor and improve the dolphins' welfare, aid in regulatory decisions, and could enrich wild dolphin research by revealing changing affective states. This review's findings are species-specific to bottlenose dolphins, but the general principles and selected measures could be adapted to other cetacean species. Our overall aim was to present current cetology knowledge in terms of measuring welfare, with the hope of stimulating researchers globally to take up the challenge.

Acknowledgements

The authors are each indebted to a number of experts in the various fields of cetology and welfare science for their discussions and encouragement. This work forms part of IC's PhD, supported by CIFRE fund no 2014/0289, and UFAW grant no 22-14/15.

References

- Atkinson S, Crocker D, Houser D and Mashburn K** 2015 Stress physiology in marine mammals: how well do they fit the terrestrial model? *Journal of Comparative Physiology B*: 1-24. <https://doi.org/10.1007/s00360-015-0901-0>
- Barber JC** 2009 Programmatic approaches to assessing and improving animal welfare in zoos and aquariums. *Zoo Biology* 28: 519-530. <https://doi.org/10.1002/zoo.20260>
- Barnard S, Pedernera C, Candeloro L, Ferri N, Velarde A and Dalla Villa P** 2016 Development of a new welfare assessment protocol for practical application in long-term dog shelters. *The Veterinary Record* 178: 18
- Bel'kovich VM, Ivanova EE, Kozarovitsky LB, Novikova EV and Kharitonov SP** 1991 Dolphin play behavior in the open sea. In: Pryor K and Norris KS (eds) *Dolphin Societies: Discoveries and Puzzles* pp 67-77. University of California Press: London, UK
- Blois-Heulin C, Rochais C, Camus S, Fureix C, Lemasson A, Lunel C, Bézard E and Hausberger M** 2015 Animal welfare: Could adult play be a false friend? *Animal Behavior and Cognition* 2: 156-185
- Blokhuis HJ** 2008 International cooperation in animal welfare: the Welfare Quality[®] project. *Acta Veterinaria Scandinavica* 50: S10. <https://doi.org/10.1186/1751-0147-50-S1-S10>
- Boissy A, Manteuffel G, Jensen MB, Moe RO, Spruijt B, Keeling LJ and Aubert A** 2007 Assessment of positive emotions in animals to improve their welfare. *Physiology & Behavior* 92: 375-397. <https://doi.org/10.1016/j.physbeh.2007.02.003>
- Brando SICA** 2010 Advances in husbandry training in marine mammal care programs. *International Journal of Comparative Psychology* 23: 777-791
- Broom DM** 1991 Assessing welfare and suffering. *Behavioural Processes* 25: 117-123. [https://doi.org/10.1016/0376-6357\(91\)90014-Q](https://doi.org/10.1016/0376-6357(91)90014-Q)
- Broom DM** 2010 Animal welfare: an aspect of care, sustainability, and food quality required by the public. *Journal of Veterinary Medical Education* 37: 83-88. <https://doi.org/10.3138/jvme.37.1.83>
- Broom DM and Johnson KG** 1993 *Stress and Animal Welfare*. Chapman & Hall: London, UK. <https://doi.org/10.1007/978-94-024-0980-2>
- Buchanan-Smith HM, Gričiute J, Daoudi S, Leonardi R and Whiten A** 2013 Interspecific interactions and welfare implications in mixed species communities of capuchin (*Sapajus apella*) and squirrel monkeys (*Saimiri sciureus*) over 3 years. *Applied Animal Behaviour Science* 147: 324-333. <https://doi.org/10.1016/j.applanim.2013.04.004>
- Butterworth A, Brakes P, Vail CS and Reiss D** 2013 A veterinary and behavioral analysis of dolphin killing methods currently used in the 'drive hunt' in Taiji, Japan. *Journal of Applied Animal Welfare Science* 16: 184-204. <https://doi.org/10.1080/1088705.2013.768925>
- Butterworth A, Clegg I and Bass C** 2012 *Untangled-marine debris: a global picture of the impact on animal welfare and of animal-focused solutions*. World Society for the Protection of Animals International: London, UK
- Castellote M and Fossa F** 2006 Measuring acoustic activity as a method to evaluate welfare in captive beluga whales (*Delphinapterus leucas*). *Aquatic Mammals* 32: 325. <https://doi.org/10.1578/AM.32.3.2006.325>
- Cattet M, Boulanger J, Stenhouse G, Powell RA and Reynolds-Hogland MJ** 2008 An evaluation of long-term capture effects in ursids: implications for wildlife welfare and research. *Journal of Mammalogy* 89(4): 973-990. <https://doi.org/10.1644/08-MAMM-A-095.1>
- Ceta-base** 2011 *Captive Cetaceans, Living Population*. Ceta-Base: United States & Canada. <http://www.ceta-base.com/publications/cetaceans-captive-us-canada-2011.pdf>
- Christiansen F and Lusseau D** 2015 Linking behavior to vital rates to measure the effects of non-lethal disturbance on wildlife. *Conservation Letters* 8: 424-431. <https://doi.org/10.1111/conl.12166>
- Clapham PJ, Leimkuhler E and Gray BK** 1995 Do humpback whales exhibit lateralized behaviour? *Animal Behaviour* 50: 73-82. <https://doi.org/10.1006/anbe.1995.0222>
- Clark FE** 2013 Marine mammal cognition and captive care: A proposal for cognitive enrichment in zoos and aquariums. *Journal of Zoo and Aquarium Research* 1: 1-6
- Clegg ILK, Borger-Turner JL and Eskelinen HC** 2015 C-Well: The development of a welfare assessment index for captive bottlenose dolphins (*Tursiops truncatus*). *Animal Welfare* 24: 267-282. <https://doi.org/10.7120/09627286.24.3.267>
- Connor RC, Smolker R and Bejder L** 2006 Synchrony, social behaviour and alliance affiliation in Indian Ocean bottlenose dolphins, *Tursiops aduncus*. *Animal Behaviour* 72: 1371-1378. <https://doi.org/10.1016/j.anbehav.2006.03.014>

- Connor RC, Wells R, Mann J and Read A** 2000 The bottlenose dolphin: social relationships in a fission-fusion society. In: Mann J, Connor R, Tyack P and Whitehead H (eds) *Cetacean Societies: Field Studies of Whales and Dolphins* pp 91-126. University of Chicago Press: Chicago, IL, USA
- Dantzer R** 1988 *Les Emotions*. Presses Universitaires de France: Paris, France
- Dawkins MS** 1980 *Animal Suffering: The Science of Animal Welfare*. Chapman and Hall: London, UK. <https://doi.org/10.1007/978-94-009-5905-7>
- Dawkins MS** 1998 Evolution and animal welfare. *Quarterly Review of Biology* 305-328. <https://doi.org/10.1086/420307>
- Dawkins MS** 2004 Using behaviour to assess animal welfare. *Animal Welfare* 13: 3-7
- Dawkins MS** 2006 A user's guide to animal welfare science. *Trends in Ecology and Evolution* 21: 77-82. <https://doi.org/10.1016/j.tree.2005.10.017>
- Dawkins M** 2015 Chapter two: animal welfare and the paradox of animal consciousness. *Advances in the Study of Behavior* 47: 5-38. <https://doi.org/10.1016/bs.asb.2014.11.001>
- Delfour F** 2006 Marine mammals in front of the mirror, or from body experiences to self-recognition. Cognitive ethological methodology combined with a phenomenological questioning. *Journal of Aquatic Mammals* 32: 517-527. <https://doi.org/10.1578/AM.32.4.2006.517>
- Delfour F** 2010 Marine mammals enact individual worlds. *International Journal of Comparative Psychology* 23: 792-810
- Delfour F and Beyer H** 2012 Assessing the effectiveness of environmental enrichment in bottlenose dolphins (*Tursiops truncatus*). *Zoo Biology* 31: 137-150. <https://doi.org/10.1002/zoo.20383>
- Delfour F and Marten K** 2006 Lateralized visual behavior in bottlenose dolphins (*Tursiops truncatus*) performing audio-visual tasks: The right visual field advantage. *Behavioural Processes* 71: 41-50. <https://doi.org/10.1016/j.beproc.2005.09.005>
- Désiré L, Boissy A and Veissier I** 2002 Emotions in farm animals: a new approach to animal welfare in applied ethology. *Behavioural Processes* 60: 165-180. [https://doi.org/10.1016/S0376-6357\(02\)00081-5](https://doi.org/10.1016/S0376-6357(02)00081-5)
- Dierauf L and Gulland FM** 2001 *CRC Handbook of Marine Mammal Medicine, Second Edition*. CRC Press: Boca Raton, FL, USA. <https://doi.org/10.1201/9781420041637>
- Dudzinski KM** 2010 Overlap between information gained from complementary and comparative studies of captive and wild dolphins. *International Journal of Comparative Psychology* 23: 566-586
- EAZA** 2015 *European Association of Zoos and Aquariums (EAZA) European StudBook for bottlenose dolphins*. Information provided by Cornelis van Elk, current European StudBook keeper for bottlenose dolphins, December 2015
- Esch HC, Sayigh LS, Blum JE and Wells RS** 2009 Whistles as potential indicators of stress in bottlenose dolphins (*Tursiops truncatus*). *Journal of Mammalogy* 90: 638-650. <https://doi.org/10.1644/08-MAMM-A-069R.1>
- Fair PA, Schaefer AM, Romano TA, Bossart GD, Lamb SV and Reif JS** 2014 Stress response of wild bottlenose dolphins (*Tursiops truncatus*) during capture-release health assessment studies. *General and Comparative Endocrinology* 206: 203-212. <https://doi.org/10.1016/j.ygcen.2014.07.002>
- Fraser D, Weary DM, Pajor EA and Milligan BN** 1997 A scientific conception of animal welfare that reflects ethical concerns. *Animal Welfare* 6: 187-205
- Galhardo L, Appleby MC, Waran NK and Dos Santos ME** 1996 Spontaneous activities of captive performing bottlenose dolphins (*Tursiops truncatus*). *Animal Welfare* 5: 373-389
- Gonyou HW** 1994 Why the study of animal behavior is associated with the animal welfare issue. *Journal of Animal Science* 72: 2171-2177
- Greene WE, Melillo-Sweeting K and Dudzinski KM** 2011 Comparing object play in captive and wild dolphins. *International Journal Comparative Psychology* 24: 292-306
- Grimm D** 2011 Are dolphins too smart for captivity? *Science* 332: 526-529. <https://doi.org/10.1126/science.332.6029.526>
- Gygax L** 1993 Spatial movement patterns and behaviour of two captive bottlenose dolphins (*Tursiops truncatus*): absence of stereotyped behaviour or lack of definition? *Applied Animal Behaviour Science* 38: 337-344. [https://doi.org/10.1016/0168-1591\(93\)90031-J](https://doi.org/10.1016/0168-1591(93)90031-J)
- Heiler J, Elwen SH, Kriesell HJ and Gridley T** 2016 Changes in bottlenose dolphin whistle parameters related to vessel presence, surface behaviour and group composition. *Animal Behaviour* 117: 167-177. <https://doi.org/10.1016/j.anbehav.2016.04.014>
- Held SD and Špinka M** 2011 Animal play and animal welfare. *Animal Behaviour* 81: 891-899. <https://doi.org/10.1016/j.anbehav.2011.01.007>
- Herman LM and Tavalga WN** 1980 The communication systems of cetaceans. In: Herman L (ed) *Cetacean Behavior: Mechanisms and Functions* pp 149-209. John Wiley and Sons: New York, NY, USA
- Herzing DL** 1996 Vocalizations and associated underwater behavior of free-ranging Atlantic spotted dolphins, *Stenella frontalis* and bottlenose dolphins, *Tursiops truncatus*. *Aquatic Mammals* 22: 61-80
- Hill H and Lackups M** 2010 Journal publication trends regarding cetaceans found in both wild and captive environments: what do we study and where do we publish? *International Journal of Comparative Psychology* 23: 414-534
- Hill H and Ramirez D** 2014 Adults play but not like their young: the frequency and types of play by belugas (*Delphinapterus leucas*) in human care. *Animal Behavior and Cognition* 1: 166-185. <https://doi.org/10.12966/abc.05.07.2014>
- Hill SP and Broom DM** 2009 Measuring zoo animal welfare: theory and practice. *Zoo Biology* 28: 531-544. <https://doi.org/10.1002/zoo.20276>
- Hoy JM, Murray PJ and Tribe A** 2010 Thirty years later: enrichment practices for captive mammals. *Zoo Biology* 29: 303-316. <https://doi.org/10.1002/zoo.20254>
- Innes WS, DeMaster DP, Rodriguez A and Crowder LB** 2005 Survival rates of marine mammals in captivity: Temporal trends and institutional analysis. *Sixteenth Biennial Conference on the Biology of Marine Mammals* pp 136. 12-16 December 2005, San Diego, CA, USA
- Janik VM and Sayigh LS** 2013 Communication in bottlenose dolphins: 50 years of signature whistle research. *Journal of Comparative Physiology A* 199: 479-489. <https://doi.org/10.1007/s00359-013-0817-7>
- Jensen A-L, Delfour F and Carter T** 2013 Anticipatory behaviour in captive bottlenose dolphins (*Tursiops truncatus*): A preliminary study. *Zoo Biology* 32: 436-444. <https://doi.org/10.1002/zoo.21077>

- Jiang Y, Luck M and Parsons ECM** 2007 Public awareness, education, and marine mammals in captivity. *Tourism Review International* 11: 237-249. <https://doi.org/10.3727/154427207783948829>
- Joblon MJ, Pokras MA, Morse B, Harry CT, Rose KS, Sharp SM, Niemeyer ME, Patchett KM, Sharp WB and Moore MJ** 2014 Body condition scoring system for Delphinids based on short-beaked common dolphins (*Delphinus delphis*). *Journal of Marine Animals and Their Ecology* 7: 5-13
- Johnson SP, Venn-Watson SK, Cassle SE, Smith CR, Jensen ED and Ridgway SH** 2009 Use of phlebotomy treatment in Atlantic bottlenose dolphins with iron overload. *Journal of the American Veterinary Medical Association* 235: 194-200. <https://doi.org/10.2460/javma.235.2.194>
- Jordan B** 2005 Science-based assessment of animal welfare: wild and captive animals. *Revue Scientifique et Technique (International Office of Epizootics)* 24: 515-528. <https://doi.org/10.20506/rst.24.2.1588>
- Joseph B and Antrim J** 2010 Special considerations for the maintenance of marine mammals in captivity. In: Kleiman DG, Thompson KV and Kirk Baer C (eds) *Wild Mammals in Captivity: Principles and Techniques for Zoo Management, 2nd Edition* pp 181-216. The University of Chicago Press: Chicago, IL, USA
- Joseph BE, Cornell LH, Simpson JG, Migaki G and Griner L** 1986 Pulmonary aspergillosis in three species of dolphin. *Zoo Biology* 5: 301-308. <https://doi.org/10.1002/zoo.1430050308>
- Karenina K, Giljov A, Baranov V, Osipova L and Krasnova V** 2010 Visual laterality of calf-mother interactions in wild whales. *PLoS ONE* 5: e13787. <https://doi.org/10.1371/journal.pone.0013787>
- Karenina K, Giljov A, Ivkovich T, Burdin A and Malashichev Y** 2013 Lateralization of spatial relationships between wild mother and infant orcas, *Orcinus orca*. *Animal Behavior* 86: 1225-1231. <https://doi.org/10.1016/j.anbehav.2013.09.025>
- Kuczaj S, Lacinak T, Fad O, Trone M, Solangi M and Ramos J** 2002 Keeping environmental enrichment enriching. *International Journal of Comparative Psychology* 15: 127-137
- Kuczaj S and Makecha R** 2008 The role of play in the evolution and ontogeny of contextually flexible communication. In: Oller K and Griebel U (eds) *Evolution of Communicative Flexibility* pp 253-278. MIT Press: Cambridge, USA. <https://doi.org/10.7551/mitpress/9780262151214.003.0012>
- Kuczaj SA and Eskelinen HC** 2014 Why do dolphins play. *Animal Behavior and Cognition* 1: 113-127. <https://doi.org/10.12966/abc.05.03.2014>
- Kuczaj SA, Highfill LE, Makecha RN and Byerly HC** 2013 Why do dolphins smile? A comparative perspective on dolphin emotions and emotional expressions. In: Watanabe S and Kuczaj S (eds) *Emotions of Animals and Humans: Comparative Perspectives* pp 63-85. Springer: New York, NY, USA
- Leliveld LM, Langbein J and Puppe B** 2013 The emergence of emotional lateralization: evidence in non-human vertebrates and implications for farm animals. *Applied Animal Behaviour Science* 145: 1-14. <https://doi.org/10.1016/j.applanim.2013.02.002>
- Lilly JC** 1963 Distress call of the bottlenose dolphin: stimuli and evoked behavioral responses. *Science* 139: 116-118. <https://doi.org/10.1126/science.139.3550.116>
- Mann J, Connor RC, Tyack PL and Whitehead H** 2000 *Cetacean Societies: Field Studies of Dolphins and Whales*. University of Chicago Press: Chicago, IL, USA
- Mann J and Kemps C** 2003 The effects of provisioning on maternal care in wild bottlenose dolphins, Shark Bay, Australia. *Books Online* 2006: 304-320
- Maple TL** 2007 Toward a science of welfare for animals in the zoo. *Journal of Applied Animal Welfare Science* 10: 63-70. <https://doi.org/10.1080/10888700701277659>
- Mason GJ** 1991 Stereotypies: a critical review. *Animal Behaviour* 41: 1015-1037. [https://doi.org/10.1016/S0003-3472\(05\)80640-2](https://doi.org/10.1016/S0003-3472(05)80640-2)
- Mason GJ and Latham NR** 2004 Can't stop, won't stop: is stereotypy a reliable animal welfare indicator? *Animal Welfare* 13: 57-69
- Mason GJ and Rushen J** 2008 *Stereotypic Animal Behaviour: Fundamentals and Applications to Welfare*. CABI: Wallingford, UK
- Mason GJ and Veasey JS** 2010 How should the psychological well-being of zoo elephants be objectively investigated? *Zoo Biology* 29: 237-255. <https://doi.org/10.1002/zoo.20256>
- Maze-Foley K and Wursig B** 2002 Patterns of social affiliation and group composition for bottlenose dolphins (*Tursiops truncatus*) in San Luis Pass, Texas. *Gulf of Mexico Science* 20: 122-134
- McCowan B, Marino L, Vance E, Walke L and Reiss D** 2000 Bubble ring play of bottlenose dolphins (*Tursiops truncatus*): implications for cognition. *Journal of Comparative Psychology* 114: 98. <https://doi.org/10.1037/0735-7036.114.1.98>
- Meehan CL and Mench JA** 2007 The challenge of challenge: can problem solving opportunities enhance animal welfare? *Applied Animal Behaviour Science* 102: 246-261. <https://doi.org/10.1016/j.applanim.2006.05.031>
- Mendl M, Burman OH, Parker RM and Paul ES** 2009 Cognitive bias as an indicator of animal emotion and welfare: emerging evidence and underlying mechanisms. *Applied Animal Behaviour Science* 118: 161-181. <https://doi.org/10.1016/j.applanim.2009.02.023>
- Mendl M, Burman OH and Paul ES** 2010 An integrative and functional framework for the study of animal emotion and mood. *Proceedings of the Royal Society of London B: Biological Sciences* 277: 2895-2904. <https://doi.org/10.1098/rspb.2010.0303>
- Miller LJ, Kuczaj S and Herzing D** 2011a Stereotypic behavior in wild marine carnivores? *Zoo Biology* 30: 365-370. <https://doi.org/10.1002/zoo.20347>
- Miller SJ, Mellen J, Greer T and Kuczaj SA** 2011b The effects of education programmes on Atlantic bottlenose dolphin (*Tursiops truncatus*) behaviour. *Animal Welfare* 20: 159-172
- Millman ST** 2007 Sickness behaviour and its relevance to animal welfare assessment at the group level. *Animal Welfare* 16: 123-125
- Moberg GP and Mench JA** 2000 *Biology of Animal Stress: Basic Principles and Implications for Animal Welfare*. CABI Publishing: Wallingford, UK. <https://doi.org/10.1079/9780851993591.0000>
- Mononen J, Moller SH, Hansen SW, Hovland AL, Koistinen T, Lidfors L, Malkmvist J and Ahola L** 2012 The development of on-farm welfare assessment protocols for foxes and mink: the WelFur project. *Animal Welfare* 21: 363-371. <https://doi.org/10.7120/09627286.21.3.363>
- Möstl E and Palme R** 2002 Hormones as indicators of stress. *Domestic Animal Endocrinology* 23: 67-74. [https://doi.org/10.1016/S0739-7240\(02\)00146-7](https://doi.org/10.1016/S0739-7240(02)00146-7)

- Müller M and Bossley M** 2002 Solitary bottlenose dolphins in comparative perspective. *Aquatic Mammals* 28: 298-307
- Neto MP, Silveira M and dos Santos ME** 2016 Training bottlenose dolphins to overcome avoidance of environmental enrichment objects in order to stimulate play activities. *Zoo Biology* 35: 210-215. <https://doi.org/10.1002/zoo.21282>
- Newberry RC** 1995 Environmental enrichment: increasing the biological relevance of captive environments. *Applied Animal Behaviour Science* 44: 229-243. [https://doi.org/10.1016/0168-1591\(95\)00616-Z](https://doi.org/10.1016/0168-1591(95)00616-Z)
- Ohi F and van der Staay FJ** 2012 Animal welfare: At the interface between science and society. *The Veterinary Journal* 192: 13-19. <https://doi.org/10.1016/j.tvjl.2011.05.019>
- Orbach DN, Packard JM, Piwetz S and Würsig B** 2015 Sex-specific variation in conspecific-acquired marking prevalence among dusky dolphins (*Lagenorhynchus obscurus*). *Canadian Journal of Zoology* 93: 383-390. <https://doi.org/10.1139/cjz-2014-0302>
- Ortiz RM and Worthy GAJ** 2000 Effects of capture on adrenal steroid and vasopressin concentrations in free-ranging bottlenose dolphins (*Tursiops truncatus*). *Comparative Biochemistry and Physiology Part A* 125: 317-324. [https://doi.org/10.1016/S1095-6433\(00\)00158-6](https://doi.org/10.1016/S1095-6433(00)00158-6)
- Overstrom NA** 1983 Association between burst-pulse sounds and aggressive behavior in captive Atlantic bottlenosed dolphins (*Tursiops truncatus*). *Zoo Biology* 2: 93-103. <https://doi.org/10.1002/zoo.1430020203>
- Pace DS** 2000 Fluke: made bubble rings as toys in bottlenose dolphin calves (*Tursiops truncatus*). *Aquatic Mammals* 26: 57-64
- Pack AA** 2010 The synergy of laboratory and field studies of dolphin behavior and cognition. *International Journal of Comparative Psychology* 23: 538-565
- Parsons KM, Durban JW, Claridge DE, Herzing DL, Balcomb KC and Noble LR** 2006 Population genetic structure of coastal bottlenose dolphins (*Tursiops truncatus*) in the northern Bahamas. *Marine Mammal Science* 22: 276-298. <https://doi.org/10.1111/j.1748-7692.2006.00019.x>
- Paul ES, Harding EJ and Mendl M** 2005 Measuring emotional processes in animals: the utility of a cognitive approach. *Neuroscience & Biobehavioral Reviews* 29: 469-491. <https://doi.org/10.1016/j.neubiorev.2005.01.002>
- Paulos RD, Trone M and Kuczaj SA II** 2010 Play in wild and captive cetaceans. *International Journal of Comparative Psychology* 23: 701-722
- Pedernera-Romano C, Valdez RA, Singh S, Chiappa X, Romano MC and Galindo F** 2006 Salivary cortisol in captive dolphins (*Tursiops truncatus*): a non-invasive. *Animal Welfare* 15: 359-362
- Perelberg A and Schuster R** 2009 Bottlenose dolphins (*Tursiops truncatus*) prefer to cooperate when petted: Integrating proximate and ultimate explanations ii. *Journal of Comparative Psychology* 123: 45-55. <https://doi.org/10.1037/a0013585>
- Pettis HM, Rolland RM, Hamilton PK, Brault S, Knowlton AR and Kraus SD** 2004 Visual health assessment of North Atlantic right whales (*Eubalaena glacialis*) using photographs. *Canadian Journal of Zoology* 82: 8-19. <https://doi.org/10.1139/z03-207>
- Pryor K and Norris KS** 1998 *Dolphin Societies: Discoveries and Puzzles*. University of California Press: London, UK
- Reif JS, Fair PA, Adams J, Joseph B, Kilpatrick DS, Sanchez R, Goldstein JD, Townsend FI Jr, McCulloch SD, Mazzoil M, Zolman ES, Hansen LJ and Bossart GD** 2008 Evaluation and comparison of the health status of Atlantic bottlenose dolphins from the Indian River Lagoon, Florida, and Charleston, South Carolina. *Journal of the American Veterinary Medical Association* 233: 299-307. <https://doi.org/10.2460/javma.233.2.299>
- Reiss D and Marino L** 2001 Mirror self-recognition in the bottlenose dolphin: A case of cognitive convergence. *Proceedings of the National Academy of Sciences* 98: 5937-5942. <https://doi.org/10.1073/pnas.101086398>
- Roche JR, Friggens NC, Kay JK, Fisher MW, Stafford KJ and Berry DP** 2009 Invited review: Body condition score and its association with dairy cow productivity, health, and welfare. *Journal of Dairy Science* 92: 5769-5801. <https://doi.org/10.3168/jds.2009-2431>
- Roe E, Buller H and Bull J** 2011 The performance of farm animal assessment. *Animal Welfare* 20: 69-78
- Rogers LJ** 2002 Lateralization in vertebrates: its early evolution, general pattern, and development. *Advances in the Study of Behavior, Vol 31* pp 107-161. Academic Press Inc: San Diego, CA, USA. [https://doi.org/10.1016/S0065-3454\(02\)80007-9](https://doi.org/10.1016/S0065-3454(02)80007-9)
- Rogers LJ** 2010 Relevance of brain and behavioural lateralization to animal welfare. *Applied Animal Behaviour Science* 127: 1-11. <https://doi.org/10.1016/j.applanim.2010.06.008>
- Rushen J, Butterworth A and Swanson JC** 2011 Animal behavior and well-being symposium: farm animal welfare assurance: science and application. *Journal of Animal Science* 89: 1219-1228. <https://doi.org/10.2527/jas.2010-3589>
- Russell JA** 2003 Core affect and the psychological construction of emotion. *Psychological Review* 110: 145. <https://doi.org/10.1037/0033-295X.110.1.145>
- Sakai M, Hishii T, Takeda S and Kohshima S** 2006 Laterality of flipper rubbing behaviour in wild bottlenose dolphins (*Tursiops aduncus*): Caused by asymmetry of eye use? *Behavioural Brain Research* 170: 204-210. <https://doi.org/10.1016/j.bbr.2006.02.018>
- Samuels A and Gifford T** 1997 A quantitative assessment of dominance relations among bottlenose dolphins. *Marine Mammal Science* 13: 70-99. <https://doi.org/10.1111/j.1748-7692.1997.tb00613.x>
- Schusterman R, Thomas JA and Wood FG** 2013 *Dolphin Cognition and Behavior: A Comparative Approach*. Psychology Press: London, UK
- Schwacke LH, Smith CR, Townsend FI, Wells RS, Hart LB, Balmer BC and Rowles TK** 2014 Health of common bottlenose dolphins (*Tursiops truncatus*) in Barataria Bay, Louisiana, following the Deepwater Horizon oil spill. *Environmental Science and Technology* 48: 93-103. <https://doi.org/10.1021/es403610f>
- Scott EM, Mann J, Watson-Capps JJ, Sargeant BL and Connor RC** 2005 Aggression in bottlenose dolphins: evidence for sexual coercion, male-male competition, and female tolerance through analysis of tooth-rake marks and behaviour. *Behaviour* 142: 21-44. <https://doi.org/10.1163/1568539053627712>
- Selye H** 1975 Stress and distress. *Comprehensive Therapy* 1: 9-13
- Siegford JM** 2013 Multidisciplinary approaches and assessment techniques to better understand and enhance zoo nonhuman animal welfare. *Journal of Applied Animal Welfare Science* 16: 300-318. <https://doi.org/10.1080/10888705.2013.827914>

- Silber GK and Fertl D** 1995 Intentional beaching by bottlenose dolphins (*Tursiops truncatus*) in the Colorado River Delta, Mexico. *Aquatic Mammals* 21: 183-186
- Small RJ and DeMaster DP** 1995 Survival of five species of captive marine mammals. *Marine Mammal Science* 11: 209-226. <https://doi.org/10.1111/j.1748-7692.1995.tb00519.x>
- Smolker R, Richards A, Connor RC, Mann J and Berggren P** 1997 Sponge carrying by dolphins (Delphinidae, *Tursiops* spp): a foraging specialization involving tool use? *Ethology* 103: 454-465. <https://doi.org/10.1111/j.1439-0310.1997.tb00160.x>
- Sneddon LU, Elwood RW, Adamo SA and Leach MC** 2014 Defining and assessing animal pain. *Animal Behaviour* 97: 201-212. <https://doi.org/10.1016/j.anbehav.2014.09.007>
- Sobel N, Supin AY and Myslobodsky MS** 1994 Rotational swimming tendencies in the dolphin (*Tursiops truncatus*). *Behavioural Brain Research* 65: 41-45. [https://doi.org/10.1016/0166-4328\(94\)90071-X](https://doi.org/10.1016/0166-4328(94)90071-X)
- Spruijt BM, van den Bos R and Pijlman FT** 2001 A concept of welfare based on reward evaluating mechanisms in the brain: anticipatory behaviour as an indicator for the state of reward systems. *Applied Animal Behaviour Science* 72: 145-171. [https://doi.org/10.1016/S0168-1591\(00\)00204-5](https://doi.org/10.1016/S0168-1591(00)00204-5)
- Stockin KA, Lusseau D, Binedell V, Wiseman N and Orams MB** 2008 Tourism affects the behavioural budget of the common dolphin *Delphinus* spp. in the Hauraki Gulf, New Zealand. *Marine Ecology-Progress Series* 355: 287. <https://doi.org/10.3354/meps07386>
- Swaigood RR** 2007 Current status and future directions of applied behavioral research for animal welfare and conservation. *Applied Animal Behaviour Science* 102: 139-162. <https://doi.org/10.1016/j.applanim.2006.05.027>
- Tamaki N, Morisaka T and Taki M** 2006 Does body contact contribute towards repairing relationships? The association between flipper-rubbing and aggressive behavior in captive bottlenose dolphins. *Behavioural Processes* 73: 209-215. <https://doi.org/10.1016/j.beproc.2006.05.010>
- Thomson CA and Geraci JR** 1986 Cortisol, aldosterone, and leucocytes in the stress response of bottlenose dolphins, *Tursiops truncatus*. *Canadian Journal of Fisheries and Aquatic Sciences* 43: 1010-1016. <https://doi.org/10.1139/f86-125>
- Ugaz C, Valdez RA, Romano MC and Galindo F** 2013 Behavior and salivary cortisol of captive dolphins (*Tursiops truncatus*) kept in open and closed facilities. *Journal of Veterinary Behavior: Clinical Applications and Research* 8: 285-290. <https://doi.org/10.1016/j.jveb.2012.10.006>
- Veasey J** 2006 Concepts in the care and welfare of captive elephants. *International Zoo Yearbook* 40: 63-79. <https://doi.org/10.1111/j.1748-1090.2006.00063.x>
- Veissier I, Butterworth A, Bock B and Roe E** 2008 European approaches to ensure good animal welfare. *Applied Animal Behaviour Science*. 113: 279-297. <https://doi.org/10.1016/j.applanim.2008.01.008>
- Venn-Watson SK, Jensen ED and Ridgway SH** 2011 Evaluation of population health among bottlenose dolphins (*Tursiops truncatus*) at the United States Navy Marine Mammal Program. *Journal of the American Veterinary Medical Association* 238: 356-360. <https://doi.org/10.2460/javma.238.3.356>
- Ventre J and Jett J** 2015 Killer whales, theme parks and controversy: an exploration of the evidence. In: Markwell K (ed) *Animals and Tourism: Understanding Diverse Relationships*. Channel View Productions: Bristol, UK
- Vicino GA and Marcacci ES** 2015 Intensity of play behavior as a potential measure of welfare: A novel method for quantifying the integrated intensity of behavior in African elephants. *Zoo Biology* 34: 492-496. <https://doi.org/10.1002/zoo.21238>
- Walker MD, Duggen G, Roulston N, Van Slack AG and Mason G** 2012 Negative affective states and their effects on morbidity, mortality and longevity. *Animal Welfare* 21: 497-509. <https://doi.org/10.7120/09627286.21.4.497>
- Waples KA and Gales NJ** 2002 Evaluating and minimising social stress in the care of captive bottlenose dolphins (*Tursiops aduncus*). *Zoo Biology* 21: 5-26. <https://doi.org/10.1002/zoo.10004>
- Watters JV** 2014 Searching for behavioral indicators of welfare in zoos: Uncovering anticipatory behavior. *Zoo Biology* 33: 251-256. <https://doi.org/10.1002/zoo.21144>
- Weary DM, Niel L, Flower FC and Fraser D** 2006 Identifying and preventing pain in animals. *Applied Animal Behaviour Science* 100: 64-76. <https://doi.org/10.1016/j.applanim.2006.04.013>
- Webster J** 2005 *Animal Welfare: Limping towards Eden*. Blackwell Publishing Ltd: Oxford, UK. <https://doi.org/10.1002/9780470751107>
- Welfare Quality**® 2009a *Welfare Quality*® Assessment Protocol for Cattle (Fattening Cattle, Dairy cows, Veal Calves). Welfare Quality® Consortium: Lelystad, The Netherlands
- Welfare Quality**® 2009b *Welfare Quality*® Assessment Protocol for Pigs. Welfare Quality® Consortium: Lelystad, The Netherlands
- Welfare Quality**® 2009c *Welfare Quality*® Assessment Protocol for Poultry. Welfare Quality® Consortium: Lelystad, The Netherlands
- Wells RS** 1991 The role of long-term study in understanding the social structure of a bottlenose dolphin community. In: Pryor K and Norris KS (eds) *Dolphin Societies: Discoveries and Puzzles* pp 199-225. University of California Press: London, UK
- Wells RS** 2009 Learning from nature: bottlenose dolphin care and husbandry. *Zoo Biology* 28: 635-651. <https://doi.org/10.1002/zoo.20252>
- Wells RS and Scott MD** 1999 Bottlenose dolphin: *Tursiops truncatus* (Montagu, 1821). In: Ridgway SH and Harrison R (eds) *Handbook of Marine Mammals: The Second Book of Dolphins and Porpoises* pp 137-182. Academic Press: San Diego, CA
- Whitham JC and Wielebnowski N** 2013 New directions for zoo animal welfare science. *Applied Animal Behaviour Science* 147: 247-260. <https://doi.org/10.1016/j.applanim.2013.02.004>
- Würbel H and Garner JP** 2007 Refinement of rodent research through environmental enrichment and systematic randomization. *NC3Rs* 9: 1-9
- Yaman S, von Fersen L, Dehnhardt G and Güntürkün O** 2003 Visual lateralization in the bottlenose dolphin (*Tursiops truncatus*): Evidence for a population asymmetry? *Behavioural Brain Research* 142: 109-114. [https://doi.org/10.1016/S0166-4328\(02\)00385-6](https://doi.org/10.1016/S0166-4328(02)00385-6)
- Yeates JW and Main DCJ** 2008 Assessment of positive welfare: a review. *The Veterinary Journal* 175: 293-300. <https://doi.org/10.1016/j.tvjl.2007.05.009>