

# Erratum

## Volume 8(4)

The author Rosalind J. Neuman wishes to offer the following correction to Table 2 on p. 397 of the following article from the previous issue: Neuman, R. J., Sitdhiraksa, N., Reich, W., Ji, T. H.-C., Joyner, C. A, Sun, L.-W., & Todd, R. D. (2005). Estimation of prevalence of DSM-IV and latent class-defined ADHD subtypes in a population-based sample of child and adolescent twins. *Twin Research and Human Genetics*, 8, 392–401.

**Table 2**  
Population Prevalence (%) of Latent Class ADHD Subtypes by Sex and Twin Type

Latent class	Total sample		Male			Female			
	(N = 1472)	Total (N = 923)	MZ (N = 275)	DZ same-sex pairs (N = 330)	DZ opposite-sex pairs (N = 318)	Total (N = 549)	MZ (N = 105)	DZ same-sex pairs (N = 125)	DZ opposite-sex pairs (N = 319)
Few symptoms N = 795	75.0 (71.1–78.8)	69.5 <sup>a</sup> (63.4–75.1)	69.5 (58.2–80.1)	65.7 (53.0–76.7)	73.0 (65.0–80.0)	82.5 <sup>a</sup> (77.6–87.1)	88.2 (77.2–94.8)	76.7 (67.6–85.1)	81.9 (73.3–89.4)
Mild inattentive N = 196	6.4 (4.6–8.4)	7.9 <sup>b</sup> (5.4–10.9)	8.4 (4.1–13.5)	9.2 (3.3–16.5)	6.5 (3.9–10.5)	4.2 <sup>b</sup> (1.9–6.8)	2.7 (0.3–6.4)	5.0 (0.6–10.8)	4.8 (1.2–9.6)
Talkative–mild impulsive N = 66	4.4 (2.6–6.4)	4.4 (1.8–7.2)	7.7 (2.7–13.2)	2.3 (0.0–7.1)	1.8 (0.0–5.4)	4.6 (2.0–7.8)	3.8 (0.2–9.1)	6.7 (1.2–13.5)	3.7 (0.3–8.7)
Mild combined N = 128	3.8 (2.6–5.3)	4.8 (3.1–7.0)	4.9 (2.0–8.9)	2.9 (1.9–4.1)	6.1 (2.5–11.0)	2.5 (1.2–4.4)	0.8 (0.0–1.6)	2.2 (0.9–3.6)	4.5 (1.0–9.5)
Mild hyperactive N = 52	3.5 (1.9–5.3)	5.3 <sup>c</sup> (2.6–8.5)	4.2 (0.5–8.9)	10.5 (3.8–19.7)	2.5 (0.4–5.9)	1.5 <sup>c</sup> (0.2–3.3)	0.0 (0.0–0.0)	3.1 (0.3–7.5)	1.8 (0.0–5.3)
Severe combined N = 118	3.3 (2.1–4.9)	3.5 (2.5–4.8)	1.7 (0.8–2.6)	3.3 (2.3–4.5)	5.5 (3.1–9.4)	2.8 (0.8–5.7)	3.8 (0.0–10.7)	3.1 (0.3–7.6)	1.3 (0.7–2.1)
Severe inattentive N = 94	2.5 (1.8–3.4)	3.7 <sup>d</sup> (2.5–5.4)	2.1 (1.1–3.0)	5.6 (2.8–9.9)	4.1 (1.8–7.6)	0.7 <sup>b</sup> (0.3–1.1)	0.8 (0.0–1.7)	1.3 (0.3–2.6)	0.3 (0.0–0.7)
Severe hyperactive N = 23	1.1 (0.3–2.2)	0.9 (0.2–2.1)	1.5 (0.0–4.2)	0.4 (0.0–0.8)	0.7 (0.2–1.2)	1.2 (0.0–3.1)	0.0 (0.0–0.0)	2.0 (0.0–6.1)	1.7 (0.0–5.2)

Note: a,b,c,d = prevalence difference in males versus females: a =  $\chi^2 = 11.3, p = 8 \times 10^{-4}$ ; b =  $\chi^2 = 3.9, p = .05$ ; c =  $\chi^2 = 4.9, p = .03$ ; d =  $\chi^2 = 14.4, p = 1.5 \times 10^{-4}$ .