

Antioxidant micronutrient supplementation increases erythrocyte membrane fluidity in adults from a rural Chinese community

Xiu-Xia Han¹, Ming Zhang², Ai-Guo Ma^{1*}, Sheng Ge³, Xue-Xiang Shi⁴, Yong-Ye Sun¹, Qiu-Zhen Wang¹ and Hui Liang¹

¹The Institute of Human Nutrition, Medical College of Qingdao University, 38# Dengzhou Road, Qingdao 266021, People's Republic of China

²Department of Clinical Nutrition, Weifang People's Hospital, Shandong, People's Republic of China

³Department of Clinical Nutrition, Shanghai 6th People's Hospital, Shanghai, People's Republic of China

⁴Qingdao Center of Disease Control, Qingdao, People's Republic of China

(Received 22 October 2010 – Revised 21 March 2011 – Accepted 22 March 2011 – First published online 7 June 2011)

Abstract

The objective of the present study was to investigate age-related differences in erythrocyte membrane fluidity (EMF) and changes in antioxidant capacity following supplementation. A total of seventy-four children were randomly divided into two groups: group A1 was the placebo-controlled group and group A2 was supplemented daily with 600 µg retinol, 1.0 mg β-carotene, 100 mg tocopherol, 300 mg ascorbic acid and 200 µg Se. A total of ninety young people were randomly divided into B1 and B2 groups, and ninety-one elderly subjects were divided into C1 and C2 groups. Groups B1 and C1 were placebo-controlled groups, and groups B2 and C2 were daily supplemented with 900 µg retinol, 1.5 mg β-carotene, 200 mg tocopherol, 500 mg ascorbic acid and 400 µg Se. Results showed that plasma malondialdehyde (MDA) was 5.35 µmol/l in children, which was lower than in young and elderly people. The MDA levels of the young and elderly individuals in the treated groups were significantly lower compared with the control groups, but the supplementation did not alter MDA levels in children. At baseline, there was a lower value of polarisation (ρ) and microviscosity (η) in children, indicating a higher EMF, than in both the young and elderly subjects. After the 2-month trial, the ρ and η values of young and elderly subjects in the treated groups decreased significantly in comparison with the placebo groups, indicating an increase in EMF. In conclusion, there was a background of higher MDA levels and lower EMF in young and elderly people than in children, which could be improved by antioxidant supplementation.

Key words: Antioxidants; Erythrocyte membrane fluidity; Oxidative stress; Children; Elderly people

Free radical species can be endogenously or exogenously produced⁽¹⁾. A major target of free radicals is the cell membrane, which is involved in the control of cell function through integral maintenance of the spatial and intermolecular arrangements of its components. Because of their susceptibility to oxidation, erythrocytes have been used as a model to investigate oxidative damage in biomembranes^(2,3).

There have been reports of oxidant–antioxidant imbalances and increased free radical activity^(4,5), which may damage cell membranes resulting changes in membrane fluidity and cell function⁽⁶⁾. Pathological processes in ageing have been associated with free radical-induced damage⁽⁷⁾. Elderly people are subject to a decrease in antioxidant defences (25%) and higher levels of oxidative stress, which causes an increase in plasma hydroperoxide levels (18%)⁽⁸⁾. A variety of antioxidant systems exist *in vivo* to remove free radicals and limit tissue

damage. Supplementation with antioxidants, such as vitamin A⁽⁹⁾, ascorbic acid or Se, reduces oxidative stress and enhances total antioxidant status^(10,11).

Many *in vitro* and *in vivo* studies have shown that several parameters related to erythrocyte function and integrity are negatively affected by increased oxidative stress. Data reported in the literature^(12,13) concerning membrane characteristics in the elderly are often conflicting, due to the difficulty in examining oxidative stress damage independently of its frequently associated pathologies. Thus, the mechanisms underlying these phenomena are not yet understood, and little information is available about erythrocyte membrane modification resulting from oxidative stress and ageing and about membrane fluidity in children, young and elderly people. Data have been reported^(14–16), indicating that senescence is associated with an increase in both oxidant generation rates

Abbreviations: η , microviscosity; ρ , polarisation; EMF, erythrocyte membrane fluidity; MDA, malondialdehyde; GSH-Px, glutathione peroxidase.

* **Corresponding author:** Professor A.-G. Ma, fax +86 532 83812434, email aiguo502@hotmail.com

and in the susceptibility of tissues to oxidative damage during ageing. Therefore, the objectives of the present study were to investigate whether there are any differences in erythrocyte membrane fluidity (EMF) between individuals of various ages and to describe any changes in antioxidant capacity and membrane fluidity in children, young and elderly people after supplementation with multiple micronutrients. We carried out the study in a rural population, where poor nutrition was expected to contribute to lack of micronutrients and increased oxidative stress, and where we could expect to observe bigger changes in the measured variables after supplementation. Nevertheless, the present results will be helpful for improving our understanding whether micronutrient supplementation is effective across all populations in decreasing oxidative stress and improving EMF.

Subjects and methods

Subjects

A total of 255 healthy persons including seventy-four children (aged 8–10 years) from a primary school, ninety young people (aged 18–22 years) from a college and ninety-one elderly (aged 60–75 years) from a community were recruited. A total of seventy-four children were randomly divided into A1 and A2 groups; group A1 (n 36) was the placebo-controlled group and group A2 was supplemented daily with 600 μg of retinol (as retinyl palmitate), 1.0 mg of β -carotene, 100 mg of tocopherol (as DL- α -tocopherol acetate), 300 mg of ascorbic acid and 200 μg of Se (as sodium selenite). A total of ninety young people were randomly divided into B1 and B2 groups, and ninety-one elderly subjects were divided into C1 and C2 groups. Groups B1 (n 45) and C1 (n 48)

were placebo-controlled groups, and groups B2 (n 45) and C2 (n 43) were supplemented with 900 μg of retinol, 1.5 mg of β -carotene, 200 mg of tocopherol, 500 mg of ascorbic acid and 400 μg of Se. The capsules used in these treatment groups were labelled and manufactured by Hurun Company, Ypsilanti, MI, USA (a Chinese food additive company in Beijing). The trial participants and the research team were unaware of the treatment assignment. The trial was deblinded after analysis of its primary outcomes. The characteristics of the subjects are shown in Table 1.

Potential study subjects were informed about the study content and design via a formal presentation and an individual interview. Written, informed consent was obtained from the adult participants and the parents of child subjects before entering the trial. The subjects underwent a clinical examination to determine their suitability for participation in the study. Exclusion criteria were severe or chronic illness; smoking; a history of gastrointestinal surgery or chronic bowel disease; a current metabolic illness, such as diabetes mellitus, untreated hyper- or hypothyroidism or a severe lipid metabolism disorder; and regular intake of nutritional supplements (i.e. vitamin/mineral supplements or micronutrient-enriched foods), unless discontinued for at least 4 weeks before entry into the study and throughout the investigation. After ascertainment of eligibility, consenting subjects were enrolled in the study, subjected to a baseline interview and were given their allocated supplements once a week, which were to be taken daily for a period of 2 months. The subjects were examined or visited once each week by teachers or co-investigators trained to replenish supplements and to monitor compliance by counting and recording the number of supplements that were taken.

Table 1. General characteristics and indicators in children, young and elderly people (Mean values and standard deviations)

Items	Children		Young people		Elderly people		P^*
	Mean	SD	Mean	SD	Mean	SD	
Characteristics							
n		74		90		91	
Age (years)	10.07	0.69	18.59	0.99	65.18	8.06	0.000
Height (cm)	138.47	7.37	168.38	8.60	161.67	8.03	0.000
Weight (kg)	32.63	6.11	59.99	9.52	62.70	9.36	0.000
BMI (kg/m^2)†	–		21.09 ^a	2.41	24.00	3.23	0.000
Micronutrient status							
β -Carotene ($\mu\text{mol}/\text{l}$)	1.90	0.31	1.99	0.50	1.86	0.38	0.090
Retinol ($\mu\text{mol}/\text{l}$)	1.80	0.26	1.84	0.40	1.89	0.28	0.240
Tocopherol ($\mu\text{mol}/\text{l}$)	12.50	3.22	13.36	2.95	13.00	1.84	0.124
Ascorbic acid ($\mu\text{mol}/\text{l}$)	30.98 ^a	13.01	27.89	10.06	24.46	14.65	0.005
Se ($\mu\text{mol}/\text{l}$)	0.92	0.15	0.80	0.43	0.84	0.44	0.151
Oxidative stress parameters							
SOD (U/ml)	99.38	19.68	92.46	32.52	96.98	17.34	0.179
GSH-Px (IU/ml)	129.83	42.12	139.15	40.06	131.38	54.42	0.370
MDA ($\mu\text{mol}/\text{l}$)	5.35 ^{a,b}	1.59	7.07	2.65	6.73	3.51	0.000
Erythrocyte membrane fluidity							
ρ	0.313 ^{a,b}	0.033	0.326 ^a	0.037	0.337	0.024	0.000
η	4.48 ^{a,b}	0.85	5.50	0.93	5.73	1.35	0.000

SOD, superoxide dismutase; GSH-Px, glutathione peroxidase; MDA, malondialdehyde; ρ , polarisation; η , microviscosity.

^{a,b} Mean values with unlike superscript letters were significantly different between elderly people and young people.

* ANOVA.

† BMI of children could not be calculated because the formula is not suitable for children.

The study was approved by the ethical review committees of the Medical College of Qingdao University, Qingdao, China (QY-ETC20031120). Written consent was given by each subject at the beginning of the trial.

Blood sample treatment

Blood samples were taken from all subjects at baseline and at the end of the trial and processed immediately for storage. Specimens were stored in polypropylene vials at -80°C for up to 3 months before batch analysis. It has been determined that vitamins C, retinol, β -carotene, vitamin E and Se are stable under these storage conditions⁽¹⁷⁾.

Micronutrient concentrations in plasma

Retinol, β -carotene and tocopherols were determined in plasma samples according to the method of Hess *et al.*⁽¹⁸⁾. Analyses were performed using HPLC equipped with an LC-10AD pump. A standard curve for each analyte was constructed from authentic standards (Sigma, St Louis, MO, USA). Standard concentrations were calculated on the basis of their known extinction coefficients. Total ascorbic acid concentrations in the subjects' plasma were determined spectrophotometrically, using the 2,4-dinitrophenylhydrazine method. Ascorbic acid was converted to dehydroascorbic acid in the presence of thiourea and CuSO_4 ^(19,20). Plasma Se levels were measured using a standardised fluorometric method⁽²¹⁾.

The activities of superoxide dismutase and glutathione peroxidase (GSH-Px) were determined in plasma⁽²²⁾ as U/ml and IU/ml, respectively. The malondialdehyde (MDA) concentration was determined using the thiobarbituric acid reaction. MDA, an end product of fatty acid peroxidation, reacts with thiobarbituric acid to form a coloured complex, and the concentrations of this complex were calculated by comparing the absorbance values of the samples with those of standard MDA solutions. Results are expressed as nmol/ml plasma⁽²³⁾.

Preparation of erythrocyte membranes

Fresh erythrocytes were washed three times with isotonic saline (0.15 M-NaCl, 10 mM-Tris-HCl, pH 7.4), and the buffy coat and plasma were removed each time. Erythrocyte membranes were prepared by haemolysing washed erythrocytes in 5 mM phosphate buffer, pH 8, according to a method described previously⁽²⁴⁾. Erythrocyte membranes (ghosts) were prepared by hypotonic haemolysis according to Raccach *et al.*⁽²⁵⁾. Membrane (ghost) proteins were determined according to Lowry *et al.*⁽²⁶⁾, and aliquots were immediately frozen and stored at -80°C until use for membrane fluidity determination.

Erythrocyte membrane fluidity measurement

EMF can be measured by fluorescence polarisation (ρ) and microviscosity (η). Erythrocytes were washed using physiological saline and suspended in a PBS solution (pH 7.4) at 0.01 mol/l. Then, 1,6-diphenyl-1,3,5-hexatriene (2×10^{-6} mol/l)

was added to the erythrocyte suspension before being incubated in a water-bath at 25°C for 15 min^(27,28). The suspension was examined using a spectrofluorophotometer (Perkin-Elmer fluorescence spectrometer, LS-50) with an excitation wavelength of 360 nm and an emission wavelength of 430 nm⁽²⁹⁾, and the ρ and η of the erythrocyte membranes were calculated using the following formula:

$$\rho = I_{VV} - GI_{VH}/I_{VV} + GI_{VH}; \quad \eta = 2\rho/0.46 - \rho.$$

The indices V and H indicate the vertical and horizontal position of the polariser in the excitation and fluorescence beams, respectively. G is an instrumental correction factor equal to (I_{VV}/I_{VH}) . The subscript H refers to the horizontally polarised excitation beam. I_{VV} and I_{VH} represent the components of the corrected polarised emissions parallel and perpendicular to the vertical direction, respectively⁽³⁰⁾.

Statistical analysis

Data are presented as means and standard deviations. Variables were assessed for normality by the skewness and kurtosis test. The indicators of the vitamin and oxidative stress parameters exhibited a normal distribution. The skewed data of ρ and η were not distributed and were logarithmically transformed for statistical analysis and then back-transformed to their natural units for presentation in tables. The differences of all of the data between groups were determined using a general linear model ANOVA. The baseline variables were compared across treatment groups. $P < 0.05$ was considered as the significance level for all tests.

Results

In the intervention study, complete data were available for 238 individuals, including seventy-four children, ninety young and ninety-one elderly people representing 93.3% of the original 255 subjects. For seventeen subjects, insufficient volumes of plasma and whole blood were available for the measurement of micronutrient concentrations, oxidative stress and membrane fluidity analysis. There were no substantial differences in any of the baseline characteristics between the placebo and supplemented groups in children, young or elderly people.

The general characteristics, micronutrient status, oxidative stress and EMF in the children, young and elderly people investigated are shown in Table 1. The average levels of retinol, tocopherol and total ascorbic acid concentrations in the plasma were over the cut-off values for deficiencies and were 0.70, 11.6 and $11.4 \mu\text{mol/l}$ ^(31,32), respectively; however, the plasma level of total ascorbic acid in children was $30.98 \mu\text{mol/l}$, which was significantly higher than in young ($27.89 \mu\text{mol/l}$) and elderly ($24.46 \mu\text{mol/l}$) people.

At baseline, no differences of the superoxide dismutase or GSH-Px activities were found in the three age populations, and there were higher levels of MDA in young ($7.07 \mu\text{mol/l}$) and elderly people ($6.73 \mu\text{mol/l}$) compared with children ($5.35 \mu\text{mol/l}$). Compared with elderly people (0.337), there

Table 2. Comparison of plasma micronutrient levels between the supplemented and control groups* in the three populations (Mean values and standard deviations)

Vitamins	Children				Young people				Elderly people			
	A1		A2		B1		B2		C1		C2	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>n</i>	34		35		42		42		46		39	
β-Carotene (μmol/l)	1.87	0.31	2.19†	0.26	2.02	0.48	2.56†	0.71	1.82	0.36	2.34†	0.61
Retinol (μmol/l)	1.82	0.26	2.28†	0.43	1.87	0.46	2.68†	0.53	1.86	0.32	2.58†	1.25
Tocopherol (μmol/l)	12.19	3.06	14.27†	3.52	13.36	3.08	18.44†	3.84	12.90	1.95	15.74†	2.96
Ascorbic acid (μmol/l)	32.96	12.43	46.44†	19.61	26.17	8.37	46.86†	20.46	25.78	14.47	39.88†	17.87
Se (μmol/l)	0.87	0.15	1.10†	0.39	0.79	0.45	1.25†	0.54	0.80	0.42	1.05†	0.42

* Six groups: A1, B1 and C1 were the control groups, and A2, B2 and C2 were the supplemented groups of children, young and elderly people, respectively.

† Mean values were significantly different after supplementation from those of controls in children, young people and elderly people ($P < 0.05$, one-way ANOVA).

was a lower ρ value (0.326) in young subjects ($P < 0.001$), and the lowest values of ρ and η were 0.313 and 4.48, respectively, in children, which indicates a high EMF, as shown in Table 1.

After 2 months of supplementation, there were significant increases in plasma β-carotene, retinol, tocopherol, ascorbic acid and Se concentrations in the treated groups compared with the control groups in all three age groups. Moreover, there were significant increases in the plasma β-carotene, retinol, tocopherol, ascorbic acid and Se concentrations in the supplemented groups after the trial, compared with the controls and the baseline of children, young and elderly people, respectively (all $P < 0.05$; Table 2).

The changes in the oxidative stress parameters investigated are presented in Table 3. Compared with the control groups of children, young and elderly people, the increases in GSH-Px activities were 44.91, 90.12 and 107.21 IU/ml in the A2, B2 and C2 groups, respectively; and the levels of MDA decreased by 2.16 and 1.63 μmol/l ($P < 0.001$) in the B2 and C2 groups, respectively; however, there were no important changes in superoxide dismutase activities following the trial.

EMF was evaluated by measuring the fluorescence ρ and η ; lower values indicate better membrane fluidity. After the 2-month trial, the decreases in ρ and η were significantly greater in the supplemented groups of young and elderly people compared with the control groups. The differences in the ρ and η values were -0.023 and -0.93 for young subjects and -0.026 and -1.15 for elderly people, respectively.

However, the differences in the ρ and η values in the supplemented group of children were -0.009 and -0.34 , respectively, which is not significantly different than the child control group.

Discussion

The present study demonstrated age-related changes in MDA levels and EMF in children, young and elderly people. There was a lower membrane fluidity in young and elderly people, which was related to an age-dependent increase in erythrocyte MDA levels⁽³³⁾. Plasma MDA concentration and carbonyl content of plasma proteins were used as an index of lipid and protein oxidation, respectively⁽³⁴⁾. After 2 months of micronutrient supplementation, there were considerable increases in GSH-Px activity and plasma antioxidant micronutrient concentration and decreases in MDA levels in the three age populations. Moreover, there were significant improvements in the EMF in the supplemented groups of young and elderly people compared with the control groups, which are attributed to an increase in the antioxidant capacity, resulting in significant protection of erythrocytes against oxidative stress⁽³⁵⁾.

In the present study, we chose to use moderate doses of antioxidants, i.e. 900 μg retinol, 1.5 mg β-carotene, 200 mg α-tocopherol, 500 mg ascorbic acid and 400 μg Se for adult subjects. These doses were selected mainly based on previous studies taking into consideration the beneficial effect

Table 3. Comparison of indicator levels between the supplemented and control groups* in the three populations (Mean values and standard deviations)

Indicators	Children				Young people				Elderly people			
	A1		A2		B1		B2		C1		C2	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>n</i>	36		34		45		42		48		46	
SOD	100.4	21.6	90.6	31.4	86.9	32.6	85.2	35.9	94.6	19.7	93.0	21.7
GSH-Px	122.7	38.6	167.6†	37.6	139.4	41.8	229.3†	75.3	130.8	49.7	238.0†	47.4
MDA	5.54	1.46	4.39	0.93	7.38	2.57	5.22†	3.31	6.67	3.59	5.04†	1.16
ρ	0.314	0.028	0.305	0.027	0.325	0.032	0.303†	0.040	0.337	0.028	0.311†	0.025
η	4.49	0.90	4.15	0.57	5.42	1.08	4.49†	1.08	5.76	1.42	4.61†	1.44

SOD, superoxide dismutase; GSH-Px, glutathione peroxidase; MDA, malondialdehyde; ρ , polarisation; η , microviscosity.

* Six groups: A1, B1 and C1 were the control groups, and A2, B2 and C2 were the supplemented groups of children, young and elderly people, respectively.

† Mean values were significantly different after supplementation from those of controls in children, young people and elderly people ($P < 0.05$, one-way ANOVA).

in reducing oxidative stress or the lipid peroxidation process in ageing without obvious adverse effects. Fluorescence polarisation has been used to assess membrane fluidity in the past four decades, but it is now frequently also applied to other areas of study, including investigations of oxidative stress⁽³⁶⁾, erythrocyte aggregation⁽³⁷⁾ and membrane protein function⁽³⁸⁾. To date, few studies have evaluated changes in EMF in large populations, possibly because erythrocyte membrane preparation is quite laborious.

Around 70% of the total Chinese population lives in, often underdeveloped, rural areas⁽³⁹⁾. The present study was carried out under difficult circumstances in a poor, rural part of China, where nutritional problems are frequent⁽⁴⁰⁾. The people still live in lower socio-economic conditions, with low intake of meat, vegetables and fruit. We selected these subjects knowing that poor diet was likely to contribute to low antioxidant and micronutrient levels, which in turn gives rise to cellular oxidative stress. We would, therefore, anticipate that changes in antioxidant micronutrient status, membrane fluidity and decreasing oxidative stress after supplementation would be clearer in such subjects. Moreover, compliance was excellent, because the study subjects were motivated by the offer of free medical care and all subjects were visited weekly by village nurses, who counted leftover capsules, provided new supplies and gave support in case of any problems or questions related to the study.

Compared with children, the lower membrane fluidity observed in elderly people at baseline should be attributed to a higher rate of peroxidation, which might be due to age-related changes in the glutathione redox system⁽⁴¹⁾. The susceptibility of membranes to peroxidation is affected by the nature and physical state of the lipid bilayer⁽⁴²⁾. The present findings are in accordance with reported data in animal models^(43,44), and it has been observed that there were lower MDA levels and higher EMF in a verbascoside-treated group compared with the placebo group⁽⁴⁵⁾. It has been reported that positive correlations between age and GSH-Px and oxidative stress increase during the ageing process⁽⁴⁶⁾. Human studies showed that supplementation of ascorbic acid and α -tocopherol is useful in preventing bone loss linked to oxidative stress in the elderly⁽⁴⁷⁾, and might be beneficial in women with a low antioxidant status⁽⁴⁸⁾. An antioxidant micronutrient combination (800 mg DL- α -tocopherol acetate, 24 mg β -carotene, 1.0 g vitamin C, 200 μ g selenomethionine, 7.2 mg riboflavin, 80 mg niacin, 60 mg Zn and 5 mg Mn) can modulate biomarkers of oxidative stress and inflammation in human subjects⁽⁴⁹⁾, and dietary supplementation with vitamin E (50 mg/d) and vitamin C (150 mg/d) for 6 months decreased plasma oxidative damage and enhanced the erythrocyte activities of catalase and glutathione reductase⁽⁵⁰⁾. Additionally, it was found that adequate supplementation with the aforementioned vitamins led to increased quality of life in haemodialysis patients, from some clinical points of view⁽⁵¹⁾. Proanthocyanidin-rich grape seed extract has also been found to be an effective anti-ageing drug in preventing the oxidative stress-associated loss of membrane surface charge, which thereby maintains erythrocyte membrane integrity and functions in elderly

individuals⁽⁵²⁾. The present results and the reports described earlier show that antioxidant supplementation is beneficial to elderly people for decreasing oxidative stress and improving EMF.

In the young subjects we investigated, who came from a college, there were higher MDA levels (7.07 μ mol/l) and lower EMF, with higher ρ and η values at baseline than in children, which may be related to a greater amount of physical exercise. This leads to a decrease in antioxidant levels, which could promote an increase both in the markers of lipoprotein peroxidation and in damage to the erythrocyte membrane, with consequential modifications to membrane fluidity⁽⁵³⁾. The oxidative stress and EMF of young subjects in the present study was greatly improved following multiple micronutrient supplementations, which is beneficial to young people when they engage in hard exercise. However, there have been few papers published about oxidative stress and micronutrient supplementation in healthy children. Although no change of EMF was found in the supplemented children group compared with the control group, the present results showed that healthy children (aged 8–10 years) exhibit higher EMF than those of young and elderly people, which should be attributed to lower oxidative stress or lower accumulation of oxidative damage.

In conclusion, the present findings indicate that the supplementation of multiple antioxidant micronutrients is more effective in decreasing oxidative stress and improving the EMF of young and elderly people than those of children, suggesting that taking these supplements is beneficial to young and elderly people. Moreover, new hypotheses on the physiological causes of ageing and the physiological outcomes of oxidant–antioxidant imbalances, such as those related to membrane fluidity, are expected to be developed.

Acknowledgements

We thank Chinese Natural Science Foundation (no. 30872103) and the Danone Nutrition Institute, Beijing, China, for financial support. X.-X. H. and A.-G. M. designed the intervention study, analysed the data and wrote the manuscript. M. Z., S. G., X.-X. S. and Y.-Y. S. conducted the investigations at field sites, data collection, analyses and interpretation, and laboratory analyses. Q.-Z. W. and H. L. performed a part of the laboratory analyses. There are no conflicts of interest.

References

1. Rice-Evans CA, Diplock AT, Symons MCR, *et al.* (1991) Techniques in free radical research. In *Laboratory Techniques in Biochemistry and Molecular Biology*, pp. 1–18 [RH Burdon and PH Knippenberg, editors]. London: Elsevier.
2. López-Revuelta A, Sánchez-Gallego JI, Hernández-Hernández A, *et al.* (2005) Increase in vulnerability to oxidative damage in cholesterol-modified erythrocytes exposed to *t*-BuOOH. *Biochim Biophys Acta* **1734**, 74–85.
3. Li Y, Qian ZJ, Ryu B, *et al.* (2009) Chemical components and its antioxidant properties *in vitro*: an edible marine brown alga, *Ecklonia cava*. *Bioorg Med Chem* **17**, 1963–1973.

4. Babu GN, Kumar A, Chandra R, *et al.* (2008) Oxidant-antioxidant imbalance in the erythrocytes of sporadic amyotrophic lateral sclerosis patients correlates with the progression of disease. *Neurochem Int* **52**, 1284–1289.
5. El-Rashidy FH, Al-Turk WA & Stohs SJ (1984) Glutathione, glutathione reductase and glutathione *S* transferase activities in erythrocytes and lymphocytes in chronic renal disease. *Res Commun Chem Pathol Pharmacol* **44**, 423–430.
6. Miguel A, Miguel A & Linares M (1988) Evidence of an increased susceptibility to lipid peroxidation in red blood cells of chronic renal failure patients. *Nephron* **50**, 64–65.
7. Southorn PA & Powis G (1988) Free radicals in medicine. II. Involvement in human disease. *Mayo Clin Proc* **63**, 390–408.
8. Goi G, Cazzola R, Tringali C, *et al.* (2005) Erythrocyte membrane alterations during ageing affect beta-D-glucuronidase and neutral sialidase in elderly healthy subjects. *Exp Gerontol* **40**, 219–225.
9. Alpsyoy L, Yildirim A & Agar G (2009) The antioxidant effects of vitamin A, C, and E on aflatoxin B₁-induced oxidative stress in human lymphocytes. *Toxicol Ind Health* **25**, 121–127.
10. Seyedrezazadeh E, Ostadrahimi A, Mahboob S, *et al.* (2008) Effect of vitamin E and selenium supplementation on oxidative stress status in pulmonary tuberculosis patients. *Respirology* **13**, 294–298.
11. Mayne ST (2003) Antioxidant nutrients and chronic disease: use of biomarkers of exposure and oxidative stress status in epidemiologic research. *J Nutr* **133**, 933S–940S.
12. Brosche T & Platt D (1990) Decrease of cholesterol concentration in human erythrocyte ghosts in elderly age. *Exp Gerontol* **25**, 23–28.
13. Prisco D, Rogasi PG, Paniccia R, *et al.* (1991) Age-related changes in red blood lipids. *Angiology* **42**, 316–322.
14. Knight JA (1998) Free radicals: their history and current status in aging and disease. *Ann Clin Lab Sci* **28**, 331–346.
15. Droge W (2003) Oxidative stress and aging. *Adv Exp Med Biol* **543**, 191–200.
16. Davydov VV, Dobaeva NM & Bozhkov AI (2004) Possible role of alteration of aldehyde's scavenger enzymes during aging. *Exp Gerontol* **39**, 11–16.
17. Comstock GW, Norkus EP, Hoffman SC, *et al.* (1995) Stability of ascorbic acid, carotenoids, retinol and tocopherol in plasma stored at -70 degrees C for 4 years. *Cancer Epidemiol Biomarkers Prev* **4**, 505–507.
18. Hess D, Keller HE, Oberlin B, *et al.* (1991) Simultaneous determination of retinol, tocopherols, carotenes and lycopene in plasma by means of high-performance liquid-chromatography on reversed phase. *Int J Vit Nutr Res* **61**, 232–238.
19. Pressman EK, Cavanaugh JL, Mingione M, *et al.* (2003) Effects of maternal antioxidant supplementation on maternal and fetal antioxidant levels: a randomized, double-blind study. *Am J Obstet Gynecol* **189**, 1720–1725.
20. USDHHS, Public Health Services IV (1979) Analytical Methods, Vitamin C. pp. 17–19. Atlanta: Centers for Disease Control and Prevention.
21. Koh T-S & Benson TH (1983) Critical re-appraisal of fluorometric method for determination of selenium in biological materials. *J Assoc Off Anal Chem* **66**, 918–926.
22. Durak I, Canbolat O, Kavutcu M, *et al.* (1996) Activities of total, cytoplasmic and mitochondrial superoxide dismutase enzymes in sera and pleural fluids from patients with lung cancer. *J Clin Lab Anal* **10**, 17–20.
23. Avci A, Atli T, Ergüder IB, *et al.* (2008) Effects of garlic consumption on plasma and erythrocyte antioxidant parameters in elderly subjects. *Gerontology* **54**, 173–176.
24. Steck TL & Kant JA (1974) Preparation of impermeable ghosts and inside out vesicles from human erythrocyte membranes. In *Methods in Enzymology*, pp. 172–173 [SP Colowick and NO Kaplan editors, editors]. London: Academic Press.
25. Raccach D, Dadoun F, Coste T, *et al.* (1996) Decreased Na/K ATPase ouabain binding sites in red blood cells of patients with insulin-dependent diabetes and healthy north African control subjects: relationship with insulin and diabetic neuropathy. *Horm Metab Res* **28**, 128–132.
26. Lowry OH, Rosebrough NJ, Farr AL, *et al.* (1951) Protein measurement with Folin phenol reagent. *J Biol Chem* **193**, 265–275.
27. Chadha VD & Dhawan DK (2009) Membrane fluidity and surface changes during initiation of 1,2-dimethylhydrazine-induced colon carcinogenesis: protection by zinc. *Oncol Res* **18**, 17–23.
28. Aricha B, Fishov I, Cohen Z, *et al.* (2004) Differences in membrane fluidity and fatty acid composition between phenotypic variants of *Streptococcus pneumoniae*. *J Bacteriol* **186**, 4638–4644.
29. Chauhan VP, Chauhan A & Wegiel J (2001) Fibrillar amyloid beta-protein forms a membrane-like hydrophobic domain. *Neuroreport* **12**, 587–590.
30. Tang X, Xia Z & Yu J (2007) An experimental study of hemolysis induced by onion (*Allium cepa*) poisoning in dogs. *J Vet Pharmacol Therap* **31**, 143–149.
31. Ma AG, Schouten EG, Wangm Y, *et al.* (2009) Micronutrient status in anemic and non-anemic Chinese women in the third trimester of pregnancy. *Asia Pac J Clin Nutr* **18**, 41–47.
32. Drewel BT, Giraud DW, Davy SR, *et al.* (2006) Less than adequate vitamin E status observed in a group of preschool boys and girls living in the United States. *J Nutr Biochem* **17**, 132–138.
33. Rizvi SI & Maurya PK (2007) Markers of oxidative stress in erythrocytes during aging in humans. *Ann N Y Acad Sci* **1100**, 373–382.
34. Berzosa C, Gómez-Trullén EM, Piedrafitra E, *et al.* (2011) Erythrocyte membrane fluidity and indices of plasmatic oxidative damage after acute physical exercise in humans. *Eur J Appl Physiol* **111**, 1127–1133.
35. Maurya PK & Rizvi SI (2009) Protective role of tea catechins on erythrocytes subjected to oxidative stress during human aging. *Nat Prod Res* **23**, 1072–1079.
36. Soto-Arriaza MA, Sotomayor CP & Lissi EA (2008) Relationship between lipid peroxidation and rigidity in L-alpha-phosphatidylcholine-DPPC vesicles. *J Colloid Interface Sci* **323**, 70–74.
37. Spengler MI, Bertoluzzo SM, Catalani G, *et al.* (2008) Study on membrane fluidity and erythrocyte aggregation in equine, bovine and human species. *Clin Hemorheol Microcirc* **38**, 171–176.
38. Kaneko T, Matsui H, Shimokawa O, *et al.* (2007) Cellular membrane fluidity measurement by fluorescence polarization in indomethacin-induced gastric cellular injury *in vitro*. *J Gastroenterol* **42**, 939–946.
39. Yang YG (2010) Population and environment sustainable development in rural China. *J Shandong Inst Min Technol (Social Science)* **3**, 55–56.
40. Ma AG, Schouten EG, Zhang FZ, *et al.* (2008) Retinol and riboflavin supplementation decreases the prevalence of anemia in Chinese pregnant women taking iron and folic acid supplements. *J Nutr* **138**, 1946–1950.

41. Rizvi SI, Pandey KB, Jha R, *et al.* (2009) Ascorbate recycling by erythrocytes during aging in humans. *Rejuvenation Res* **12**, 3–6.
42. Clemens MR & Waller HD (1987) Lipid peroxidation in erythrocytes. *Chem Phys Lipids* **45**, 251–268.
43. Oh HY, Lim S, Lee JM, *et al.* (2007) A combination of soy isoflavone supplementation and exercise improves lipid profiles and protects antioxidant defense-systems against exercise-induced oxidative stress in ovariectomized rats. *Biofactors* **29**, 175–185.
44. Harris DE (1992) Regulation of antioxidant enzymes. *FASEB J* **6**, 2675–2683.
45. Liu MJ, Li JX, Guo HZ, *et al.* (2003) The effects of verbasco-side on plasma lipid peroxidation level and erythrocyte membrane fluidity during immobilization in rabbits: a time course study. *Life Sci* **73**, 883–892.
46. Erden-Inal M, Sunal E & Kanbak G (2002) Age-related changes in the glutathione redox system. *Cell Biochem Funct* **20**, 61–66.
47. Ruiz-Ramos M, Vargas LA, Fortoul Van der Goes TI, *et al.* (2010) Supplementation of ascorbic acid and alpha-tocopherol is useful to preventing bone loss linked to oxidative stress in elderly. *J Nutr Health Aging* **14**, 467–472.
48. McCance DR, Holmes VA, Maresh MJ, *et al.* (2010) Vitamins C and E for prevention of pre-eclampsia in women with type 1 diabetes (DAPIT): a randomised placebo-controlled trial. *Lancet* **376**, 259–266.
49. Hopkins MH, Fedirko V, Jones DP, *et al.* (2010) Antioxidant micronutrients and biomarkers of oxidative stress and inflammation in colorectal adenoma patients: results from a randomized, controlled clinical trial. *Cancer Epidemiol Biomarkers Prev* **19**, 850–858.
50. Ferrer MD, Tauler P, Sureda A, *et al.* (2010) Variegated porphyria induces plasma and neutrophil oxidative stress: effects of dietary supplementation with vitamins E and C. *Br J Nutr* **103**, 69–76.
51. Mydlik M & Derzsiová K (2008) Vitamins and quality of life in hemodialysis patients. *J Nephrol* **21**, S129–S133.
52. Sangeetha P, Balu M, Haripriya D, *et al.* (2005) Age associated changes in erythrocyte membrane surface charge: modulatory role of grape seed proanthocyanidins. *Exp Gerontol* **40**, 820–828.
53. Brzeszczynska J, Pieniazek A, Gwozdziński L, *et al.* (2008) Structural alterations of erythrocyte membrane components induced by exhaustive exercise. *Appl Physiol Nutr Metab* **33**, 1223–1231.