

P. Graham. *Institute of Child Health, London, UK*

Ethical issues around the use of integrative approaches in child and adolescent psychiatry arise in relation to diverse aspects of practice but especially in collaborative work with other disciplines. Child and adolescent psychiatrists work with nurses, social workers, psychologists, psychotherapists, family practitioners, paediatricians, those working in the juvenile justice system and educationists. Different ethical issues arise in relation to work with different disciplines and examples will be given. But there are common ethical issues arising from different standards of confidentiality and the communication of information as well as in the exercise of medical responsibility. Ethical issues can sometimes hinder the delivery of effective, evidence-based care. This is paradoxical for failure to deliver such care itself offends against ethical principles.

### PS01.02

Collaboration between industrialized and developing countries in psychiatry

D. Moussaoui. *Ibn Rushd Niversity Psychiatric Centre, Casablanca, Morocco*

There is no possibility of progress for psychiatry in the world if there is a too wide gap between the quality of psychiatric practice and research in developing and in industrialized countries. The reason for that is because bad image which might emerge from developing countries, in a media globalized world, impacts negatively on the mental health work all over the world. Examples will be given for such statement.

On the other hand, collaboration existed for a long period of time between these two categories of countries in the education of psychiatrists. It is necessary, but at the same time creates major problems, such as the brain drain which is a direct consequence of learning psychiatry abroad for students coming from developing countries. There are more psychiatrists from Pakistan, India or South Africa practicing in the UK or the USA than in their country of origin. Another problem is the lack of transcultural sensitivity, making the training of these students insufficient when they come back home.

On the other hand, collaboration in research must take into account the cultural specificities of the developing country, and in anyway should avoid a “safari research”. Respect of ethical guidelines in all collaborative studies is an essential ingredient of success.

### PS01.03

Ethical questions in integrating standard therapies and alternative therapies in psychiatry

H. Helmchen. *Klinik Für Psychiatrie Und Psychotherapie, Charite, Berlin, Germany*

Increasing predominance of evidence-based therapies and standardisation of their clinical application as well as financial limitations narrow the therapists view and leaving out of account the patients' individual specificities and demands. Consequently patients try to get their subjective needs met by alternative approaches, such as self-medication, procedures of unknown quality, consultation of healers outside the medical professions. According to the prevailing ethical principles the following ethical questions will be discussed:

- Respect of the patients' dignity and autonomy means to take him/her seriously. However, what are the limits of taking the view of the patient, particularly in cases of a discrepancy between the

patients will and his welfare, and especially in cases of incapacity to decide competently on therapeutic alternatives?

- Harm avoidance should keep away from the patient unproven approaches with both the risks of unknown unwanted effects and of omission of an efficient treatment. But how to sail safe through the narrowness of *Skylla and Charybdis*: to convince the patient of the advantages of the proposed quality proven approach without chasing him away out of medical services to unqualified and uncontrolled approaches?
- With regard to justice it seems clear that approaches without proven quality should not be paid by insurance companies in order not to reduce the limited resources for the whole of all its members. However, will this be acceptable also in cases in which there are no efficient treatments, but the alternative approach demanded by the patient may improve his quality of life?

---

## Symposium: ADHD across the lifespan: Genetics, learning, comorbidity and circadian rhythm

---

### S21.01

Attention deficit/hyperactivity disorder and dyslexia: evidence for shared genetic susceptibility

C.L. Barr<sup>1,2</sup>, K. Wigg<sup>1</sup>, N. Laurin<sup>1</sup>, V. Misener<sup>2</sup>, T. Cate-Carter<sup>2</sup>, L.W. Maureen<sup>2</sup>, E. Kerr<sup>2</sup>, J. Couto<sup>1</sup>. <sup>1</sup> *Genetics and Development Division, The Toronto Western Hospital, Toronto, ON, Canada* <sup>2</sup> *Program in Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, ON, Canada*

**Objective:** Many individuals ascertained for developmental dyslexia (DD) are also diagnosed with attention-deficit hyperactivity disorder (ADHD) and approximately 20% of individuals with ADHD will have evidence for DD. The basis for this overlap is not completely understood but twin studies have provide support for common genetic influences, particularly for inattention symptoms. Genetic linkage studies have found significant evidence for linkage of DD to chromosomes 1p34-p36, 15q, 6p21.3-22, 2p15-16, 6q11.2-q12 and 18p11.2. Evidence for linkage/association to ADHD has also been found to overlap for some of these regions. The objective of this study is to identify genes contributing to both.

**Methods:** We examined evidence for the involvement of specific genes in these chromosomal regions using two samples of families, one ascertained through a proband with DD (n= 273 families) and the other through a proband with ADHD (n= 390 families).

**Results:** Our studies of the 6p region indicate that the sample of DD families is associated to markers in this region and to ADHD but not to the same markers within the linked region. For the 15q region, we have found significant evidence for association for both the ADHD and reading phenotypes in both samples (Wigg et al., 2004; Wigg et al., 2005). We have also found evidence for the gene for the dopamine receptor D1 to be associated to the inattention symptoms in both samples

**Conclusions:** While the studies of the overlap in ADHD and DD are preliminary, they are promising in that they will ultimately help to disentangle the causal relationship.