

Sensory and cognitive deficits in schizophrenia may be the result of top-down regulation failure.

Disclosure: No significant relationships.

Keywords: schizophrenia; visual and olfactory impairments; cognition functions

O262

Self - evaluation of social adaptation in patients with schizophrenia and metabolic syndrome

V. Khamina^{1,2,*}, E. Kornetova¹, A. Kornetov², A. Goncharova¹ and A. Boiko¹

¹Mental Health Research Institute, Tomsk National Research Medical Center of Russian Academy of Sciences, Tomsk, Russian Federation and ²Department Of Fundamental Psychology And Behavioral Medicine, Siberian State Medical University, Tomsk, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.435

Introduction: Metabolic syndrome (MS) is an often co-occurring condition that occurs during antipsychotic therapy and impairs social functioning

Objectives: We tried to conduct a self - evaluation of social adaptation in patients with schizophrenia and MS

Methods: We examined 150 patients with schizophrenia receiving antipsychotic therapy at the clinics of Mental Health Research Institute. The study was supported by a grant from the Russian Science Foundation 18-15-00011. The IDF criteria were used to diagnose metabolic syndrome. We used «The social adaptation self - evaluation scale» (SASS).

Results: 63 patients (42%) had MS and 87 patients (58%) did not. In the subgroup of patients with MS, 59 people (93.65%) had disabilities or were unemployed, in the group without MS - 82 (94.26%) patients. There were no statistically significant differences between the groups ($p \geq 0.05$). In the patients with schizophrenia and concomitant MS, the median SSAS scores was 35 [29; 39], which corresponds mainly to a high level of self - evaluation of social adaptation. At the same time, in patients with schizophrenia and without MS, on the contrary, the self - evaluation of social adaptation was 30 [23; 38] points ($p = 0.03914$). Perhaps this is due to the great attention from relatives and doctors of general somatic practice and the primary medical network in connection with the risk of developing severe somatic pathology.

Conclusions: Patients with MS can give a higher assessment of social adaptation, despite a objectively low social status.

Disclosure: No significant relationships.

Keywords: social adaptation; schizophrenia; Metabolic syndrome

O263

Social cognition as a mediator between cognitive reserve and psychosocial functioning in patients with first episode psychosis

I. González-Ortega^{1*}, S. Alberich-Mesa², E. Echeburúa^{2,3}, M. Bernardo⁴, B. Cabrera⁴, S. Amoretti⁴, A. Lobo⁵, C. Arango⁶, I. Corripio⁷, E. Vieta⁸, E. De La Serna⁹, R. Rodríguez-Jiménez¹⁰,

R. Segarra¹¹, J.M. López-Ilundain¹², A. Sánchez-Torres¹², M. Cuesta¹² and A. González-Pinto¹

¹Department Of Psychiatry, Araba University Hospital, Araba University Hospital, BIOARABA RESEARCH INSTITUTE, University OF THE BASQUE COUNTRY, CIBERSAM, Vitoria-Gasteiz, Spain; ²Departament Of Psychiatry, Araba University Hospital, Araba University Hospital, BIOARABA RESEARCH INSTITUTE, University OF THE BASQUE COUNTRY, CIBERSAM, Vitoria-Gasteiz, Spain; ³Psychology, UNIVERSITY OF THE BASQUE COUNTRY, BIODONOSTIA, CIBERSAM, San Sebastián, Spain; ⁴Psychiatry, Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, CIBERSAM, Barcelona, Spain; ⁵Medicine And Psychiatry, University of Zaragoza, Aragon Institute for Health Sciences (IIS Aragón), Zaragoza, Spain; ⁶Child And Adolescent Department. Institute Of Psychiatry And Mental Health, Hospital General Universitario Gregorio Marañón. CIBERSAM, Madrid, Spain; ⁷Psychiatry, Institut d'Investigació Biomèdica-Sant Pau (IIB-SANT PAU), Barcelona, Spain; ⁸Institute Of Neuroscience, Hospital Clinic, University Of Barcelona, Bipolar and Depressive Disorders Unit, IDIBAPS, CIBERSAM, Barcelona, Spain; ⁹Child And Adolescent Psychiatry, Hospital Clinic of Barcelona, CIBERSAM, Barcelona, Spain; ¹⁰Psychiatry, 12 de Octubre Hospital Research Institute, CIBERSAM, Madrid, Spain; ¹¹Neurosciences, University of the Basque Country, Cruces University Hospital, Biocruces Bizkaia Health Research Institute, CIBERSAM, Leioa, Spain and ¹²Psychiatry, Navarre Hospital Complex, IdiSNA, Navarre Institute for Health Research, Pamplona, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.436

Introduction: Social cognition has been associated with functional outcome in patients with first episode psychosis (FEP). Social cognition has also been associated with neurocognition and cognitive reserve. Although cognitive reserve, neurocognitive functioning, social cognition, and functional outcome are related, the direction of their associations is not clear.

Objectives: The aim of the study was to analyze the influence of social cognition as a mediator between cognitive reserve and cognitive domains on functioning in FEP both at baseline and at 2 years.

Methods: The sample of the study was composed of 282 FEP patients followed up for 2 years. To analyze whether social cognition mediates the influence of cognitive reserve and cognitive domains on functioning, a path analysis was performed. The statistical significance of any mediation effects was evaluated by bootstrap analysis.

Results: At baseline, as neither cognitive reserve nor the cognitive domains studied were related to functioning, the conditions for mediation were not satisfied. Nevertheless, at 2 years of follow-up, social cognition acted as a mediator between cognitive reserve and functioning. Likewise, social cognition was a mediator between verbal memory and functional outcome. The results of the bootstrap analysis confirmed these significant mediations (95% bootstrapped CI (-10.215 to -0.337) and (-4.731 to -0.605) respectively).

Conclusions: Cognitive reserve and neurocognition are related to functioning, and social cognition mediates in this relationship.

Disclosure: This work was supported by the Carlos III Institute of Health and European Fund for Regional Development (PI08/1213, PI11/ 01977, PI14/01900, PI08/01026, PI11/02831, PI14/01621, PI08/1161, PI16/ 00359, PI16/01164, PI18/00805), the Basque Foundation for Health

Keywords: first episode psychosis; social cognition; cognitive reserve; functioning